

**Supplementary table:** Sexual and contraceptive use history of BioCycle study participants by sexual activity category

	Sexual Activity Category					<i>p</i> -value <sup>a</sup>
	Total Cohort	Sexual inactive	Not sexually active during study	Weekly or less	Greater than weekly	
n (%)	258	59 (23)	64 (25)	71 (28)	64 (25)	-----
<b>Sexual history</b>						
Ever sexually active						
Yes	192 (74)	0 (0)	64 (100)	65 (92)	63 (98)	<0.01
No	63 (24)	59 (100)	0 (0)	4 (6)	0 (0)	
Missing	3 (1)			2 (3)	1 (2)	
Currently sexually active <sup>b</sup>						
Yes	135 (70)	NA	17 (27)	57 (88)	61 (97)	<0.01
No	57 (30)	NA	47 (73)	8 (12)	2 (3)	
Sexual orientation <sup>b</sup>						
Heterosexual	180 (94)	NA	58 (91)	63 (97)	59 (94)	0.66
Lesbian	7 (4)	NA	4 (6)	1 (2)	2 (3)	
Bisexual	5 (3)	NA	2 (3)	1 (2)	2 (3)	
Ever used contraceptives <sup>b</sup>						
Yes	177 (92)	NA	55 (86)	62 (95)	60 (95)	0.06
No	14 (7)	NA	9 (14)	3 (5)	2 (3)	
Missing	1 (1)	NA	0 (0)	0 (0)	1 (2)	
<b>Current contraceptive method<sup>b,c</sup></b>						
Any male condom use	68 (38)	NA	10 (18)	29 (47)	29 (48)	<0.01
Male condom use only	47	NA	5	21	21	<0.01
Male condom use plus other methods	21	NA	5	8	8	0.80
Other methods only	17 (10)	NA	1 (2)	6 (10)	10 (17)	0.02
No/unknown contraceptive method	92 (52)	NA	44 (80)	27 (44)	21 (35)	<0.01

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

<sup>a</sup> *P*-values calculated using Fisher's exact test.

<sup>b</sup> Only women who reported ever being sexually active were asked this question.

<sup>c</sup> Only women who reported ever using contraceptives were asked about current contraceptive method.