	Participants who tested positive for rapid HCV testing n = 7										
	Subject 1	Subject 2	Subject 3	<u>Subject 4</u>	Subject 5	<u>Subject 6*</u>	Subject 7*				
Age	25	29	32	34	41	44	51				
Sex	Male	Male	Male	Male	Male	Female	Male				
Source of medical care	ED	Private clinic/practice	Hospital or community health clinics	ED	ED	Hospital or community health clinics	Urgent care center				
Study arm	No intervention	No intervention	Intervention	No intervention	Intervention	Intervention	Intervention				
HIV testing history											
Ever tested and how	Never donated blood or tested for HIV	Donated blood but never tested for HIV	Tested but not part of blood donations	Donated blood but never tested for HIV							
Time elapsed since last test	N/A	N/A	≤ 2 years but > 1 year ago	≤ 5 years but > 2 years ago	≤ 6 months ago	> 5 years ago	N/A				
Reason for last HIV test	N/A	N/A	Blood or blood product transfusion	Jail/prison	Jail/prison	Jail/prison	N/A				
Reason for not testing for HIV	Hadn't considered it	Never been asked or offered a test	N/A	N/A	N/A	N/A	Never been asked or offered a test				
HCV testing history											
Ever tested and how	Never donated blood or tested for HCV	Donated blood but never tested for HCV	Tested but not part of blood donations	Donated blood but never tested for HCV							
Time elapsed since last test	N/A	N/A	≤ 2 years but > 1 year ago	\leq 5 years but > 2 years	≤ 6 months ago	> 5 years ago	N/A				
Reason for last HCV test	N/A	N/A	Blood or blood product transfusion	Jail/prison	Jail/prison	Jail/prison	N/A				
Reason for not testing for HCV	Hadn't considered it	Never been asked or offered a test	N/A	N/A	N/A	N/A	Never been asked or offered a test				
Past 3 months drug use ASSIST category	Marijuana, Cocaine or crack, Methamphetamines, Hallucinogens, Illicit opioids, Benzodizepines, Methadone or	Illicit opioids, Benzodiazepines	Marijuana	Marijuana	Marijuana	Cocaine or crack, Prescription opioid analgesics	Marijuana, Cocaine or crack, Illicit opiods				

Data Supplement 10: Demographic, clinical, and other characteristics and risk-taking behaviors of InVITED participants with a reactive HCV antibody test

	buprenorphine, Prescription opioid analgesics						
Injection Drug Use	Yes, past 3 months	Never	Never	Never	Yes but not in the past 3 months	Never	Yes, past 3 months
Shared needles	Yes	No	No	No	No	No	Yes
Total drug ASSIST score	135	45	29	16	18	55	64
WHO recommended drug intervention	Intensive intervention	Brief intervention	Brief intervention	Brief intervention	Brief intervention	Intensive intervention	Brief intervention
Sexual risk-taking behaviors Type of partners	Main, casual, exchange	Main	Main, casual	None	Main, casual	Main, exchange	Don't know
Had anal sex with men	No	No	No	No	No	N/A	No
Did not use condoms	Yes	Yes	Yes	N/A	Yes	Used no condoms with main partners, but used condoms with exchange partners	N/A
Partners had HIV or AIDS	Don't know	No	Don't know	N/A	No	No	N/A
Partners had other STD	Don't know	Don't know	Don't know	N/A	No	Yes	N/A
Partners ever injected drugs	Don't know	Don't know	Don't know	N/A	Yes	Yes	N/A
No condoms use with partners	20 main, 20 casual,	5 main	8 main, 13 casual	N/A	20 main, 25	6 main	N/A

AIDS = Acquired Immune Deficiency Syndrome; ASSIST = Alcohol, Smoking and Substance Involvement Screening Test; ED = Emergency Department; HCV = Hepatitis C Virus; N/A = Not Applicable; STD = Sexual Transmitted Disease; WHO = World Health Organization