

SUPPLEMENTARY TABLE S4. DOSAGE IN SIX SUBJECTS, WHO DIED DURING THE STUDY PERIOD,
IN THE 12 MONTHS PERIOD PRIOR TO DEATH

| | <i>Highest daily dose ever after the age of 5</i> | <i>Number of months prior to death</i> | | | | | | | | | | | <i>At the time of death</i> | |
|---------|---|--|-----------|-----------|----------|----------|----------|----------|----------|----------|----------|----------|---------------------------------|----------|
| | | <i>12</i> | <i>11</i> | <i>10</i> | <i>9</i> | <i>8</i> | <i>7</i> | <i>6</i> | <i>5</i> | <i>4</i> | <i>3</i> | <i>2</i> | | <i>1</i> |
| Child 1 | 15 mg | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| Child 2 | 30 mg | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | NU |
| Child 3 | 30 mg | NU | NU | NU | NU | 1 | 3 | 3 | 2 | 3 | 2 | 2 | 3 | 3 |
| Child 4 | 36 mg | NU | NU | 1 | 1 | 1 | NU | NU | NU | NU | NU | NU | NU | NU |
| Child 5 | 54 mg | NU | NU | NU | NU | NU | NU | NU | NU | NU | NU | NU | NU | NU |
| Child 6 | 54 mg | 1 | 1 | 1 | 1 | 2 | 2 | 1 | 1 | 1 | 1 | 2 | 2 | 2 |

NU corresponds to non-use

1 corresponds to low dose (<15 mg per day)

2 corresponds to medium dose (15–30 mg per day)

3 corresponds to high dose (>30 mg per day)