

Supporting Information

for *Macromol. Biosci.*, DOI: 10.1002/mabi.201200384

An anti-angiogenic reverse thermal gel as a drug delivery system for age-related wet macular degeneration

Daewon Park, Veeral Shah, Britta M. Rauck, Thomas R. Friberg, Yadong Wang*

Dr. D. Park
Department of Bioengineering, University of Colorado Denver Anschutz Medical Campus
Aurora, CO 80045, USA

Dr. V. Shah, Dr. T.R. Friberg
Department of Ophthalmology, University of Pittsburgh School of Medicine
Pittsburgh, PA 15219, USA

B.M. Rauck, Prof. Y. Wang
Department of Bioengineering, Chemical Engineering, Surgery and the McGowan Institute
for Regenerative Medicine, University of Pittsburgh
Pittsburgh, PA 15219, USA
E-mail: yaw20@pitt.edu




Figure S1. The bevacizumab standard curve, concentration vs absorption, measured between 0.2 ng/ml - 3.125 µg/ml. For the calculation, the linear region between 12.2 ng/ml - 3.125 µg/ml was used.




Figure S2. *In vitro* cytotoxicity studied using primary bovine corneal endothelium cells. Phase images (100 \times) of Calcein AM-treated cells cultured in (A) control TCPS and (B) ESHU conditions. No differences were observed between control and ESHU-exposed cells.