

Table S1. Characteristics of patients with and without testosterone measurement

Factor	With testosterone measurement (N=2419)	Without testosterone measurement (N=23281)
Age*	67.3±11.3	72.1±11.5
African American	268(11.1)	2293(9.8)
CKDEPI eGFR at Inception*	49.1±9.5	47.0±10.5
eGFR groups *		
45-59	1776(73.4)	15106(64.9)
30-44	511(21.1)	6134(26.3)
15-29	132(5.5)	2041(8.8)
BMI *	29.4±5.4	28.9±5.5
BMI group *		
<18.5 kg/m2	10(0.41)	169(0.73)
18.5-24.9 kg/m2	457(18.9)	4969(21.3)
25-29.9 kg/m2	997(41.2)	9232(39.7)
30+ kg/m2	931(38.5)	7781(33.4)
Missing	24(0.99)	1130(4.9)
Smoking Status*		
No	2123(87.8)	17503(75.2)
Yes	176(7.3)	1744(7.5)
Missing	120(5.0)	4034(17.3)
Diabetes *	744(30.8)	5219(22.4)
Hypertension *	2243(92.7)	19100(82.0)
Cerebrovascular disease	252(10.4)	2301(9.9)
Malignancy	673(27.8)	6340(27.2)
CAD	714(29.5)	6755(29.0)
CHF *	330(13.6)	2321(10.0)
Hyperlipidemia *	2055(85.0)	17707(76.1)

CKD EPI: chronic kidney disease epidemiology collaboration equation, BMI: body mass index, CAD: coronary artery disease, CHF: chronic heart failure

Values presented as Mean ± SD with t-test; or N(%) with chi-square test unless otherwise noted.

*Significantly different between patients with and without testosterone data