

Registration or ID Number	
----------------------------------	--

Age	Gender

Mode of Transport: How was injured person traveling?

Pedestrian	Non-motorized vehicle (bike, cart)	Motorcycle
Car	Pick-up, minibus, jeepney	Truck
Bus	Train/subway	Unknown
Other – specify		
All applicable codes:		

Type of Road User: What was the injured person doing?

Pedestrian	Driver	Passenger
Unknown	Other – specify:	
All applicable codes:		

Injuries Sustained

Fracture	Sprain/Strain	Cut or open wounds
Contusion/bruising	Burn	Concussion/head trauma
Internal injury	Broken bones	Unknown
Other – specify		
All applicable codes:		

Injury Type

Unintentional (direct result of traffic incident)	Intentional (deliberately inflicted on one person by another)	Medical case only	Medico-legal case
Other – specify			
All applicable codes:			

Severity of Injuries

Mild injuries (small cuts or lacerations, bruising)	Moderate injuries (sprain/strain, deep cuts or open wounds, mild burns)	Severe injuries (fractures, dislocations severe contusions, moderate burns)
Very severe injuries (concussion/loss of consciousness, broken bones, internal injury/bleeding, severe burns, amputation, paralysis)	Unknown	Other –specify
Other – specify:		
All applicable codes		

Injury Severity Score

Injury	Abbreviated Injury Scale (AIS)
Minor	1
Moderate	2
Serious	3
Severe	4
Critical	5
Unsurvivable	6

Region	Injury Description	AIS	ISS
Head and Neck			
Verbal response			
Motor response			
Total ISS (top three squared and added)			

Assessment of Impaired Consciousness (GCS Score)

Mild head injury	GCS of 13-15
Moderate head injury	GCS of 9-12
Severe head injury	GCS of 8 or less

Response	Condition	Score
Eye opening	Spontaneously	4
	To speech	3
	To pain	2
	None	1
Verbal response	Orientated	5
	Confused	4
	Inappropriate	3
	Incomprehensible	2
	None	1
Motor response	Obeys commands	6
	Localizes to pain	5
	Withdraws from pain	4
	Flexion to pain	3
	Extension to pain	2
	None	1
	Total GCS	

Alcohol Use

Patient presented clinical signs of intoxication	
--	--

If Alcohol (1) then complete the following:

Assessment of Intoxication

Mild intoxication (50mg/dL): Smell of alcohol on breath, Exuberance/loss of emotional restraint/depressed mood, impairment of attention and/or judgment, conjunctival injection and/or flushed face	Moderate intoxication (100mg/dL): All or any of mild intoxication symptoms PLUS impairment of speech (e.g. slurring), impairment of motor coordination, disturbances in behavioural response, disturbances in emotional responses, impaired ability to cooperate	Severe intoxication (200mg/dL): All or any of mild/moderate intoxication symptoms PLUS horizontal gaze, nystagmus
Very severe intoxication (300mg/dL): All or any of mild/moderate/severe intoxication symptoms PLUS stuporous, heavy sleeping/heavy snoring	Extremely severe intoxication (400mg/dL): All or any of mild/moderate/severe/very severe intoxication symptoms PLUS irregular breathing, inhalation of vomit, comatose/inability to be aroused	Unknown
Other – specify:		

All applicable codes:

Patient Assessment

Status of intake and medical history	
Status of blood pressure	
Non-invasive procedures performed	
Laboratory testing for BAC performed	

If lab testing (1), then

Did blood test determine presence of alcohol?	
---	--

Results

Patient treated and sent home	Patient admitted to hospital, no surgery required	Patient admitted to hospital, surgery required
Patient referred to a specialist or another facility	Unknown	
Other – specify:		
All applicable codes:		