

Supplementary Table 3. Oligonucleotides used for the construction of expression vectors.

(Sequences are given 5' → 3')

name	sequence
oh-BglII TEVBEAFFw	AGCGCGAGATCTGAAAACCTGTATTTTCAGGGAATGCC
oh-HindBEAFRev	GGCCGCAAGCTTCAGTGATGATGATGATGATGTCCCTGAAAATACAGG T
TEVBEAFFw	TATTTTCAGGGAATGCCCAAGGGTCGTGTTA
TEVBEAFRev	CTGAAAATACAGGTTTTCTCATCCTTGCAAGCGT
ChromatorattB1TEFW	GGGACAAGTTTGTACAAAAAAGCAGGCTTCGAAAACCTGTATTTTCA GGGCATGTTGGCACAGGAGATTTACC
ChromatorattB2Rev	GGGACCACTTTGTACAAGAAAGCTGGGTGTTACGTTGGGATGTTGA GCGTCTTA
CP190attB1 Fw	GGGACAAGTTTGTACAAAAAAGCAGGCTTCGAAAACCTGTATTTTCA GGGCATGGGTGAAGTCAAGTCCGTG
CP190attB2Rev	GGGACCACTTTGTACAAGAAAGCTGGGTGTTATAGCTCCTCCTTCGC CGCCG
chrom-C-601-926attB1	GGGACAAGTTTGTACAAAAAAGCAGGCTTCGAAAACCTGTATTTTCA GGGCGGAACCTCCGCAACATCGCCAAC
chrom-C-attB1	GGGACAAGTTTGTACAAAAAAGCAGGCTTCATCGATGAAACGGACG AAG