

**Extended metAFLP approach in studies of the tissue culture induced variation (TCIV)
in case of triticale**

Molecular Breeding

Joanna Machczyńska¹, Renata Orłowska¹, Janusz Zimny², Piotr Tomasz Bednarek*¹

¹Department of Plant Physiology and Biochemistry

²Department of Plant Biotechnology and Cytogenetics

Plant Breeding and Acclimatization Institute - National Research Institute, 05-870 Błonie,
Radzików, Poland

*Corresponding author: Piotr Tomasz Bednarek - p.bednarek@ihar.edu.pl

Online Resource 7 Arrangement of events represented by individual 4-digit codes for regenerants derived via anther (RA), shed-microspore (RM), immature zygotic embryo cultures (RE) and for all regenerants taken together (Σ).

4-digit code	RA	RM	RE	Σ
0000	1046	1225	1106	3377
0001	182	204	165	551
0010	266	344	316	926
0011	429	563	349	1341
0100	119	199	73	391
0101	228	262	294	784
0110	66	92	31	189
0111	539	561	656	1756

1000	192	220	238	650
1001	43	83	50	176
1010	151	238	167	556
1011	339	491	378	1208
1100	168	217	117	502
1101	272	290	297	859
1110	275	350	298	923
1111	10285	12181	10649	33115
