

Additional file 1: Pre-treatment scores of patients, community norms, Cronbach's Alpha and criterion C for each outcome measure

	N	M	SD	M	SD	Cron-	Crite-	%
	Patients	Patients	Patients	Norm	Norm	bach's α	riterion C	Norma-
						Patients		tive
								patients
EDI-2								
Drive for Thinness	434	29.60	8.65	17.3	6.8	.89	22.71	21.8
Bulimia	434	16.73	9.26	10.6	3.4	.92	12.25	41.7
Body Dissatisfaction	429	38.21	9.40	30.2	10.3	.85	34.39	37.0
Ineffectiveness	429	37.33	10.78	23.5	5.7	.91	28.28	21.3
Perfectionism	434	23.17	6.27	16.5	5.7	.77	19.68	25.1
Interpersonal Mistrust	432	24.01	6.89	18.4	4.4	.84	20.59	29.6
Interoceptive Awareness	434	36.25	9.63	22	5.7	.84	27.30	16.9
Maturity Fears	431	27.78	8.04	20.8	4.7	.83	23.37	26.9
Asceticism	429	26.23	7.51	16.5	4	.78	19.88	17.3
Impulse regulation	427	28.75	8.10	23.6	6.3	.77	25.85	36.4
Social Insecurity	433	28.52	6.86	22	4.7	.78	24.65	26.4
EDI-2 Global score	433	315.96	63.86	223.2	38.1	.96	257.86	20.6
BSI GSI	431	1.37	0.72	0.35	0.23	.96	0.60	15.7
BDI-2	381	28.37	12.22	7.4	7.3	.92	15.24	13.9

Notes: EDI-2=Eating disorder inventory-2, BSI=Brief symptom inventory, GSI=Global severity index, BDI-2=Beck depression inventory-2, norms for the healthy samples were taken from the applicable norms [24, 26, 28]