

Table S1 Sampling localities for all wild-derived inbred laboratory strains used in this study.

Subspecies	Wild Derived Inbred		
	Line ID	Country	Locality
<i>M. m. castaneus</i>	CAST/EIJ ^a	Thailand	Thonburi
	CIM/MPL	India	Masinagudi
	CKN/MPL	Kenya	Nairobi
	CKS/MPL	Kenya	Shanzu
	CTP/MPL ^b	Thailand	Pathumthani
	DKN/MPL	Kenya	Nairobi
	MDG/MPL	Madagascar	Manakasina
	MPR/MPL ^b	Pakistan	Rawalpindi
<i>M. m. domesticus</i>	BIK/MPL	Israel	Kefar Galim
	BZO/MPL	Algeria	Oran
	DCP/MPL	Cyprus	Paphos
	DJO/MPL	Italy	Orcetto
	DMZ/MPL	Morocco	Azemmour
	LEWES/EIJ	USA	Delaware
	WLA/MPL	France	Toulouse
	WSB/EIJ ^a	USA	Maryland
<i>M. m. musculus</i>	BID/MPL ^b	Iran	Birdjand
	CZECHII/EIJ	Czechoslovakia	
	MBK/MPL	Bulgaria	Kranevo
	MBT/MPL	Bulgaria	Général Toshevo
	MCZ/MPL	Czech Republic	Bialowieza
	MDH/MPL	Denmark	Hov
	MPB/MPL	Poland	Prague
	PWK/PhJ ^c	Czech Republic	Lhotka
<i>M. caroli</i>	CAROLI/EIJ	Thailand	
<i>M. spretus</i>	SPRET/EIJ ^c	Spain	Cadiz

^aData were taken from the Wellcome Trust Mouse Genomes Project.

^bData were excluded from further analyses due to admixture.

^cData from transcriptome sequencing was combined with data from the Wellcome Trust Mouse Genomes Project.