

Table S12 Average measures of differentiation in regions containing testis specific genes and all other regions for all pairwise comparisons of *Mus musculus* subspecies.

Subspecies	Subspecies	Regions	n	\bar{F}_{st} (SD)	t	\bar{D}_{xy} (SD)	t	$\bar{\delta}$ (SD)	t
<i>M. m. castaneus</i>	<i>M. m. domesticus</i>	Contain testis specific genes	520	0.24 (0.16)	0.41	0.42 (0.14)	0.29	0.41 (0.14)	0.16
		All Others	1226	0.24 (0.19)		0.42 (0.16)		0.41 (0.16)	
<i>M. m. castaneus</i>	<i>M. m. musculus</i>	Contain testis specific genes	515	0.29 (0.19)	1.73*	0.47 (0.15)	1.47	0.46 (0.16)	1.64*
		All Others	1247	0.27 (0.20)		0.45 (0.16)		0.44 (0.17)	
<i>M. m. domesticus</i>	<i>M. m. musculus</i>	Contain testis specific genes	518	0.43 (0.22)	2.15*	0.56 (0.18)	1.81*	0.56 (0.18)	1.93*
		All Others	1364	0.41 (0.24)		0.55 (0.19)		0.54 (0.20)	

*P<=0.05 in 1-sided t-tests comparing measures from regions containing testis specific regions and all others.