

1. Location of your practice:

Town

County

2. Roughly how many cases of bovine babesiosis (redwater) did you attend in the last 12 months?

3. At what time of year did they mostly occur?

- Mainly in the spring
- Mainly in the summer
- Mainly in the autumn
- Mainly in the winter
- At any time of year

Other (please specify)

4. Approx how many of these cases:

a) Needed to be revisited?

b) Required blood transfusions?

c) Resulted in mortalities?

COMMENTS

5. What were the chief presenting clinical signs?

	most often	sometimes	rarely	never
Haemoglobinuria (redwater)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Fever	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Jaundice	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Diarrhoea	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Constipation	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Rapid breathing (tachypnea)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Increased heart rate	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Edema of the brisket	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Other (please specify)

6. Do you think there has there been a change in the number of clinical babesiosis cases in the last 10 years?

- Yes, they have increased
- Yes, they have decreased
- No, they have not changed
- Don't know

Comment

7. Do you think there has there been a change in the severity of bovine babesiosis cases in the last 10 years?

- Yes, infections tend to be more severe
- Yes, infections tend to be less severe
- No, the severity has not changed
- Don't know

Comment

8. How are most cases of bovine babesiosis diagnosed? Please rank according to your experience

	most often	sometimes	rarely	never
Case history	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Previous history of babesiosis on the farm	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Treatment success	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Blood smear (in-house)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Referral to a regional laboratory	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Other (please specify)

9. How effective is treatment of bovine babesiosis with imidocarb dipropionate (Imizol)?

- Highly effective
- Effective
- Not effective
- Not at all effective
- Don't know

Comment

10. Do you use Imizol prophylactically?

- Yes
- No

11. If yes, how effective is the drug as a prophylactic agent?

- Highly effective
- Effective
- Not effective
- Not at all effective

12. Do you use any other drugs as prophylactic agents against bovine babesiosis? If yes, please specify:

13. Would you be amenable to receive an email seeking further information? If yes, please provide your name and contact details:

Name

Email/Telephone



Many thanks for taking the time to complete this survey!