

Supporting Material

Computational Studies of the Effect of the S23D/S24D Troponin I Mutation on Cardiac Troponin Structural Dynamics

Yuanhua Cheng^{1,2}, Steffen Lindert^{2,4}, Peter Kekenyes-Huskey^{2,4}, Vijay S. Rao¹, R. John Solaro⁶,
Paul R. Rosevear⁵, Rommie Amaro², Andrew D. McCulloch^{2,3}, J. Andrew McCammon^{2,4},
Michael Regnier^{1,*}

¹University of Washington, Department of Bioengineering, Seattle, WA, USA

²University of California San Diego, National Biomedical Computational Resource (NBCR),
LaJolla, CA, USA

³University of California San Diego, Department of Bioengineering, LaJolla, CA, USA

⁴University of California San Diego, Department of Pharmacology, LaJolla, CA, USA

⁵University of Cincinnati, Department of Molecular Genetics, Biochemistry, and Microbiology,
Cincinnati, OH, USA

⁶University of Illinois at Chicago, Department of Physiology and Biophysics, College of Medicine,
Chicago, IL, USA

Running Head: Troponin phosphomimics simulations

Corresponding Author:

Michael Regnier, Ph.D. Department of Bioengineering, University of Washington, Box 358056, 850
Republican St. Seattle, WA 98195. Email: mregnier@uw.edu

Contents	Page
Figure S1. The root-mean-square-displacement (RMSD) values of the backbone atoms (C, CA, N) for the two models with respect to the first snapshot as a function of time.	3
Figure S2. Distances between Ca ²⁺ and its coordinating cTnC site II residues (Asp 65, Asp 67, Asp 73 and Glu 76) over the course of each MD simulation for two complexes. Here, the 1 st run result is shown in black, the 2 nd run result is in red, and the 3 rd run result is in blue.	4
Figure S3. Contact maps of residue-residue pairs between the B- (residues 38-48) and C-helices (residues 54-64) of cTnC during each 150 ns MD simulation for two models. The blue end of the spectrum (value 0) reflects no contact between residue-residue pair, while the red end of the spectrum (value 1) represents 100% contact between residue-residue pair.	5
Figure S4. Contact maps of residue-residue pairs between NcTnC (cTnC residues 1-89) and NcTnI (cTnI residues 1-41) during each 150 ns MD simulation for two models.	6
Figure S5. The average angles and distances between the B- (residues 38-48) and C-helices (residues 54-64) of cTnC during 150 ns MD simulations for two models.	7
Figure S6. Contact maps of residue-residue pairs between NcTnC and the switch peptide of cTnI (cTnI residues 148-164) during each 150 ns MD simulation for two models.	8
Figure S7. The average angles and distances between the A- (residues 14-28) and B-helices (residues 38-48) of cTnC during 150 ns MD simulations for two models.	9
Figure S8. The average solvent accessible surface area (SASA) values for two models.	10
Figure S9. Contact maps of residue-residue pairs between cTnC and the inhibitory peptide of cTnI (cTnI residues 138-147) during each 150 ns MD simulation for two models.	11

RMSD WT vs cTnI-S23D/S24D




Figure S1




Figure S2




Figure S3




Figure S4




Figure S5




Figure S6




Figure S7




Figure S8




Figure S9