



Table S2. The frequency of PIN and cancer in mouse prostates

<b>Strain</b>	<b>Age</b>	<b>#Mice</b>	<b>Phenotype</b>
$\delta$ -Cat <sup>-/-</sup>	≤ 6 weeks	3	3/3 normal
	≤ 3 months	2	2/2 normal
	≥ 6 months	2	2/2 normal
	≥ 11 months	2	2/2 normal
No Myc/ $\delta^{+/+}$	≤ 6 weeks	4	4/4 normal
	≥ 6 months	3	3/3 normal
Myc/ $\delta^{+/+}$	≤ 4 weeks	5	5/5 epithelial hyperplasia
	≤ 6 weeks	5	5/5 low grade mPIN
	≥ 6 months	5	5/5 cribriform mPIN
	≥ 8 months	3	1/3 invasive cancer 2/3 cribriform PIN
Myc/ $\delta^{-/-}$	≤ 6 weeks	3	3/3 cribriform mPIN
	≥ 6 months	5	5/5 cribriform mPIN
	≥ 8 months	3	3/3 invasive cancer
Myc/ $\delta^{-/-}$	≤ 6 weeks	4	4/4 cribriform mPIN
	≥ 6 months	3	3/3 invasive cancer
	≥ 8 months	3	3/3 invasive cancer