

BMJ Open

Sexual violence and neonatal outcomes: a population-based cohort study

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2014-005935
Article Type:	Research
Date Submitted by the Author:	19-Jun-2014
Complete List of Authors:	Henriksen, Lena; Oslo University Hospital, Department for Women's and Children's Health Schei, Berit; Norwegian University of Science and Technology, Department of Public Health and General Practice, Faculty of Medicine; St Olavs University Hospital, Department of Obstetrics and Gynaecology Vangen, Siri; Oslo University Hospital, Department for Women's and Children's Health Lukasse, Mirjam; Oslo and Akershus University College, Norway, Faculty of Health Sciences
Primary Subject Heading:	Obstetrics and gynaecology
Secondary Subject Heading:	Obstetrics and gynaecology, Epidemiology
Keywords:	Fetal medicine < OBSTETRICS, Maternal medicine < OBSTETRICS, EPIDEMIOLOGY

SCHOLARONE™
Manuscripts

only

TITLE PAGE

Title: Sexual violence and neonatal outcomes: a population-based cohort study

Lena Henriksen¹, Berit Schei^{2,3}, Siri Vangen⁴, Mirjam Lukasse⁵

Corresponding author: Lena Henriksen¹, Section of Obstetrics, Department for Women's and Children's Health, Oslo University Hospital, Postboks 4950 Nydalen, N-0424 Oslo, Norway.

E-mail: uxhelz@ous-hf.no Phone: +47 90068485.

Co-authors: Berit Schei², Department of Public Health and General Practice, Faculty of Medicine, The Norwegian University of Science and Technology (NTNU), Håkon Jarls gate 11, N-7489 Trondheim, Norway. ³Department of Obstetrics and Gynecology, St.Olavs Hospital, Trondheim University Hospital, Postbox 3250 Sluppen, N-7006 Trondheim, Norway.

Siri Vangen⁴, National Resource Centre for Women's Health, Department for Women's and Children's Health, Oslo University Hospital, Postboks 4950 Nydalen, N-0424 Oslo, Norway.

Mirjam Lukasse⁵, Department of Health, Nutrition and Management, Faculty of Health Sciences, Oslo and Akershus University College of Applied Sciences, Postboks 4 St. Olavs plass, N-0130 Oslo, Norway.

Keywords: Sexual violence, rape, premature birth, low birth weight, small for gestational age, MoBa.

Word count: 3644

ABSTRACT

Objective: The objective of this study was to explore the association between sexual violence and neonatal outcomes.

Design: National cohort study.

Setting: Women were recruited to the Norwegian Mother and Child Cohort Study (MoBa) when attending routine ultrasound examinations from 1999-2008.

Population: A total of 76 870 pregnant women.

Methods: Sexual violence and maternal characteristics were self-reported in postal questionnaires during pregnancy. Neonatal outcomes were retrieved from the Medical Birth Registry of Norway (MBRN). Risk estimations were performed with linear and logistic regression analysis.

Outcome Measures: Gestational age at birth, birth weight, preterm birth (PTB), low birth weight (LBW) and small for gestational age (SGA).

Results: Of 76 870 women, 18.4% reported a history of sexual violence. A total of 4.7% had a premature birth, 2.7% had children with a birth weight <2500 g and 8.1% children were small for gestational age. Women reporting both moderate and severe sexual violence (rape) had a significantly reduced gestational length (2 days) when the birth was provider-initiated in an analysis adjusted for age, parity, education, smoking, BMI and mental distress. Those exposed to severe sexual violence had a significantly reduced gestational length of 0.51 days with a spontaneous start of birth. Crude estimates showed that severe sexual violence was associated with PTB, LBW and SGA. When controlling for the above-mentioned socio-demographic and behavioural factors, the association was no longer significant.

Conclusions: Sexual violence was not associated with adverse neonatal outcomes. Moderate and severe violence had a small but significant effect on gestational age; however, the clinical influence of this finding is most likely limited. Women exposed to sexual violence in this study reported more of the socio-demographic and behavioural factors associated with PTB, LBW and SGA compared with non-abused women.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- This study was based on information from a large population-based study described as sufficient for large-scale epidemiologic studies.
- All outcomes were collected prospectively from a quality-assessed birth register.
- The setting, with small social and health inequalities, was suitable to isolate the effect of sexual violence on adverse neonatal outcomes.
- A non-validated instrument for measuring of the exposure variable was a limitation to this study.

INTRODUCTION

Preterm birth is a common and costly health problem.[1, 2] Approximately one in ten babies are born preterm worldwide, and prematurity is considered to be the leading cause of death for newborns.[2] Low birth weight (LBW) can be a consequence of preterm birth (PTB) or intra uterine growth restriction, the latter leading to the birth of small for gestational age (SGA) infants.[3] Research has identified some biological risk factors for PTB and LBW: ethnicity, multiple pregnancies, a previous preterm birth and uterine or placental abnormalities.[1, 3] Studies also emphasise behavioural and social factors as important but less understood factors for both PTB and LBW.[1] These factors can be maternal age, socio-economic status, maternal weight, substance abuse, stress, depression and violence.[1, 2, 4]

Violence against women is a significant public health problem, and a recent report from the World Health Organization (WHO) state that 35% of women worldwide have experienced either physical and/or sexual intimate partner violence or non-partner sexual violence.[5] A pregnancy does not protect women from violence, and the prevalence of physical or sexual violence during pregnancy ranges from 3.4-11% in high-income countries.[6] It is recognised that violence has an adverse impact on women's physical, sexual, reproductive, and mental health.[5, 7] A connection between PTB or LBW and violence against women has been reported, but the association is both supported and contradicted.[5, 8-20] Studies have primarily addressed physical abuse during pregnancy and PTB or LBW [8, 10, 11, 16-19, 21] or child sexual abuse and PTB/LBW.[9, 12] Results from a new meta-analysis published in the recent WHO report [5] have demonstrated an association between intimate partner violence, including both physical and sexual abuse, and PTB with an adjusted odds ratio (AOR) of 1.41 (95% CI 1.21-1.62) and AOR of 1.16 (95% CI: 1.02-1.29) with LBW.[5]

1
2
3 However, studies on sexual violence and neonatal outcomes are limited, and few population-
4 based studies with large sample sizes that enable controlling for confounding variables have
5 been conducted [5, 11] Thus, there is a need to study the potential effect of lifetime sexual
6 violence on outcomes for newborns using a large cohort. The Norwegian Mother and Child
7 Cohort Study (MoBa) is a population-based prospective cohort study of pregnant women that
8 includes measurements of lifetime sexual violence and other relevant covariates. In this study,
9 we assessed the relationship between sexual violence and gestational age at birth and birth
10 weight. Additionally, we explored the associations between sexual violence and PTB, LBW
11 and SGA.
12
13
14
15
16
17
18
19
20
21
22
23
24

25 **METHODS**

26
27 This study was a sub-project in the MoBa study that was conducted by the Norwegian
28 Institute of Public health from 1999-2008.[22] All pregnant women in Norway were eligible
29 to participate in MoBa, and they were recruited during their routine foetal ultrasound
30 examination. Of the invited women, 40.6% consented to participate. Data were obtained
31 through extensive self-administered questionnaires that contained demographic factors,
32 general health, reproductive history and questions about maternal health status during
33 pregnancy. Our analyses were based on questionnaire 1 (Q1), which was completed during
34 (approximately) gestational week 17, and questionnaire 3 (Q3), which was completed during
35 (approximately) gestational week 30. Data from MoBa were linked with data from the
36 Medical Birth Registry of Norway (MBRN), which provided information on pregnancy and
37 birth outcome. The current study is based on version VI of the quality-assured data files
38 released for research in 2011. The MoBa study is described in detail elsewhere.[22]
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Study population

This study included women who had a singleton birth between 22 and 44 weeks of gestation, who completed both Q1 and Q3, had MBRN data available and participated for the first time (79 363 women). While a pregnancy is the observation unit in the MoBa study, women are the observation unit in our study; hence, the exclusion of 13 475 pregnancies of women who participated more than once. We excluded 703 women who did not answer the questions on sexual violence. Further, we excluded records with missing data on gestational length (n= 297) and birth weight (n=41). We also excluded 6 children with birth weight <500 g and 4 children with birth weight >6000 g, leaving a study sample of 76 870 women (Figure 1).

Ethical statement

Informed consent was obtained from each women participating in the study. The Regional Committee for Medical Research Ethics (Ref.SAFH 95/313 RTL) and the Norwegian Data Inspectorate approved the study.

Variables

Exposure variable

The exposure variable was collected from Q1; here, the women were asked if they had been 1) pressured to perform sexual acts, 2) forced with violence or 3) raped. A positive answer was defined as having experienced sexual violence. Women with more than one positive answer were classified according to the most severe level reported. The answers were coded into three levels of severity for the sexual violence: 1) mild, 2) moderate and 3) severe, and by the timing of the abuse (previously or recent). Recent meant exposed to sexual violence in the current pregnancy or the last 6 or 12 months before pregnancy. More details about the exposure variable can be found in our previous studies.[23;24]

Outcome variables

All outcome variables were obtained from the MBRN. Gestational age at birth in days was based on ultrasound at (approximately) gestational week 18. For women with no ultrasound, the gestational age was based on the last menstrual period (1.7%). PTB was defined as a gestational age <37, LBW as a birth weight <2500 grams, and SGA was defined as birth weight below the 10th percentile for the gestational age at birth. SGA was calculated using Norwegian specific foetal growth tables by Skjerven et al.[25]

Adjusting variables

Maternal age, parity, socio-economical status, smoking and body mass index (BMI) were considered as possible confounding factors and were adjusted for. All adjusting variables were taken from the MoBa. In Q1, age was categorised into 5 groups: younger than 20 years, 20–24 years, 25–29 years, 30–34 years or 35 years and older. As a proxy for socio-economic status, we used the woman's education in years (categorised into 4 groups): primary (<12 years), secondary (12 years), higher ≤ 4 years (13–16 years) and higher >4 years (≥ 17 years). Parity was dichotomised into nulli- and multiparous women. Smoking was categorised as no smoking or smoking, which included both daily and occasional smoking. BMI was grouped into 4 categories: <20, 20-24.9, 25.0-29.9 or ≥ 30.0 kg/m². We also adjusted for mental distress because it is considered to be associated with both the exposure and the outcome.[4,5] Mental distress was measured using 5 items from The Hopkins symptoms checklist (SCL-5) with a cut-off at ≥ 2.0 points, as suggested by Strand [26] and obtained from Q3.

Because of the co-occurrence of different violence types,[7] we examined the effect of physical and emotional abuse as a child or as an adult in the multivariable statistical models.

1
2
3 Information on adult physical violence was taken from Q1 and consisted of a positive answer
4 to whether women as an adult had experienced being slapped, hit, kicked or otherwise
5 bothered in a physical manner. Child physical violence was taken from Q3 and consisted of a
6 positive answer to the question “Have you experienced physical violence before the age of
7 18?” Emotional abuse as a child (<18) or as an adult (≥ 18) consisted of a positive answer to
8 either, “Someone has over a long period of time systematically tried to subdue, degrade or
9 humiliate you” or “Someone has threatened to hurt you or someone close to you” or both.
10
11
12
13
14
15
16
17
18
19

20
21 Previous preterm birth and inadequate antenatal care are considered to be associated with the
22 exposure and the outcomes.[2, 3, 5] Because a previous preterm birth may be a result of
23 sexual violence prior to the related pregnancy, we did not control for a previous preterm birth.
24 In Norway, the majority of women attend antenatal care, a free and a well-integrated part of
25 the public health system; therefore, we did not control for antenatal care. Ethnicity was not
26 considered a relevant confounding factor in our study because the majority of the MoBa
27 participants are ethnic Norwegian.
28
29
30
31
32
33
34
35
36
37

38 **Statistical Analysis**

39
40 Characteristics were presented as percentages within the entire sample and the different
41 outcomes. Linear regression was performed to assess differences in birth weight and
42 gestational age for children born to women with and without a history of mild, moderate and
43 severe sexual violence. The association between sexual violence and PTB, LBW and SGA
44 was estimated with crude and adjusted odds ratios using logistic regression analysis. All
45 analyses were adjusted for maternal age, parity, education, smoking, BMI and mental distress
46 in the first step. Birth weight was additionally adjusted for gestational age. We further
47 adjusted for other types of violence in the second step. We initially tested the correlation
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 between other types of violence and sexual violence because of co-occurrence, and all
4
5 Pearson's correlation coefficients were below the generally accepted cut-off of <0.4 for use as
6
7 a covariate in the regression analyses.[27] Post protocol, we stratified the sample into
8
9 spontaneous start of birth and provider-initiated start of birth (induced start of birth or elective
10
11 caesarean section) for gestational age because a provider-initiated start could influence the
12
13 time point of birth. Information on how the birth started was taken from MBRN. We
14
15 additionally performed a sensitivity analysis in which we examined the association between
16
17 sexual violence and SGA and LBW among women who had a spontaneous birth at term (≥ 37
18
19 weeks) because we wanted to examine the effect of violence in a group of women who were
20
21 considered to be low risk according to gestational age and start of birth. When we examined
22
23 the timing of the sexual violence, we compared women who were exposed to recent (sexual
24
25 violence during pregnancy or the last 6 or 12 months) and those exposed to previous sexual
26
27 violence during pregnancy or the last 6 or 12 months) and those exposed to previous sexual
28
29 violence to non-abused women. We also examined the timing among women reporting recent
30
31 and previous severe sexual violence (rape) for all outcomes. The prevalence of missing data
32
33 was generally low with 2.5% for BMI, 3.7% for education, and 0.7% for smoking during
34
35 pregnancy. Because of this, no imputing methods for missing data were used,[28] except for
36
37 the missing data for the SCL-5 (3.2%), which were replaced by the series mean. The results of
38
39 the logistic regression analyses remained approximately the same when performed with the
40
41 complete exclusion of missing data compared with using the imputed missing data for SCL-5.
42
43
44
45
46
47 The comparison group for all analyses was women not reporting any sexual violence. All
48
49 analyses were performed with the statistical package SPSS for WINDOWS (SPSS Inc.,
50
51 Chicago, IL, USA) version 18. P-values <0.05 were considered statistically significant.
52
53
54
55
56
57
58
59
60

RESULTS

The mean gestational age at birth in the total sample was 279.6 days (standard deviation 11.9 days), and the mean birth weight was 3592.7 grams (standard deviation 547.1 g). Table 1 displays the characteristics in the total sample and by the different outcomes.

Table 1. Background characteristics in the total sample and by preterm birth (PTB): gestational age <week 37, low birth weight (LBW): weight <2500 g and small for gestational age (SGA) weight below the 10th percentile by gestational age at birth in the Mother and Child Cohort

	Total	PTB	LBW	SGA
	N=76	N=3620	N=2107	N=6257
	870	%	%	%
	%			
<hr/>				
Age				
<20	1.5	2.2	2.6	1.9
20–24	12.2	12.7	13.8	13.2
25–29	36.2	34.6	34.8	36.4
30–34	35.7	33.7	33.2	34.2
≥35	14.4	16.7	15.6	14.3
Education				
Primary	2.4	3.1	3.9	2.7
Secondary	34.8	38.5	38.8	35.0
Higher ≤4 years	37.8	35.6	37.3	36.1
Higher >4 years	21.3	19.2	20.5	22.2
Missing	3.7	3.6	3.7	3.9
Parity				
Nulliparous	54.9	61.7	68.2	70.7
Multiparous	45.1	38.3	31.8	29.3
Smoking				
No	90.9	89.3	85.8	85.6
Yes	8.5	10.1	13.6	13.6
Missing	0.7	0.6	0.6	0.7
Pre-pregnancy body mass index				
<20	12.4	12.8	15.6	19.1
20–24.9	54.9	49.9	50.8	55.6
25–29.9	21.0	22.0	19.0	15.2
≥30	9.2	12.6	11.8	7.6
Missing	2.5	2.7	2.9	2.4
Mental distress				

No	93.3	92.3	91.3	92.3
Yes	6.8	7.7	8.7	7.7
Adult physical violence				
No	85.5	83.9	83.6	84.7
Yes	14.5	16.1	16.4	15.3
Child physical violence				
No	94.5	94.4	94.1	94.3
Yes	5.5	5.6	5.9	5.5
Adult emotional abuse				
No	83.6	82.1	81.3	82.5
Yes	16.4	17.9	18.7	17.5
Child emotional abuse				
No	86.2	85.1	84.7	85.5
Yes	13.8	14.9	15.3	14.5

The prevalence of adverse neonatal outcomes was generally highest in the youngest (<20 y) and the oldest age groups (≥ 35 y) among smokers and women with primary school education.

A BMI ≥ 30 was associated with PTB and LBW and BMI <20 with SGA. Women who reported mental distress also reported more PTB, LBW and SGA.

Among the 76 870 women enrolled, 9263 (12.1%) reported a history of mild sexual violence, 2102 (2.8%) moderate and 2746 (3.5%) severe. Women with a history of sexual violence were significantly younger and they were more likely to have primary school education. Additionally, these women more frequently reported smoking, a BMI ≥ 30 and mental distress. These women more often experienced other types of violence both as children and adults (data not provided in tables).

A lower gestational age at birth was observed for newborns from women who reported moderate and severe sexual violence with approximately two days when birth was provider-initiated (Table 2).

Table 2. Differences in gestational age for spontaneous and provider-initiated start of birth and birth weight for non-exposed women and women exposed to mild, moderate and severe sexual violence

	N= 76 870 (%)	Mean	Crude estimate β (95% CI)	Adjusted estimate β (95% CI)*
Gestational age				
No sexual violence	62 699 (81.6)	279.7 days	1	1
<i>Mild sexual violence</i>				
Spontaneous start	7323 (11.5)	279.9 days	0.09 (-0.16; 0.33)	0.22 (-0.04; -0.48)
Provider-initiated start	1940 (12.5)	280.3 days	0.36 (-0.37; 1.08)	0.64 (-0.11; 1.39)
<i>Moderate sexual violence</i>				
Spontaneous start	1670 (2.7)	279.3 days	-0.40 (-0.87; 0.06)	-0.28 (-0.78; 0.21)
Provider-initiated start	492 (3.2)	277.6 days	-2.13 (-3.41; -0.84)	-2.02 (-3.39; -0.67)
<i>Severe sexual violence</i>				
Spontaneous start	2048 (3.3)	278.7 days	-0.91 (-1.37; -0.44)	-0.65 (-1.15; -0.16)
Provider-initiated start	698 (4.5)	277.5 days	-2.24 (-3.47; -1.01)	-1.92 (-3.22; -0.62)
Birth weight				
No sexual violence	62 699 (81.6)	3594 g	1	1
Mild sexual violence	9263 (12.1)	3597 g	2.03 (-9.26; 13.32)	0.04 (-9.16; 9.23)**
Moderate sexual violence	2102 (2.8)	3582 g	-13.61 (-34.74; 7.51)	6.11 (-11.30; 23.52)**
Severe sexual violence	2746 (3.6)	3556 g	-38.33 (-59.17; -17.49)	-0.76 (-18.05; 16.53)**

*Adjusted for maternal age, parity, education, smoking, body mass index and mental distress

**Additional adjustment for gestational age

Among women with a spontaneous start of birth, the gestational age was approximately one half of a day shorter when women reported severe sexual violence. These findings were significant in an adjusted analysis. A crude analysis showed that women who reported a history of severe violence delivered on average 38.3 g lighter children, a difference that disappeared when controlling for gestational age, mother's age, parity, education, smoking, BMI and mental distress. There were no differences regarding birth weight between women with a history of mild or moderate sexual violence compared with non-abused women.

Results from the logistic regression analysis are presented in Table 3.

Table 3. Odds of preterm birth (PTB), low birth weight (LBW) and small for gestational age (SGA) with 95% confidence intervals (CI) according to the different levels of sexual violence

	N (%)	Prevalence (%)	Crude odds ratio (95% CI)	Adjusted odds ratio (95% CI)*
PTB		3620 (4.7)		

No sexual violence	62 699 (81.6)	2931 (4.7)	1	1
Mild sexual violence	9263 (12.1)	412 (4.4)	0.95 (0.85-1.06)	0.93 (0.83-1.03)
Moderate sexual violence	2102 (2.8)	115 (5.3)	1.15 (0.95-1.39)	1.14 (0.93-1.39)
Severe sexual violence	2746 (3.6)	162 (5.9)	1.28 (1.08-1.51)	1.15 (0.97-1.37)
LBW		2107 (2.7)		
No sexual violence	62 699 (81.6)	1681 (2.7)	1	1
Mild sexual violence	9263 (12.1)	257 (2.8)	1.04 (0.91-1.18)	0.98 (0.85-1.12)
Moderate sexual violence	2102 (2.8)	75 (3.5)	1.30 (1.03-1.65)	1.19 (0.93-1.53)
Severe sexual violence	2746 (3.6)	94 (3.4)	1.29 (1.04-1.59)	1.07 (0.85-1.34)
SGA		6257 (8.1)		
No sexual violence	62 699 (81.6)	5061 (8.1)	1	1
Mild sexual violence	9263 (12.1)	768 (8.3)	1.03 (0.95-1.12)	1.00 (0.91-1.08)
Moderate sexual violence	2102 (2.8)	178 (8.3)	1.02 (0.87-1.19)	0.95 (0.80-1.12)
Severe sexual violence	2746 (3.6)	250 (9.1)	1.14 (1.00-1.30)	1.05 (0.91-1.21)

*Adjusted for maternal age, parity, education, smoking, body mass index and mental distress

Women who reported severe sexual violence had higher odds of PTB, LBW and SGA in a crude analysis, an association that was attenuated and no longer significant when adjusted for maternal age, parity, education, smoking, BMI and mental distress. Other types of violence, both as a child and adult, had small attenuating effects on the odds ratios and were not included in the final models.

The sensitivity analysis, in which we examined the association between a history of sexual violence and SGA and LBW in a sub-sample of women who had a spontaneous term birth, showed the same pattern as in the total sample reported in Table 3. Women who reported severe sexual violence had higher odds of LBW and SGA in a crude analysis but not in the adjusted analysis (Data not provided in tables).

A crude analysis was used to examine if the timing of the violence was associated with adverse outcome. Women who reported recent sexual violence had a higher risk for LBW (OR 1.60 95% CI 1.04-2.17) compared with non-abused women. The association was no longer significant in the adjusted analysis. In our study, 684 (0.9%) women reported recent

1
2
3 sexual violence (mild, moderate and severe) and 13 487 (17.5%) previous sexual violence
4
5 (Supplementary Table S1). There was no association between recent severe sexual violence
6
7 (rape) and adverse neonatal outcome (Supplementary Table S2). There were 66 (0.1%)
8
9 women who reported rape during pregnancy or the last 6 or 12 months in this study.
10
11

12 13 14 **DISCUSSION**

15 16 **Main outcome**

17
18 We found that moderate and severe sexual violence were associated with a reduction in
19
20 gestational age at birth. The largest effect was observed when birth was provider-initiated
21
22 among women exposed to moderate or severe violence. These women had an approximately
23
24 two-day reduction in gestational age. There was no significant association between sexual
25
26 violence and PTB, LBW or SGA in the adjusted analysis.
27
28

29 30 31 **Strength**

32
33 This study, based on information from a large population-based study, the Norwegian Mother
34
35 and Child Cohort study (MoBa), which is linked to the Medical Birth Registry (MBRN), gave
36
37 a unique opportunity to assess the association between sexual violence and outcome for
38
39 newborns. The validity of the data in the MoBa has in earlier research been described as
40
41 sufficient for large-scale epidemiologic studies.[29, 30] Our study was strengthened by the
42
43 fact that the information on the different outcome variables was collected prospectively from
44
45 the quality-assessed MBRN.[31] The outcomes in this study are part of a complex
46
47 phenomenon that has several different risk factors. [2] The setting in this study, with small
48
49 social and health inequalities, may therefore be suitable to isolate the effect of sexual violence
50
51 on adverse neonatal outcomes.
52
53
54
55
56
57
58
59
60

Limitations

There are also limitations to our study. The participation rate of 40.6% in MoBa is low, and MoBa suffers to some extent from selection bias. The women included in the study are older, have more education, smoke less and are less likely to be of a non-Norwegian origin than the Norwegian population. A recent study has found that this may affect the prevalence estimates, but there was no evidence that the exposure-outcome associations were affected by selection bias.[32] Furthermore, as preterm birth is associated with ethnicity, the ethnic homogeneous sample in MoBa may limit the generalisability of our findings. The lack of a validated instrument for measuring the exposure is a limitation to this study, and violence measured in modules as part of a larger questionnaire, as that in MoBa, may achieve a lower disclosure rate.[5] However, a similar prevalence to that found in our study was reported in a Nordic study examining sexual violence and health.[33]

To our knowledge, no studies have examined the influence of lifetime sexual violence reported during pregnancy on the gestational age at birth for newborns. There were minor differences in the gestational age between abused and non-abused women in this study, and the clinical importance of our findings for the health of the newborn is most likely limited. However, the difference between the provider-initiated and spontaneous initiation of birth may be of interest. Shorter provider-initiated pregnancies may suggest an increase in elective inductions and elective caesarean sections for those exposed to violence. This is supported by others [34, 35] and in our previous study on sexual violence and maternal outcome.[36] Studies have emphasised the meaning of control for abused women when giving birth,[37] and choosing a planned start of birth may help the abused women remain in control.

1
2
3 Unlike the current meta-analysis mentioned in a WHO report (not yet published),[5] we did
4
5 not find sexual violence to be associated with PTB and LBW in adjusted analysis. To our
6
7 knowledge, no study has found an association between violence and SGA. The studies
8
9 included in the meta-analysis investigated both sexual and physical intimate partner violence,
10
11 in contrast to our study, which investigated lifetime sexual violence. Studies on the health
12
13 effects of non-partner sexual violence are limited with less robust data than for intimate
14
15 partner violence.[5] Our data from the MoBa does not differentiate between intimate partner
16
17 violence and sexual abuse by non-partners, but it is not unlikely that the question about severe
18
19 sexual violence (rape) primarily reflects non-partner sexual abuse. Some studies have shown
20
21 significant associations for adverse neonatal outcomes in a crude analysis; these became non-
22
23 significant when adjusting for socio-demographic and behavioural factors,[38, 39] as ours.
24
25 The exposure in these studies involved intimate partner violence during pregnancy, and the
26
27 study by Webster et al. [39] also included lifetime physical violence.
28
29
30
31
32
33

34 Several pathways between sexual violence and adverse neonatal outcomes are suggested, both
35
36 direct and indirect.[5] The direct pathway of violence during pregnancy can lead to a preterm
37
38 birth, and examples of indirect pathways include more health-risk behaviours, depression and
39
40 stress/anxiety.[5, 40] Both experienced violence and living in an abusive environment can
41
42 cause increased stress levels, which could be on the pathway between abuse and adverse
43
44 neonatal outcome. Maternal exposure to stress can influence the hypothalamic pituitary axis
45
46 (HPA) hormones, and it is suggested that changes in these hormones may cause negative
47
48 outcomes, such as a reduction in gestational age and foetal growth restriction.[4, 41]
49
50

51 The prevalence of abuse during pregnancy is small in our study and may have decreased the
52
53 power to detect an association between violence during pregnancy and adverse neonatal
54
55 outcomes, an association that is supported by other studies.[10, 14, 42, 43] In our study,
56
57
58
59
60

1
2
3 sexual violence was assessed approximately in gestational week 17, and events of violence
4
5 after that may have been missed. The studies that report the highest prevalence of violence
6
7 have measured this several times during pregnancy. [6] Studies have reported a violence
8
9 prevalence during pregnancy of between 3 to 19%. [6, 44] A direct comparison to our study is
10
11 difficult because these numbers estimate both physical and sexual abuse during pregnancy;
12
13 they may suggest that sexual violence during pregnancy is underreported in our study.
14
15 Underreporting among the non-exposed may have caused a misclassification that has
16
17 diminished the associations between sexual violence and neonatal outcomes in our study.
18
19 Living in an abusive relationship may also stop women from disclosing the violence.
20
21 However, our prevalence of recent sexual (0.9%) violence is similar to the prevalence of 1%
22
23 reported in a national population based study on violence among Norwegian women. [45] This
24
25 number reflects the prevalence of sexual violence reported during the last year, not especially
26
27 among pregnant women.
28
29
30
31
32
33

34 **Conclusion**

35
36 Overall, our findings provide no evidence for an association between lifetime sexual violence
37
38 and adverse neonatal outcomes. A small significant effect on the gestational age at birth was
39
40 detected, but the clinical importance of this is most likely limited for the health of the
41
42 newborn. PTB, LBW and SGA all have complex origins with multiple possible pathways. [1,
43
44 2] Although we did not find an association between sexual violence and PTB, LBW or SGA
45
46 in adjusted analyses, crude analyses in our study suggested that sexual violence may be a risk
47
48 factor for adverse neonatal outcomes for some women; however, for the majority of women,
49
50 the relationship is confounded by other risk factors. Antenatal care is one of the most
51
52 important entry points in which women seek health care without necessarily disclosing
53
54 ongoing exposure to violence or a history of sexual violence. It is recommended that
55
56
57
58
59
60

1
2
3 caregivers and clinicians ask women about exposure to violence when assessing conditions
4
5 that may be complicated by violence,[46] and more knowledge about this conditions is
6
7 needed. Additionally, antenatal care may offer opportunities for women to receive help, both
8
9 if they are exposed to violence and in providing assistance to change behavioural factors
10
11 contributing to adverse outcomes.
12

13 14 15 16 **Competing interests**

17
18 None.
19

20 21 22 23 **Contributions to authorship**

24
25 LH participated in the conception and design of the study, performed the analysis and drafted
26
27 the manuscript. BS participated in the conception of the study, advised on the statistical
28
29 analyses and drafting of the manuscript. SV advised on the statistical analyses and the drafting
30
31 of the manuscript. ML participated in the conception and design of the study, advised on the
32
33 statistical analyses and participated in drafting the manuscript. All authors read and approved
34
35 the final version.
36
37
38
39

40 41 **Funding**

42
43 The Norwegian Mother and Child Cohort Study is supported by the Norwegian Ministry of
44
45 Health and the Ministry of Education and Research, NIH/NIEHS (contract no N01-ES-
46
47 75558), NIH/NINDS (grant no.1 U01 NS 047537-01 and grant no.2 U01 NS 047537-06A1),
48
49 and the Norwegian Research Council/FUGE (grant no. 151918/S10). The first author of this
50
51 article is supported by The Research Council of Norway (grant no. 204292/H10), and the
52
53 second author is supported by the South-Eastern Norway Regional Health Authority (grant
54
55 no. 2012010).
56
57
58
59
60

Acknowledgement

We are grateful to all the participating families in Norway who take part in this ongoing cohort study.

Data Sharing Statement

No additional data are available.

Reference List

- (1) Green NS, Damus K, Simpson JL, et al. Research agenda for preterm birth: recommendations from the March of Dimes. *Am J Obstet Gynecol* 2005;193:626-35.
- (2) March of Dimes, PMNCH, Save The Children, WHO. Born Too Soon: The Global Action Report on Preterm Birth. Geneva: World Health Organization 2012
- (3) Wilcox AJ. On the importance-and the unimportance-of birth weight. *Int J Epidemiol* 2001;30:1233-41.
- (4) Lazinski MJ, Shea AK, Steiner M. Effects of maternal prenatal stress on offspring development: a commentary. *Arch Womens Ment Health* 2008;11:363-75.
- (5) Garcia-Moreno C, Pallitto C, Devries K, et al. Global and regional estimates of violence against women: prevalence and health effects of intimate partner violence and non-partner sexual violence. Geneva: World Health Organization; 2013.
- (6) Campbell J, Garcia-Moreno C, Sharps P. Abuse During Pregnancy in Industrialized and Developing Countries. *Violence Against Women* 2004;10:770-89.
- (7) Garcia-Moreno C, Jansen HA, Ellsberg M, et al. Prevalence of intimate partner violence: findings from the WHO multi-country study on women's health and domestic violence. *Lancet* 2006;368:1260-9.
- (8) Sanchez SE, Alva AV, Diez CG, et al. Risk of spontaneous preterm birth in relation to maternal exposure to intimate partner violence during pregnancy in Peru. *Matern Child Health J* 2013;17:485-92.
- (9) Leeners B, Stiller R, Block E, et al. Pregnancy complications in women with childhood sexual abuse experiences. *J Psychosom Res* 2010;69:503-10.
- (10) Nunes MA, Camey S, Ferri CP, et al. Violence during pregnancy and newborn outcomes: a cohort study in a disadvantaged population in Brazil. *Eur J Public Health* 2011;21(1):92-7.
- (11) Rodrigues T, Rocha L, Barros H. Physical abuse during pregnancy and preterm delivery. *Am J Obstet Gynecol* 2008;198:171-6.
- (12) Noll JG, Schulkin J, Trickett PK, et al. Differential pathways to preterm delivery for sexually abused and comparison women. *J Pediatr Psychol* 2007;32:1238-48.

- 1
- 2
- 3 (13) Silverman JG, Decker MR, Reed E, et al. A. Intimate partner violence victimization
- 4 prior to and during pregnancy among women residing in 26 U.S. states: associations
- 5 with maternal and neonatal health. *Am J Obstet Gynecol* 2006;195:140-8.
- 6
- 7 (14) Murphy CC, Schei B, Myhr TL, et al. Abuse: a risk factor for low birth weight? A
- 8 systematic review and meta-analysis. *CMAJ* 2001;164:1567-72.
- 9
- 10 (15) Fried LE, Cabral H, Amaro H, et al. Lifetime and during pregnancy experience of
- 11 violence and the risk of low birth weight and preterm birth. *J Midwifery Womens*
- 12 *Health* 2008 ;53:522-8.
- 13
- 14 (16) Janssen PA, Holt VL, Sugg NK, et al. Intimate partner violence and adverse
- 15 pregnancy outcomes: a population-based study. *Am J Obstet Gynecol* 2003;188:1341-
- 16 7.
- 17
- 18 (17) Urquia ML, O'Campo PJ, Heaman MI, et al. Experiences of violence before and
- 19 during pregnancy and adverse pregnancy outcomes: an analysis of the Canadian
- 20 Maternity Experiences Survey. *BMC Pregnancy Childbirth* 2011;11:42.
- 21
- 22 (18) Altarac M, Strobino D. Abuse during pregnancy and stress because of abuse during
- 23 pregnancy and birthweight. *J Am Med Womens Assoc* 2002;57:208-14.
- 24
- 25 (19) Cokkinides VE, Coker AL, Sanderson M, et al. Physical violence during pregnancy:
- 26 maternal complications and birth outcomes. *Obstet Gynecol* 1999;93:661-6.
- 27
- 28 (20) Grimstad H, Schei B, Backe B, et al. Physical abuse and low birthweight: a case-
- 29 control study. *Br J Obstet Gynaecol* 1997;104:1281-7.
- 30
- 31 (21) Kaye DK, Mirembe FM, Bantebya G, et al. Domestic violence during pregnancy and
- 32 risk of low birthweight and maternal complications: a prospective cohort study at
- 33 Mulago Hospital, Uganda. *Trop Med Int Health* 2006;11:1576-84.
- 34
- 35 (22) Magnus P, Irgens LM, Haug K, et al. Cohort profile: the Norwegian Mother and Child
- 36 Cohort Study (MoBa). *Int J Epidemiol* 2006;35:1146-50.
- 37
- 38 (23) Lukasse M, Henriksen L, Vangen S, et al. Sexual violence and pregnancy-related
- 39 physical symptoms. *BMC Pregnancy Childbirth* 2012;12:83.
- 40
- 41 (24) Henriksen L, Vangen S, Schei B, et al. Sexual violence and antenatal hospitalization.
- 42 *Birth* 2013;40:281-8.
- 43
- 44 (25) Skjaerven R, Gjessing HK, Bakketeig LS. et al. Birthweight by gestational age in
- 45 Norway. *Acta Obstet Gynecol Scand* 2000;79:440-9.
- 46
- 47 (26) Strand BH, Dalgard OS, Tambs K, et al. Measuring the mental health status of the
- 48 Norwegian population: a comparison of the instruments SCL-25, SCL-10, SCL-5 and
- 49 MHI-5 (SF-36). *Nord J Psychiatry* 2003;57:113-8.
- 50
- 51 (27) Field A. Discovering statistics using SPSS. Los Angeles: SAGE; 2009.
- 52
- 53 (28) Veierød MB, Lydersen S, Laake P, et at. Medical statistics: in clinical and
- 54 epidemiological research. Oslo: Gyldendal akademisk; 2012.
- 55
- 56
- 57
- 58
- 59
- 60

- 1
2
3 (29) Kvalvik LG, Nilsen RM, Skjaerven R, et al. Self-reported smoking status and plasma
4 cotinine concentrations among pregnant women in the Norwegian Mother and Child
5 Cohort Study. *Pediatr Res* 2012 ;72:101-7.
6
7 (30) Furu K, Karlstad O, Skurtveit S, et al. High validity of mother-reported use of
8 antiasthmatics among children: a comparison with a population-based prescription
9 database. *J Clin Epidemiol* 2011;64:878-84.
10
11 (31) Irgens LM. The Medical Birth Registry of Norway. Epidemiological research and
12 surveillance throughout 30 years. *Acta Obstet Gynecol Scand* 2000;7:435-9.
13
14 (32) Nilsen RM, Vollset SE, Gjessing HK, et al. Self-selection and bias in a large
15 prospective pregnancy cohort in Norway. *Paediatr Perinat Epidemiol* 2009 23:597-
16 608.
17
18 (33) Hilden M, Schei B, Swahnberg K, et al. A history of sexual abuse and health: a Nordic
19 multicentre study. *BJOG* 2004;111:1121-7.
20
21 (34) Nerum H, Halvorsen L, Straume B, et al. Different labour outcomes in primiparous
22 women that have been subjected to childhood sexual abuse or rape in adulthood: a
23 case-control study in a clinical cohort. *BJOG* 2013;120:487-495.
24
25 (35) Lukasse M, Vangen S, Oian P, et al. Childhood abuse and caesarean section among
26 primiparous women in the Norwegian Mother and Child Cohort Study. *BJOG*
27 2010;117:1153-7.
28
29 (36) Henriksen L, Schei B, Vangen S, et al. Sexual violence and mode of delivery: a
30 population-based cohort study. *BJOG*. In press 2014.
31
32 (37) Montgomery E. Feeling Safe: A Metasynthesis of the Maternity Care Needs of
33 Women Who Were Sexually Abused in Childhood. *Birth* 2013;40:88-95.
34
35 (38) Campbell J, Torres S, Ryan J, et al. Physical and nonphysical partner abuse and other
36 risk factors for low birth weight among full term and preterm babies: a multiethnic
37 case-control study. *Am J Epidemiol* 1999;150:714-26.
38
39 (39) Webster J, Chandler J, Battistutta D. Pregnancy outcomes and health care use: effects
40 of abuse. *Am J Obstet Gynecol* 1996;174:760-7.
41
42 (40) Leeners B, Neumaier-Wagner P, Quarg AF, et al. Childhood sexual abuse (CSA)
43 experiences: an underestimated factor in perinatal care. *Acta Obstet Gynecol Scand*
44 2006;85:971-6.
45
46 (41) Valladares E, Pena R, Ellsberg M, et al. Neuroendocrine response to violence during
47 pregnancy--impact on duration of pregnancy and fetal growth. *Acta Obstet Gynecol*
48 *Scand* 2009;88:818-23.
49
50 (42) Faramarzi M, Esmaelzadeh S, Mosavi S. Prevalence, maternal complications and birth
51 outcome of physical, sexual and emotional domestic violence during pregnancy. *Acta*
52 *Med Iran* 2005;43:115-22.
53
54
55
56
57
58
59
60

- 1
2
3 (43) Valladares E, Ellsberg M, Pena R, et al. Physical partner abuse during pregnancy: a
4 risk factor for low birth weight in Nicaragua. *Obstet Gynecol* 2002;100:700-5.
5
6 (44) Sharps PW, Laughon K, Giangrande SK. Intimate partner violence and the
7 childbearing year: maternal and infant health consequences. *Trauma Violence Abuse*
8 *2007;8:105-16.*
9
10 (45) Neroien AI, Schei B. Partner violence and health: results from the first national study
11 on violence against women in Norway. *Scand J Public Health* 2008;36:161-8.
12
13 (46) World health organisation. Responding to intimate partner violence and sexual
14 violence against women: WHO clinical and policy guidelines. Geneva: WHO; 2013.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1. Inclusion and exclusion process




Table S1: Odds of preterm birth (PTB), low birth weight (LBW) and small for gestational age (SGA) with 95% confidence intervals (CI) according to previous and recent sexual violence

	N (%)	Prevalence (%)	Crude odds ratio (95% CI)	Adjusted odds ratio (95% CI)
PTB		3620 (4.7)		
No sexual violence	62 699 (81.6)	2931 (4.7)	1	1
Previous sexual violence	13487 (17.5)	650 (4.8)	1.03 (0.95-1.13)	0.99 (0.90-1.09)
Recent sexual violence	684 (0.9)	39 (5.7)	1.23 (0.89-1.71)	1.17 (0.83-1.65)
LBW		2107 (2.7)		
No sexual violence	62 699 (81.6)	1681 (2.7)	1	1
Previous sexual violence	13 487 (17.5)	397 (2.9)	1.10 (0.99-1.23)	1.0 (0.86-1.12)
Recent sexual violence	684 (0.9)	29 (4.2)	1.60 (1.04-2.33)	1.45 (0.97-2.17)
SGA		6257 (8.1)		
No sexual violence	62 699 (81.6)	5061 (8.1)	1	1
Previous sexual violence	13 487 (17.5)	1136 (8.4)	1.05 (0.98-1.12)	0.98 (0.92-1.06)
Recent sexual violence	684 (0.9)	60 (8.8)	1.10 (0.84-1.43)	1.18 (0.90-1.56)

*Adjusted for maternal age, parity, education, smoking, body mass index and mental distress

Table S2: Odds of preterm birth (PTB), low birth weight (LBW) and small for gestational age (SGA) with 95% confidence intervals (CI) according to previous and recent severe violence (rape)

	N (%)	Prevalence (%)	Crude odds ratio (95% CI)	Adjusted odds ratio (95% CI)*
PTB		3620 (4.7)		
No rape	74 124 (96.4)	3458 (4.7)	1	1
Previous rape	2680 (3.5)	157 (5.9)	1.27 (1.08-1.50)	1.14 (0.97-1.36)
Recent rape	66 (0.1)	5 (7.6)	1.68 (0.67-4.17)	1.72 (0.68-4.33)
LBW		2107 (2.7)		
No rape	74 124 (96.4)	2013(2.7)	1	1
Previous rape	2680 (3.5)	91 (3.4)	1.26 (1.02-1.56)	1.06 (0.84-1.33)
Recent rape	66 (0.1)	3 (4.5)	1.71 (0.54-5.44)	1.55 (0.48-5.02)
SGA		6257 (8.1)		
No rape	74 124 (96.4)	6007 (8.1)	1	1
Previous rape	2680 (3.5)	247 (9.2)	1.15 (1.01-1.32)	1.06 (0.92-1.23)
Recent rape	66 (0.1)	3 (4.5)	0.54 (0.17-1.72)	0.55 (0.17-1.79)

*Adjusted for maternal age, parity, education, smoking, body mass index and mental distress

STROBE 2007 (v4) Statement—Checklist of items that should be included in reports of cohort studies

Section/Topic	Item #	Recommendation	Reported on page #
Title and abstract	1	(a) Indicate the study’s design with a commonly used term in the title or the abstract	page 1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	page 2
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	page 3 and
Objectives	3	State specific objectives, including any prespecified hypotheses	page 4
Methods			
Study design	4	Present key elements of study design early in the paper	From page 4
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection in addition reference is given for further reading of detailed information on MoBa study (referece 22)	Page 4
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up	NA
		(b) For matched studies, give matching criteria and number of exposed and unexposed	NA
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	Presented on pages 5-7
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	Page 5-7
Bias	9	Describe any efforts to address potential sources of bias controlling for confounding and covariates	pages 6 and 7.
Study size	10	Explain how the study size was arrived at	figure 1 and page5
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	Page 6
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	page 7 and 8
		(b) Describe any methods used to examine subgroups and interactions	page 8
		(c) Explain how missing data were addressed	page 8
		(d) If applicable, explain how loss to follow-up was addressed	NA
		(e) Describe any sensitivity analyses	page 8
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed All listed in Figure 1	
		(b) Give reasons for non-participation at each stage	Page 5

		(c) Consider use of a flow diagram Figure 1	
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	Table 1: page 9 and 10
		(b) Indicate number of participants with missing data for each variable of interest	Table 1
		(c) Summarise follow-up time (eg, average and total amount)	NA
Outcome data	15*	Report numbers of outcome events or summary measures over time	Table 1
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	Page 6-7, 11 - 12 and Table 2 and 3
		(b) Report <i>category</i> boundaries when continuous variables were categorized	Page 6 and Table 1
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	page 12-13 Supplementary Tables 1 and 2
Discussion			
Key results	18	Summarise key results with reference to study objectives	page 13
Limitations			
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	Page 13-16
Generalisability	21	Discuss the generalisability (external validity) of the study results	Page 14
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	Page 17

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

BMJ Open

Sexual violence and neonatal outcomes: a Norwegian population-based cohort study

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2014-005935.R1
Article Type:	Research
Date Submitted by the Author:	05-Sep-2014
Complete List of Authors:	Henriksen, Lena; Oslo University Hospital, Department for Women's and Children's Health Schei, Berit; Norwegian University of Science and Technology, Department of Public Health and General Practice, Faculty of Medicine; St Olavs University Hospital, Department of Obstetrics and Gynaecology Vangen, Siri; Oslo University Hospital, Department for Women's and Children's Health Lukasse, Mirjam; Oslo and Akershus University College, Norway, Faculty of Health Sciences
Primary Subject Heading:	Obstetrics and gynaecology
Secondary Subject Heading:	Obstetrics and gynaecology, Epidemiology
Keywords:	Fetal medicine < OBSTETRICS, Maternal medicine < OBSTETRICS, EPIDEMIOLOGY

SCHOLARONE™
Manuscripts

Only

TITLE PAGE

Title: Sexual violence and neonatal outcomes: a Norwegian population-based cohort study

Lena Henriksen¹, Berit Schei^{2,3}, Siri Vangen⁴, Mirjam Lukasse⁵

Corresponding author: Lena Henriksen¹, Section of Obstetrics, Department for Women's and Children's Health, Oslo University Hospital, Postboks 4950 Nydalen, N-0424 Oslo, Norway.

E-mail: uxhelz@ous-hf.no Phone: +47 90068485.

Co-authors: Berit Schei², Department of Public Health and General Practice, Faculty of Medicine, The Norwegian University of Science and Technology (NTNU), Håkon Jarls gate 11, N-7489 Trondheim, Norway. ³Department of Obstetrics and Gynecology, St.Olavs Hospital, Trondheim University Hospital, Postbox 3250 Sluppen, N-7006 Trondheim, Norway.

Siri Vangen⁴, National Resource Centre for Women's Health, Department for Women's and Children's Health, Oslo University Hospital, Postboks 4950 Nydalen, N-0424 Oslo, Norway.

Mirjam Lukasse⁵, Department of Health, Nutrition and Management, Faculty of Health Sciences, Oslo and Akershus University College of Applied Sciences, Postboks 4 St. Olavs plass, N-0130 Oslo, Norway.

Keywords: Sexual violence, rape, premature birth, low birth weight, small for gestational age, MoBa.

Word count: 4249

ABSTRACT

Objective: The objective of this study was to explore the association between sexual violence and neonatal outcomes.

Design: National cohort study.

Setting: Women were recruited to the Norwegian Mother and Child Cohort Study (MoBa) when attending routine ultrasound examinations from 1999-2008.

Population: A total of 76 870 pregnant women.

Methods: Sexual violence and maternal characteristics were self-reported in postal questionnaires during pregnancy. Neonatal outcomes were retrieved from the Medical Birth Registry of Norway (MBRN). Risk estimations were performed with linear and logistic regression analysis.

Outcome measures: Gestational age at birth, birth weight, preterm birth (PTB), low birth weight (LBW) and small for gestational age (SGA).

Results: Of 76 870 women, 18.4% reported a history of sexual violence. A total of 4.7% had a premature birth, 2.7% had children with a birth weight <2500 g and 8.1% children were small for gestational age. Women reporting both moderate and severe sexual violence (rape) had a significantly reduced gestational length (2 days) when the birth was provider-initiated in an analysis adjusted for age, parity, education, smoking, BMI and mental distress. Those exposed to severe sexual violence had a significantly reduced gestational length of 0.51 days with a spontaneous start of birth. Crude estimates showed that severe sexual violence was associated with PTB, LBW and SGA. When controlling for the above-mentioned socio-demographic and behavioural factors, the association was no longer significant.

Conclusions: Sexual violence was not associated with adverse neonatal outcomes. Moderate and severe violence had a small but significant effect on gestational age; however, the clinical influence of this finding is most likely limited. Women exposed to sexual violence in this study reported more of the socio-demographic and behavioural factors associated with PTB, LBW and SGA compared with non-abused women.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- This study was based on information from a large population-based study described as sufficient for large-scale epidemiologic studies.
- All outcomes were collected prospectively from a quality-assessed birth register.
- The setting, with small social and health inequalities, was suitable to isolate the effect of sexual violence on adverse neonatal outcomes.
- A non-validated instrument for measuring of the exposure variable was a limitation to this study.

INTRODUCTION

Preterm birth is a common and costly health problem.[1, 2] Approximately one in ten babies are born preterm worldwide, and prematurity is considered to be the leading cause of death for newborns.[2] Low birth weight (LBW) can be a consequence of preterm birth (PTB) or intra uterine growth restriction, the latter leading to the birth of small for gestational age (SGA) infants.[3] Research has suggested some biological risk factors for PTB and LBW: multiple pregnancies, a previous preterm birth and uterine or placental abnormalities.[1, 3] Studies also emphasise other, less understood factors for both PTB and LBW.[1] These are maternal age, socio-economic status, ethnicity, maternal weight, substance abuse, stress, depression and violence.[1, 2, 4]

Violence against women is a significant public health problem, and a recent report from the World Health Organization (WHO) state that 35% of women worldwide have experienced either physical and/or sexual intimate partner violence or non-partner sexual violence.[5] A pregnancy does not protect women from violence, and the prevalence of physical or sexual violence during pregnancy ranges from 3.4-11% in high-income countries.[6] It is recognised that violence has an adverse impact on women's physical, sexual, reproductive, and mental health.[5, 7]

A connection between PTB or LBW and violence against women has been reported, but the association is both supported and contradicted.[5, 8-20] Several pathways between sexual violence and adverse pregnancy outcomes are suggested.[5] A direct pathway of sexual violence can result in immediate complications as bleeding and rupture of membranes that can lead to a preterm birth.[5, 16] Other, more indirect pathways are suggested mediated by stress

1
2
3 and stress responses [18] or by behavioural factors such as smoking or substance abuse, used
4
5 to cope with the negative consequences of violence.[5]
6
7

8
9
10 Studies have primarily addressed physical abuse during pregnancy and PTB or LBW [8, 10,
11 11, 16-19, 21] or child sexual abuse and PTB/LBW.[9, 12] Results from a new meta-analysis
12
13 published in the recent WHO report [5] have demonstrated an association between intimate
14
15 partner violence, including both physical and sexual abuse, and PTB with an adjusted odds
16
17 ratio (AOR) of 1.41 (95% CI 1.21-1.62) and AOR of 1.16 (95% CI: 1.02-1.29) with LBW.[5]
18
19
20 However, studies that have investigated the impact of sexual violence on neonatal outcomes
21
22 specifically, are limited and few population-based studies with large sample sizes that enable
23
24 controlling for confounding variables have been conducted [5, 11] The Norwegian Mother
25
26 and Child Cohort Study (MoBa) is a population-based prospective cohort study of pregnant
27
28 women that includes measurements of lifetime sexual violence, sexual violence during
29
30 pregnancy and other relevant covariates which makes it suitable to examine associations
31
32 between sexual violence and neonatal outcome. In this study, we assessed the relationship
33
34 between sexual violence and gestational age at birth and birth weight. Additionally, we
35
36 explored the associations between sexual violence and PTB, LBW and SGA.
37
38
39
40
41

42 **METHODS**

43
44
45 This study was a sub-project in the MoBa study that was conducted by the Norwegian
46
47 Institute of Public health from 1999-2008.[22] All pregnant women in Norway were eligible
48
49 to participate in MoBa, and they were recruited during their routine fetal ultrasound
50
51 examination. Of the invited women, 40.6% consented to participate. Data were obtained
52
53 through extensive self-administered questionnaires that contained demographic factors,
54
55 general health, reproductive history and questions about maternal health status during
56
57
58
59
60

1
2
3 pregnancy. Our analyses were based on questionnaire 1 (Q1), which was completed during
4 (approximately) gestational week 17, and questionnaire 3 (Q3), which was completed during
5 (approximately) gestational week 30. Data from MoBa were linked with data from the
6
7
8 Medical Birth Registry of Norway (MBRN), which provided information on pregnancy and
9
10
11 birth outcome. The current study is based on version VI of the quality-assured data files
12
13 released for research in 2011. The MoBa study is described in detail elsewhere.[22] The
14
15 questionnaires that were used in MoBa are available at the internet-site:
16
17
18 <http://www.fhi.no/studier/den-norske-mor-og-barn-undersokelsen/sporreskjemaer>
19
20
21
22

23 **Study population**

24
25 This study included women who had a singleton birth between 22 and 44 weeks of gestation,
26
27 who completed both Q1 and Q3, had MBRN data available and participated for the first time
28
29 (79 363 women). While a pregnancy is the observation unit in the MoBa study, women are
30
31 the observation unit in our study; hence, the exclusion of 13 475 pregnancies of women who
32
33 participated more than once. We excluded 703 women who did not answer the questions on
34
35 sexual violence. Further, we excluded records with missing data on gestational length
36
37 (n= 297) and birth weight (n=41). We also excluded 6 children with birth weight <500 g and 4
38
39 children with birth weight >6000 g, leaving a study sample of 76 870 women (Figure 1).
40
41
42
43
44

45 **Ethical statement**

46
47 Informed consent was obtained from each women participating in the study. The Regional
48
49 Committee for Medical Research Ethics (Ref.SAFH 95/313 RTL) and the Norwegian Data
50
51 Inspectorate approved the study.
52
53
54
55
56
57
58
59
60

Variables

Exposure variable

The exposure variable was collected from Q1. Women were asked if they had been pressured or forced to sexual relations. There were four possible answer options: 1) No, never 2) Yes, pressured, 3) Yes, forced with violence or 4) Yes, raped. A positive answer was defined as having experienced sexual violence. Women with more than one positive answer were classified according to the most severe level reported. The answering options were coded into three levels of severity for the sexual violence: 1) mild (pressured), 2) moderate (forced with violence) and 3) severe (raped). Women could also indicate when the violence had taken place: 1) during this pregnancy, 2) during the last six month before pregnancy or 3) earlier. Approximately 1 700 women who filled out the first version of Q1 had the answering options earlier and during the last 12 month when assessing time. We therefore created the variables previous and recent sexual violence, recent containing sexual violence during last 12 month, including the current pregnancy. Among the women who participated several times we included the first pregnancy only to ensure that the exposure was included only once per woman. More details about the exposure variable can be found in our previous studies [23;24] and in supplementary table 1.

Outcome variables

All outcome variables were obtained from the MBRN. Gestational age at birth in days was based on ultrasound at (approximately) gestational week 18. For women with no ultrasound, the gestational age was based on the last menstrual period (1.7%). PTB was defined as a gestational age <37, LBW as a birth weight <2500 grams, and SGA was defined as birth weight below the 10th percentile for the gestational age at birth. SGA was calculated using Norwegian specific fetal growth tables by Skjerven et al.[25]

Adjusting variables

Maternal age, parity, socio-economical status, smoking and body mass index (BMI) were considered as possible confounding factors and were adjusted for. All adjusting variables were taken from the MoBa. In Q1, age was categorised into 5 groups: younger than 20 years, 20–24 years, 25–29 years, 30–34 years or 35 years and older. As a proxy for socio-economic status, we used the woman's education in years (categorised into 4 groups): primary (<12 years), secondary (12 years), higher ≤ 4 years (13–16 years) and higher >4 years (≥ 17 years). Parity was dichotomised into nulli- and multiparous women. Smoking was categorised as no smoking or smoking, which included both daily and occasional smoking. BMI was grouped into 4 categories: <20, 20-24.9, 25.0-29.9 or ≥ 30.0 kg/m². We also adjusted for mental distress because it is considered to be associated with both the exposure and the outcome.[4,5] Mental distress was measured using 5 items from The Hopkins symptoms checklist (SCL-5) with a cut-off at ≥ 2.0 points, as suggested by Strand [26] and obtained from Q3.

Because of the co-occurrence of different violence types,[7] we examined the effect of physical and emotional abuse as a child or as an adult in the multivariable statistical models. Information on adult physical violence was taken from Q1 and consisted of a positive answer to whether women as an adult had experienced being slapped, hit, kicked or otherwise bothered in a physical manner. Child physical violence was taken from Q3 and consisted of a positive answer to the question “Have you experienced physical violence before the age of 18?” Emotional abuse as a child (<18) or as an adult (≥ 18) consisted of a positive answer to either, “Someone has over a long period of time systematically tried to subdue, degrade or humiliate you” or “Someone has threatened to hurt you or someone close to you” or both.

1
2
3 Previous preterm birth and inadequate antenatal care are considered to be associated with the
4 exposure and the outcomes.[2, 3, 5] Because a previous preterm birth may be a result of
5 sexual violence prior to the related pregnancy, we did not control for a previous preterm birth.
6
7 In Norway, the majority of women attend antenatal care, a free and a well-integrated part of
8 the public health system; therefore, we did not control for antenatal care. Ethnicity was not
9 considered a relevant covariate in our study because the majority of the MoBa participants are
10 ethnic Norwegian.
11
12
13
14
15
16
17
18
19

20 21 **Statistical Analysis**

22 Characteristics were presented as percentages within the entire sample and the different
23 outcomes. Linear regression was performed to assess differences in birth weight and
24 gestational age for children born to women with and without a history of mild, moderate and
25 severe sexual violence. The association between sexual violence and PTB, LBW and SGA
26 was estimated with crude and adjusted odds ratios using logistic regression analysis. Sexual
27 violence was analysed as a categorical variable: 1=mild sexual violence, 2= moderate sexual
28 violence and 3=severe sexual violence with no sexual violence as the reference group. All
29 analyses were adjusted for maternal age, parity, education, smoking, BMI and mental distress
30 in the first step. Birth weight was additionally adjusted for gestational age. We further
31 adjusted for other types of violence in the second step. We initially tested the correlation
32 between other types of violence and sexual violence because of co-occurrence, and all
33 Pearson's correlation coefficients were below the generally accepted cut-off of <0.4 for use as
34 a covariate in the regression analyses.[27] Post protocol, we stratified the sample into
35 spontaneous start of birth and provider-initiated start of birth (induced start of birth or elective
36 caesarean section) for gestational age because a provider-initiated start could influence the
37 time point of birth. Information on how the birth started was taken from MBRN. We
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 additionally performed a sensitivity analysis in which we examined the association between
4 sexual violence and SGA and LBW among women who had a spontaneous birth at term (≥ 37
5 weeks) because we wanted to examine the effect of violence in a group of women who were
6 considered to be low risk according to gestational age and start of birth. When we examined
7 the timing of the sexual violence, we compared women who were exposed to recent sexual
8 violence (within the last 12 months) and those exposed to previous sexual violence to non-
9 abused women. We also examined the timing among women reporting recent and previous
10 severe sexual violence (rape) for all outcomes. The prevalence of missing data was generally
11 low with 2.5% for BMI, 3.7% for education, and 0.7% for smoking during pregnancy.
12 Because of this, no imputing methods for missing data were used,[28] except for the missing
13 data for the SCL-5 (3.2%), which were replaced by the series mean. The results of the logistic
14 regression analyses remained approximately the same when performed with the complete
15 exclusion of missing data compared with using the imputed missing data for SCL-5.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34 The comparison group for all analyses was women not reporting any sexual violence. All
35 analyses were performed with the statistical package SPSS for WINDOWS (SPSS Inc.,
36 Chicago, IL, USA) version 18. P-values < 0.05 were considered statistically significant.
37
38
39
40
41
42

43 RESULTS

44
45 The mean gestational age at birth in the total sample was 279.6 days (standard deviation 11.9
46 days), and the mean birth weight was 3592.7 grams (standard deviation 547.1 g). Table 1
47 displays the characteristics in the total sample and by the different outcomes.
48
49
50

51 Table 1. Background characteristics in the total sample and by preterm birth (PTB): gestational age < 37 ,
52 low birth weight (LBW): weight < 2500 g and small for gestational age (SGA) weight below the 10th percentile
53 by gestational age at birth in the Mother and Child Cohort
54

	Total	PTB	LBW	SGA
	N=76 870	N=3620	N=2107	N=6257
	%	%	%	%

1					
2					
3	Age				
4	<20	1.5	2.2	2.6	1.9
5	20–24	12.2	12.7	13.8	13.2
6	25–29	36.2	34.6	34.8	36.4
7	30–34	35.7	33.7	33.2	34.2
8	≥35	14.4	16.7	15.6	14.3
9					
10					
11	Education				
12	Primary	2.4	3.1	3.9	2.7
13	Secondary	34.8	38.5	38.8	35.0
14	Higher ≤4 years	37.8	35.6	37.3	36.1
15	Higher >4 years	21.3	19.2	20.5	22.2
16	Missing	3.7	3.6	3.7	3.9
17					
18					
19	Parity				
20	Nulliparous	54.9	61.7	68.2	70.7
21	Multiparous	45.1	38.3	31.8	29.3
22					
23	Smoking				
24	No	90.9	89.3	85.8	85.6
25	Yes	8.5	10.1	13.6	13.6
26	Missing	0.7	0.6	0.6	0.7
27					
28					
29	Pre-pregnancy body mass				
30	index				
31	<20	12.4	12.8	15.6	19.1
32	20–24.9	54.9	49.9	50.8	55.6
33	25–29.9	21.0	22.0	19.0	15.2
34	≥30	9.2	12.6	11.8	7.6
35	Missing	2.5	2.7	2.9	2.4
36					
37					
38	Mental distress				
39	No	93.3	92.3	91.3	92.3
40	Yes	6.8	7.7	8.7	7.7
41					
42	Adult physical violence				
43	No	85.5	83.9	83.6	84.7
44	Yes	14.5	16.1	16.4	15.3
45					
46	Child physical violence				
47	No	94.5	94.4	94.1	94.3
48	Yes	5.5	5.6	5.9	5.5
49					
50	Adult emotional abuse				
51	No	83.6	82.1	81.3	82.5
52	Yes	16.4	17.9	18.7	17.5
53					
54	Child emotional abuse				
55	No	86.2	85.1	84.7	85.5
56					
57					
58					
59					
60					

Yes	13.8	14.9	15.3	14.5
-----	------	------	------	------

The prevalence of adverse neonatal outcomes was generally highest in the youngest (<20 y) and the oldest age groups (≥ 35 y) among smokers and women with primary school education. A BMI ≥ 30 was associated with PTB and LBW and BMI <20 with SGA. Women who reported mental distress also reported more PTB, LBW and SGA.

Among the 76 870 women enrolled, 9263 (12.1%) reported a history of mild sexual violence, 2102 (2.8%) moderate and 2746 (3.5%) severe. Women with a history of sexual violence were significantly younger and they were more likely to have primary school education. Additionally, these women more frequently reported smoking, a BMI ≥ 30 and mental distress. These women more often experienced other types of violence both as children and adults (data not provided in tables).

A lower gestational age at birth was observed for newborns from women who reported moderate and severe sexual violence with approximately two days when birth was provider-initiated (Table 2).

Table 2. Differences in gestational age for spontaneous and provider-initiated start of birth and birth weight for non-exposed women and women exposed to mild, moderate and severe sexual violence

	N= 76 870 (%)	Mean	Crude estimate β (95% CI)	Adjusted estimate β (95% CI)*
Gestational age				
No sexual violence	62 699 (81.6)	279.7 days	1	1
<i>Mild sexual violence</i>				
Spontaneous start	7323 (11.5)	279.9 days	0.09 (-0.16; 0.33)	0.22 (-0.04; -0.48)
Provider-initiated start	1940 (12.5)	280.3 days	0.36 (-0.37; 1.08)	0.64 (-0.11; 1.39)
<i>Moderate sexual violence</i>				
Spontaneous start	1670 (2.7)	279.3 days	-0.40 (-0.87; 0.06)	-0.28 (-0.78; 0.21)
Provider-initiated start	492 (3.2)	277.6 days	-2.13 (-3.41; -0.84)	-2.02 (-3.39; -0.67)
<i>Severe sexual violence</i>				
Spontaneous start	2048 (3.3)	278.7 days	-0.91 (-1.37; -0.44)	-0.65 (-1.15; -0.16)

Provider-initiated start	698 (4.5)	277.5 days	-2.24 (-3.47; -1.01)	-1.92 (-3.22; -0.62)
Birth weight				
No sexual violence	62 699 (81.6)	3594 g	1	1
Mild sexual violence	9263 (12.1)	3597 g	2.03 (-9.26; 13.32)	0.04 (-9.16; 9.23)**
Moderate sexual violence	2162 (2.8)	3582 g	-13.61 (-34.74; 7.51)	6.11 (-11.30; 23.52)**
Severe sexual violence	2746 (3.6)	3556 g	-38.33 (-59.17; -17.49)	-0.76 (-18.05; 16.53)**

*Adjusted for maternal age, parity, education, smoking, body mass index and mental distress

**Additional adjustment for gestational age

Among women with a spontaneous start of birth, the gestational age was approximately one half of a day shorter when women reported severe sexual violence. These findings were significant in an adjusted analysis. A crude analysis showed that women who reported a history of severe violence delivered on average 38.3 g lighter children, a difference that disappeared when controlling for gestational age, mother's age, parity, education, smoking, BMI and mental distress. There were no differences regarding birth weight between women with a history of mild or moderate sexual violence compared with non-abused women.

Results from the logistic regression analysis are presented in Table 3.

Table 3. Odds of preterm birth (PTB), low birth weight (LBW) and small for gestational age (SGA) with 95% confidence intervals (CI) according to the different levels of sexual violence

	N (%)	Prevalence (%)	Crude odds ratio (95% CI)	Adjusted odds ratio (95% CI)*
PTB		3620 (4.7)		
No sexual violence	62 699 (81.6)	2931 (4.7)	1	1
Mild sexual violence	9263 (12.1)	412 (4.4)	0.95 (0.85-1.06)	0.93 (0.83-1.03)
Moderate sexual violence	2162 (2.8)	115 (5.3)	1.15 (0.95-1.39)	1.14 (0.93-1.39)
Severe sexual violence	2746 (3.6)	162 (5.9)	1.28 (1.08-1.51)	1.15 (0.97-1.37)
LBW		2107 (2.7)		
No sexual violence	62 699 (81.6)	1681 (2.7)	1	1
Mild sexual violence	9263 (12.1)	257 (2.8)	1.04 (0.91-1.18)	0.98 (0.85-1.12)
Moderate sexual violence	2162 (2.8)	75 (3.5)	1.30 (1.03-1.65)	1.19 (0.93-1.53)
Severe sexual violence	2746 (3.6)	94 (3.4)	1.29 (1.04-1.59)	1.07 (0.85-1.34)
SGA		6257 (8.1)		
No sexual violence	62 699 (81.6)	5061 (8.1)	1	1
Mild sexual violence	9263 (12.1)	768 (8.3)	1.03 (0.95-1.12)	1.00 (0.91-1.08)
Moderate sexual violence	2162 (2.8)	178 (8.3)	1.02 (0.87-1.19)	0.95 (0.80-1.12)

Severe sexual violence	2746 (3.6)	250 (9.1)	1.14 (1.00-1.30)	1.05 (0.91-1.21)
------------------------	------------	-----------	------------------	------------------

*Adjusted for maternal age, parity, education, smoking, body mass index and mental distress

Women who reported severe sexual violence had higher odds of PTB, LBW and SGA in a crude analysis, an association that was attenuated and no longer significant when adjusted for maternal age, parity, education, smoking, BMI and mental distress. Other types of violence, both as a child and adult, had small attenuating effects on the odds ratios and were not included in the final models.

The sensitivity analysis, in which we examined the association between a history of sexual violence and SGA and LBW in a sub-sample of women who had a spontaneous term birth, showed the same pattern as in the total sample reported in Table 3. Women who reported severe sexual violence had higher odds of LBW and SGA in a crude analysis but not in the adjusted analysis (Data not provided in tables).

A crude analysis was used to examine if the timing of the violence was associated with adverse outcome. Women who reported recent sexual violence had a higher risk for LBW (OR 1.60 95% CI 1.04-2.17) compared with non-abused women. The association was no longer significant in the adjusted analysis. In our study, 684 (0.9%) women reported recent sexual violence (mild, moderate and severe) and 13 487 (17.5%) previous sexual violence (Supplementary Table S2). There was no association between recent severe sexual violence (rape) and adverse neonatal outcome (Supplementary Table S3). There were 66 (0.1%) women who reported recent rape in this study.

DISCUSSION

Main outcome

We found that moderate and severe sexual violence were associated with a reduction in gestational age at birth. The largest effect was observed when birth was provider-initiated among women exposed to moderate or severe violence. These women had an approximately two-day reduction in gestational age. There was no significant association between sexual violence and PTB, LBW or SGA in the adjusted analysis.

Strength

This study, based on information from a large population-based study, the Norwegian Mother and Child Cohort study (MoBa), which is linked to the Medical Birth Registry (MBRN), gave a unique opportunity to assess the association between sexual violence and outcome for newborns. The validity of the data in MoBa has in earlier research been described as sufficient for large-scale epidemiologic studies.[29, 30] Our study was strengthened by the fact that the information on the different outcome variables was collected prospectively from the quality-assessed MBRN.[31] The outcomes in this study are part of a complex phenomenon that has several different risk factors.[2] The setting in this study, with small social and health inequalities, may therefore be suitable to isolate the effect of sexual violence on adverse neonatal outcomes.

Limitations

There are also limitations to our study. The participation rate of 40.6% in MoBa is low, and MoBa suffers to some extent from selection bias. The women included in the study are older, have more education, smoke less and are less likely to be of a non-Norwegian origin than the Norwegian population. Although it is likely that there is a socioeconomic gradient that

1
2
3 influence prevalence estimates, a recent study by Nilsen et al. found no evidence that the
4 exposure-outcome associations in the MoBa study were affected by selection bias.[32] This
5 socioeconomic gradient may also limit the generalisability of our findings. The lack of a
6 validated instrument for measuring the exposure is a limitation to this study, and violence
7 measured in modules as part of a larger questionnaire, as that in MoBa, may achieve a lower
8 disclosure rate.[5] However, a similar prevalence to that found in our study was reported in a
9 Nordic study examining sexual violence and health.[33] The exposure was measured in
10 gestational week 17, and sexual violence during pregnancy after this is therefore not included.
11
12 In addition, we have no information on the context and frequency of the violence or
13 information regarding the perpetrator.
14
15
16
17
18
19
20
21
22
23
24
25
26

27 To our knowledge, no studies have examined the influence of lifetime sexual violence
28 reported during pregnancy on the gestational age at birth for newborns. There were minor
29 differences in the gestational age between abused and non-abused women in this study, and
30 the clinical importance of our findings for the health of the newborn is most likely limited.
31
32 However, the difference between the provider-initiated and spontaneous initiation of birth
33 may be of interest. Shorter provider-initiated pregnancies may suggest an increase in elective
34 inductions and elective caesarean sections for those exposed to violence. This is supported by
35 others [34, 35] and in our previous study on sexual violence and maternal outcome.[36]
36
37 Studies have emphasised the meaning of control for abused women when giving birth,[37]
38 and choosing a planned start of birth may help the abused women remain in control.
39
40
41
42
43
44
45
46
47
48
49
50

51 Unlike the current meta-analysis (not yet published), mentioned in a WHO report [5] we did
52 not find sexual violence to be associated with PTB and LBW in adjusted analysis. To our
53 knowledge, no study has found an association between violence and SGA. The studies
54
55
56
57
58
59
60

1
2
3 included in the meta-analysis were limited to sexual and/or physical intimate partner violence.
4
5 However, our findings are supported by a Canadian population based study with a sample of 6
6
7 421 pregnant women [17] and a prospective cohort study including 1 555 women from the
8
9 US.[15] The exposure in these studies was physical and sexual violence prior to pregnancy
10
11 and in pregnancy without being limited to an intimate partner.[15, 17] These studies showed
12
13 no association between violence and PTB or LBW.
14
15

16
17
18 The nature of the exposure measured in the MoBa study makes it difficult to directly compare
19
20 our findings to others, mainly because we examined lifetime sexual violence by any
21
22 perpetrator, not limited to intimate partner. Sexual violence is considered to be traumatic for
23
24 the victim regardless if the perpetrator is a partner or not.[38] When an intimate partner is the
25
26 perpetrator, sexual violence may in addition be accompanied by controlling behaviour and
27
28 include both physical and emotional abuse.[38] Rape by strangers is usually a single violent
29
30 event with a higher risk of physical injury. Both forms of violence are associated with adverse
31
32 health effects,[5, 38] but the effect may differ. Unfortunately, we were not able to examine the
33
34 effect of the perpetrator in this study because the MoBa study does not provide this
35
36 information. However, research suggests that a substantial proportion of sexual violence
37
38 occurs within an intimate relationship.[38] It is not unlikely that the question about severe
39
40 sexual violence (rape) primarily reflects non-partner sexual abuse and that mild sexual
41
42 violence (pressured to sexual acts) may be a more psychological exposure. Our crude analyses
43
44 showed different results, with a significantly higher OR for adverse neonatal outcome among
45
46 women who reported severe sexual violence, thus supporting the idea that the different levels
47
48 of violence are different exposures. The use of a more comprehensive instrument when
49
50 measuring the exposure, with multiple response options regarding context, frequency and
51
52 perpetrator would have clarified this further and provided more comprehensive knowledge
53
54
55
56
57
58
59
60

1
2
3 about the nature of the violence. Nevertheless, we had the opportunity to control for both
4
5 emotional and physical abuse in preliminary analysis and this did not change the ORs.
6
7

8
9
10 The prevalence of abuse during pregnancy is small in our study and may have decreased the
11
12 power to detect an association between violence during pregnancy and adverse neonatal
13
14 outcomes, an association that is supported by other studies.[10, 14, 39, 40] Living in an
15
16 abusive relationship may have stopped women from disclosing the violence. In our study,
17
18 sexual violence was assessed approximately in gestational week 17, and events of violence
19
20 after that have been missed. Some studies suggest that the risk of sexual violence may
21
22 increase with the length of the pregnancy for women that are exposed.[41] The studies that
23
24 report the highest prevalence of violence have measured this several times during
25
26 pregnancy.[6] Studies have reported a violence prevalence during pregnancy of between 3 to
27
28 19%.[6, 42], including both physical and sexual violence. Underreporting among the non-
29
30 exposed may have caused a misclassification that has diminished the associations between
31
32 sexual violence and neonatal outcomes in our study. Since the exposure was collected before
33
34 the outcome, it is unlikely that misclassification was related to the outcome, thus resulting in a
35
36 nondifferential misclassification that has biased the result towards the null. However, our
37
38 prevalence of recent sexual (0.9%) violence is similar to the prevalence of 1% reported in a
39
40 survey that assessed intimate partner violence among a representative sample of Norwegian
41
42 women.[43] This number reflects the prevalence of sexual violence reported during the last
43
44
45
46
47 year.
48

49
50
51 Several pathways between sexual violence and adverse neonatal outcomes are suggested, both
52
53 direct and indirect.[5] The direct pathway of violence during pregnancy can cause immediate
54
55 complications as bleeding, rupture of membranes and PTB, and examples of indirect
56
57
58
59
60

1
2
3 pathways include more health-risk behaviours, depression and stress/anxiety.[5, 44] Both
4
5 experienced violence and living in an abusive environment can cause increased stress levels,
6
7 which could be on the pathway between abuse and adverse neonatal outcome. Maternal
8
9 exposure to stress can influence the hypothalamic pituitary axis (HPA) hormones, and it is
10
11 suggested that changes in these hormones may cause negative outcomes, such as a reduction
12
13 in gestational age and fetal growth restriction.[4, 45] It has been proposed that mental distress
14
15 and symptoms of depression are on the causal pathway between violence and adverse health
16
17 outcome, yet it has also been suggested that women with mental health difficulties are more
18
19 likely to be victims of violence.[46] Because the relationship may be bi-directional, we chose
20
21 to control for mental distress in our study. Similarly, the health risk behaviours smoking and
22
23 BMI may be on the pathway between sexual violence and neonatal outcome. Nevertheless, we
24
25 kept these covariates in the regression analysis because they are especially related to birth
26
27 weight and PTB.[3, 47]
28
29
30
31
32
33

34 **Conclusion**

35
36 Overall, our findings provide no evidence for an association between lifetime sexual violence
37
38 and adverse neonatal outcomes. A small significant effect on the gestational age at birth was
39
40 detected, but the clinical importance of this is most likely limited for the health of the
41
42 newborn. PTB, LBW and SGA all have complex origins with multiple possible pathways.[1,
43
44 2] Although we did not find an association between sexual violence and PTB, LBW or SGA
45
46 in adjusted analyses, crude analyses in our study suggested that sexual violence may be a risk
47
48 factor for adverse neonatal outcomes for some women; however, for the majority of women,
49
50 the relationship was confounded by other risk factors. It is possible that these factors were the
51
52 result of prior exposure to violence, but this could not be assessed in this study. Antenatal care
53
54 is one of the most important entry points in which women seek health care without necessarily
55
56
57
58
59
60

1
2
3 disclosing ongoing exposure to violence or a history of sexual violence. It is recommended
4
5 that caregivers and clinicians ask women about exposure to violence when assessing
6
7 conditions that may be complicated by violence,[48] and more knowledge about this
8
9 conditions is needed. Additionally, antenatal care may offer opportunities for women to
10
11 receive help, both if they are exposed to violence and in providing assistance to change
12
13 behavioural factors contributing to adverse outcomes.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgement

We are grateful to all the participating families in Norway who take part in this ongoing cohort study.

Contributions to authorship

LH participated in the conception and design of the study, performed the analysis and drafted the manuscript. BS participated in the conception of the study, advised on the statistical analyses and drafting of the manuscript. SV advised on the statistical analyses and the drafting of the manuscript. ML participated in the conception and design of the study, advised on the statistical analyses and participated in drafting the manuscript. All authors read and approved the final version.

Funding

The Norwegian Mother and Child Cohort Study is supported by the Norwegian Ministry of Health and the Ministry of Education and Research, NIH/NIEHS (contract no N01-ES-75558), NIH/NINDS (grant no.1 U01 NS 047537-01 and grant no.2 U01 NS 047537-06A1), and the Norwegian Research Council/FUGE (grant no. 151918/S10). The first author of this article is supported by South-Eastern Norway Regional Health Authority (grant no. 2012010) and the second author is supported by the The Research Council of Norway (grant no. 204292/H10).

Data Sharing Statement

No additional data are available.

Competing interests

None.

Reference List

- (1) Green NS, Damus K, Simpson JL, et al. Research agenda for preterm birth: recommendations from the March of Dimes. *Am J Obstet Gynecol* 2005;193:626-35.
- (2) March of Dimes, PMNCH, Save The Children, WHO. Born Too Soon: The Global Action Report on Preterm Birth. Geneva: World Health Organization 2012
- (3) Wilcox AJ. On the importance-and the unimportance-of birth weight. *Int J Epidemiol* 2001;30:1233-41.
- (4) Lazinski MJ, Shea AK, Steiner M. Effects of maternal prenatal stress on offspring development: a commentary. *Arch Womens Ment Health* 2008;11:363-75.
- (5) Garcia-Moreno C, Pallitto C, Devries K, et al. Global and regional estimates of violence against women: prevalence and health effects of intimate partner violence and non-partner sexual violence. Geneva: World Health Organization; 2013.
- (6) Campbell J, Garcia-Moreno C, Sharps P. Abuse During Pregnancy in Industrialized and Developing Countries. *Violence Against Women* 2004;10:770-89.
- (7) Garcia-Moreno C, Jansen HA, Ellsberg M, et al. Prevalence of intimate partner violence: findings from the WHO multi-country study on women's health and domestic violence. *Lancet* 2006;368:1260-9.
- (8) Sanchez SE, Alva AV, Diez CG, et al. Risk of spontaneous preterm birth in relation to maternal exposure to intimate partner violence during pregnancy in Peru. *Matern Child Health J* 2013;17:485-92.
- (9) Leeners B, Stiller R, Block E, et al. Pregnancy complications in women with childhood sexual abuse experiences. *J Psychosom Res* 2010;69:503-10.
- (10) Nunes MA, Camey S, Ferri CP, et al. Violence during pregnancy and newborn outcomes: a cohort study in a disadvantaged population in Brazil. *Eur J Public Health* 2011;21(1):92-7.
- (11) Rodrigues T, Rocha L, Barros H. Physical abuse during pregnancy and preterm delivery. *Am J Obstet Gynecol* 2008;198:171-6.
- (12) Noll JG, Schulkin J, Trickett PK, et al. Differential pathways to preterm delivery for sexually abused and comparison women. *J Pediatr Psychol* 2007;32:1238-48.
- (13) Silverman JG, Decker MR, Reed E, et al. A. Intimate partner violence victimization prior to and during pregnancy among women residing in 26 U.S. states: associations with maternal and neonatal health. *Am J Obstet Gynecol* 2006;195:140-8.
- (14) Murphy CC, Schei B, Myhr TL, et al. Abuse: a risk factor for low birth weight? A systematic review and meta-analysis. *CMAJ* 2001;164:1567-72.
- (15) Fried LE, Cabral H, Amaro H, et al. Lifetime and during pregnancy experience of violence and the risk of low birth weight and preterm birth. *J Midwifery Womens Health* 2008 ;53:522-8.

- 1
2
3 (16) Janssen PA, Holt VL, Sugg NK, et al. Intimate partner violence and adverse
4 pregnancy outcomes: a population-based study. *Am J Obstet Gynecol* 2003;188:1341-
5 7.
- 6
7 (17) Urquia ML, O'Campo PJ, Heaman MI, et al. Experiences of violence before and
8 during pregnancy and adverse pregnancy outcomes: an analysis of the Canadian
9 Maternity Experiences Survey. *BMC Pregnancy Childbirth* 2011;11:42.
- 10
11 (18) Altarac M, Strobino D. Abuse during pregnancy and stress because of abuse during
12 pregnancy and birthweight. *J Am Med Womens Assoc* 2002;57:208-14.
- 13
14 (19) Cokkinides VE, Coker AL, Sanderson M, et al. Physical violence during pregnancy:
15 maternal complications and birth outcomes. *Obstet Gynecol* 1999;93:661-6.
- 16
17 (20) Grimstad H, Schei B, Backe B, et al. Physical abuse and low birthweight: a case-
18 control study. *Br J Obstet Gynaecol* 1997;104:1281-7.
- 19
20 (21) Kaye DK, Mirembe FM, Bantebya G, et al. Domestic violence during pregnancy and
21 risk of low birthweight and maternal complications: a prospective cohort study at
22 Mulago Hospital, Uganda. *Trop Med Int Health* 2006;11:1576-84.
- 23
24 (22) Magnus P, Irgens LM, Haug K, et al. Cohort profile: the Norwegian Mother and Child
25 Cohort Study (MoBa). *Int J Epidemiol* 2006;35:1146-50.
- 26
27 (23) Lukasse M, Henriksen L, Vangen S, et al. Sexual violence and pregnancy-related
28 physical symptoms. *BMC Pregnancy Childbirth* 2012;12:83.
- 29
30 (24) Henriksen L, Vangen S, Schei B, et al. Sexual violence and antenatal hospitalization.
31 *Birth* 2013;40:281-8.
- 32
33 (25) Skjaerven R, Gjessing HK, Bakketeig LS. et al. Birthweight by gestational age in
34 Norway. *Acta Obstet Gynecol Scand* 2000;79:440-9.
- 35
36 (26) Strand BH, Dalgard OS, Tambs K, et al. Measuring the mental health status of the
37 Norwegian population: a comparison of the instruments SCL-25, SCL-10, SCL-5 and
38 MHI-5 (SF-36). *Nord J Psychiatry* 2003;57:113-8.
- 39
40 (27) Field A. Discovering statistics using SPSS. Los Angeles: SAGE; 2009.
- 41
42 (28) Veierød MB, Lydersen S, Laake P, et at. Medical statistics: in clinical and
43 epidemiological research. Oslo: Gyldendal akademisk; 2012.
- 44
45 (29) Kvalvik LG, Nilsen RM, Skjaerven R, et al. Self-reported smoking status and plasma
46 cotinine concentrations among pregnant women in the Norwegian Mother and Child
47 Cohort Study. *Pediatr Res* 2012 ;72:101-7.
- 48
49 (30) Furu K, Karlstad O, Skurtveit S, et al. High validity of mother-reported use of
50 antiasthmatics among children: a comparison with a population-based prescription
51 database. *J Clin Epidemiol* 2011;64:878-84.
- 52
53 (31) Irgens LM. The Medical Birth Registry of Norway. Epidemiological research and
54 surveillance throughout 30 years. *Acta Obstet Gynecol Scand* 2000;7:435-9.
- 55
56
57
58
59
60

- 1
2
3 (32) Nilsen RM, Vollset SE, Gjessing HK, et al. Self-selection and bias in a large
4 prospective pregnancy cohort in Norway. *Paediatr Perinat Epidemiol* 2009 23:597-
5 608.
6
7 (33) Hilden M, Schei B, Swahnberg K, et al. A history of sexual abuse and health: a Nordic
8 multicentre study. *BJOG* 2004;111:1121-7.
9
10 (34) Nerum H, Halvorsen L, Straume B, et al. Different labour outcomes in primiparous
11 women that have been subjected to childhood sexual abuse or rape in adulthood: a
12 case-control study in a clinical cohort. *BJOG* 2013;120:487-495.
13
14 (35) Lukasse M, Vangen S, Oian P, et al. Childhood abuse and caesarean section among
15 primiparous women in the Norwegian Mother and Child Cohort Study. *BJOG*
16 2010;117:1153-7.
17
18 (36) Henriksen L, Schei B, Vangen S, et al. Sexual violence and mode of delivery: a
19 population-based cohort study. *BJOG* 2014;12:1237-1244
20
21 (37) Montgomery E. Feeling Safe: A Metasynthesis of the Maternity Care Needs of
22 Women Who Were Sexually Abused in Childhood. *Birth* 2013;40:88-95.
23
24 (38) Abrahams N, Devries K, Watts C, et al. Worldwide prevalence of non-partner sexual
25 violence: a systematic review. *Lancet*. 2014;383(9929):1648-54. Epub 2014/02/18.
26
27
28
29 (39) Faramarzi M, Esmaelzadeh S, Mosavi S. Prevalence, maternal complications and birth
30 outcome of physical, sexual and emotional domestic violence during pregnancy. *Acta*
31 *Med Iran* 2005;43:115-22.
32
33 (40) Valladares E, Ellsberg M, Pena R, et al. Physical partner abuse during pregnancy: a
34 risk factor for low birth weight in Nicaragua. *Obstet Gynecol* 2002;100:700-5.
35
36 (41) Shah PS, Shah J. Maternal exposure to domestic violence and pregnancy and birth
37 outcomes: a systematic review and meta-analyses. *J Womens Health (Larchmt)*.
38 2010;19:2017-31.
39
40 (42) Sharps PW, Laughon K, Giangrande SK. Intimate partner violence and the
41 childbearing year: maternal and infant health consequences. *Trauma Violence Abuse*
42 2007;8:105-16.
43
44 (43) Neroien AI, Schei B. Partner violence and health: results from the first national study
45 on violence against women in Norway. *Scand J Public Health* 2008;36:161-8.
46
47 (44) Leeners B, Neumaier-Wagner P, Quarg AF, et al. Childhood sexual abuse (CSA)
48 experiences: an underestimated factor in perinatal care. *Acta Obstet Gynecol Scand*
49 2006;85:971-6.
50
51 (45) Valladares E, Pena R, Ellsberg M, et al. Neuroendocrine response to violence during
52 pregnancy-impact on duration of pregnancy and fetal growth. *Acta Obstet Gynecol*
53 *Scand* 2009;88:818-23.
54
55
56
57
58
59
60

- 1
2
3 (46) Devries KM, Mak JY, Bacchus LJ, et al. Intimate partner violence and incident
4 depressive symptoms and suicide attempts: a systematic review of longitudinal
5 studies. *PLoS Med* 2013;10:e1001439
6
7 (47) McDonald SD, Han Z, Mulla S, Beyene J. Overweight and obesity in mothers and risk
8 of preterm birth and low birth weight infants: systematic review and meta-analyses.
9 *BMJ* 2010;341:3428.
10
11 (48) World health organisation. Responding to intimate partner violence and sexual
12 violence against women: *WHO clinical and policy guidelines*. Geneva: WHO; 2013.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TITLE PAGE

Title: Sexual violence and neonatal outcomes: a [Norwegian](#) population-based cohort study

Lena Henriksen¹, Berit Schei^{2,3}, Siri Vangen⁴, Mirjam Lukasse⁵

Corresponding author: Lena Henriksen¹, Section of Obstetrics, Department for Women's and Children's Health, Oslo University Hospital, Postboks 4950 Nydalen, N-0424 Oslo, Norway.

E-mail: uxhelz@ous-hf.no Phone: +47 90068485.

Co-authors: Berit Schei², Department of Public Health and General Practice, Faculty of Medicine, The Norwegian University of Science and Technology (NTNU), Håkon Jarls gate 11, N-7489 Trondheim, Norway. ³Department of Obstetrics and Gynecology, St.Olavs Hospital, Trondheim University Hospital, Postbox 3250 Sluppen, N-7006 Trondheim, Norway.

Siri Vangen⁴, National Resource Centre for Women's Health, Department for Women's and Children's Health, Oslo University Hospital, Postboks 4950 Nydalen, N-0424 Oslo, Norway.

Mirjam Lukasse⁵, Department of Health, Nutrition and Management, Faculty of Health Sciences, Oslo and Akershus University College of Applied Sciences, Postboks 4 St. Olavs plass, N-0130 Oslo, Norway.

Keywords: Sexual violence, rape, premature birth, low birth weight, small for gestational age, MoBa.

Word count: [42493644](#)

ABSTRACT

Objective: The objective of this study was to explore the association between sexual violence and neonatal outcomes.

Design: National cohort study.

Setting: Women were recruited to the Norwegian Mother and Child Cohort Study (MoBa) when attending routine ultrasound examinations from 1999-2008.

Population: A total of 76 870 pregnant women.

Methods: Sexual violence and maternal characteristics were self-reported in postal questionnaires during pregnancy. Neonatal outcomes were retrieved from the Medical Birth Registry of Norway (MBRN). Risk estimations were performed with linear and logistic regression analysis.

Outcome measures: Gestational age at birth, birth weight, preterm birth (PTB), low birth weight (LBW) and small for gestational age (SGA).

Results: Of 76 870 women, 18.4% reported a history of sexual violence. A total of 4.7% had a premature birth, 2.7% had children with a birth weight <2500 g and 8.1% children were small for gestational age. Women reporting both moderate and severe sexual violence (rape) had a significantly reduced gestational length (2 days) when the birth was provider-initiated in an analysis adjusted for age, parity, education, smoking, BMI and mental distress. Those exposed to severe sexual violence had a significantly reduced gestational length of 0.51 days with a spontaneous start of birth. Crude estimates showed that severe sexual violence was associated with PTB, LBW and SGA. When controlling for the above-mentioned socio-demographic and behavioural factors, the association was no longer significant.

Conclusions: Sexual violence was not associated with adverse neonatal outcomes. Moderate and severe violence had a small but significant effect on gestational age; however, the clinical influence of this finding is most likely limited. Women exposed to sexual violence in this study reported more of the socio-demographic and behavioural factors associated with PTB, LBW and SGA compared with non-abused women.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- This study was based on information from a large population-based study described as sufficient for large-scale epidemiologic studies.
- All outcomes were collected prospectively from a quality-assessed birth register.
- The setting, with small social and health inequalities, was suitable to isolate the effect of sexual violence on adverse neonatal outcomes.
- A non-validated instrument for measuring of the exposure variable was a limitation to this study.

INTRODUCTION

Preterm birth is a common and costly health problem.[1, 2] Approximately one in ten babies are born preterm worldwide, and prematurity is considered to be the leading cause of death for newborns.[2] Low birth weight (LBW) can be a consequence of preterm birth (PTB) or intra uterine growth restriction, the latter leading to the birth of small for gestational age (SGA) infants.[3] Research has ~~suggested identified~~ some biological risk factors for PTB and LBW: ~~ethnicity~~, multiple pregnancies, a previous preterm birth and uterine or placental abnormalities.[1, 3] Studies also emphasise ~~other, less understood behavioural and social factors as important but less understood~~ factors for both PTB and LBW.[1] These ~~are factors can be~~ maternal age, socio-economic status, ~~ethnicity~~, maternal weight, substance abuse, stress, depression and violence.[1, 2, 4]

Violence against women is a significant public health problem, and a recent report from the World Health Organization (WHO) state that 35% of women worldwide have experienced either physical and/or sexual intimate partner violence or non-partner sexual violence.[5] A pregnancy does not protect women from violence, and the prevalence of physical or sexual violence during pregnancy ranges from 3.4-11% in high-income countries.[6] It is recognised that violence has an adverse impact on women's physical, sexual, reproductive, and mental health.[5, 7]

A connection between PTB or LBW and violence against women has been reported, but the association is both supported and contradicted.[5, 8-20] ~~Several pathways between sexual violence and adverse pregnancy outcomes are suggested.~~ [5] ~~A direct pathway of sexual violence can result in immediate complications as bleeding and rupture of membranes that can~~

1
2
3 lead to a preterm birth.[5, 16] Other, more indirect pathways are suggested mediated by stress
4 and stress responses [18] or by behavioural factors such as smoking or substance abuse, used
5 to cope with the negative consequences of violence.[5]
6
7
8
9

10
11 Studies have primarily addressed physical abuse during pregnancy and PTB or LBW [8, 10,
12 11, 16-19, 21] or child sexual abuse and PTB/LBW.[9, 12] Results from a new meta-analysis
13 published in the recent WHO report [5] have demonstrated an association between intimate
14 partner violence, including both physical and sexual abuse, and PTB with an adjusted odds
15 ratio (AOR) of 1.41 (95% CI 1.21-1.62) and AOR of 1.16 (95% CI: 1.02-1.29) with LBW.[5]
16
17

18
19 However, ~~studies on sexual violence and neonatal outcomes~~ studies that have investigated the
20 impact of sexual violence on neonatal outcomes specifically, are limited and few population-
21 based studies with large sample sizes that enable controlling for confounding variables have
22 been conducted [5, 11] ~~Thus, there is a need to study the potential effect of lifetime sexual~~
23 ~~violence on outcomes for newborns using a large cohort.~~ The Norwegian Mother and Child
24 Cohort Study (MoBa) is a population-based prospective cohort study of pregnant women that
25 includes measurements of lifetime sexual violence, sexual violence during pregnancy and
26 other relevant covariates which makes it suitable to examine associations between sexual
27 violence and neonatal outcome. In this study, we assessed the relationship between sexual
28 violence and gestational age at birth and birth weight. Additionally, we explored the
29 associations between sexual violence and PTB, LBW and SGA.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

49 **METHODS**

50
51 This study was a sub-project in the MoBa -study that was conducted by the Norwegian
52 Institute of Public health from 1999-2008.[22] All pregnant women in Norway were eligible
53 to participate in MoBa, and they were recruited during their routine foetal ultrasound
54
55
56
57
58
59
60

1
2
3 examination. Of the invited women, 40.6% consented to participate. Data were obtained
4
5 through extensive self-administered questionnaires that contained demographic factors,
6
7 general health, reproductive history and questions about maternal health status during
8
9 pregnancy. Our analyses were based on questionnaire 1 (Q1), which was completed during
10
11 (approximately) gestational week 17, and questionnaire 3 (Q3), which was completed during
12
13 (approximately) gestational week 30. Data from MoBa were linked with data from the
14
15 Medical Birth Registry of Norway (MBRN), which provided information on pregnancy and
16
17 birth outcome. The current study is based on version VI of the quality-assured data files
18
19 released for research in 2011. The MoBa study is described in detail elsewhere.[22] [The](#)
20
21 [questionnaires that were used in MoBa are available at the internet-site:](#)
22
23 <http://www.fhi.no/studier/den-norske-mor-og-barn-undersokelsen/sporreskjemaer>
24
25
26
27
28
29
30
31
32
33

34 **Study population**

35
36 This study included women who had a singleton birth between 22 and 44 weeks of gestation,
37
38 who completed both Q1 and Q3, had MBRN data available and participated for the first time
39
40 (79 363 women). While a pregnancy is the observation unit in the MoBa study, women are
41
42 the observation unit in our study; hence, the exclusion of 13 475 pregnancies of women who
43
44 participated more than once. We excluded 703 women who did not answer the questions on
45
46 sexual violence. Further, we excluded records with missing data on gestational length
47
48 (n= 297) and birth weight (n=41). We also excluded 6 children with birth weight <500 g and 4
49
50 children with birth weight >6000 g, leaving a study sample of 76 870 women (Figure 1).
51
52
53
54
55

56 **Ethical statement**

57
58
59
60

1
2
3 Informed consent was obtained from each women participating in the study. The Regional
4
5 Committee for Medical Research Ethics (Ref.SAFH 95/313 RTL) and the Norwegian Data
6
7 Inspectorate approved the study.
8
9

10 11 12 13 14 **Variables**

15 16 Exposure variable

17
18 The exposure variable was collected from Q1 ~~;~~ ~~here, the w~~Women were asked if they had
19
20 been ~~1)~~ pressured or forced to ~~perform~~ sexual relations. There were four possible answer
21
22 options: 1) No, never acts, 2) Yes, pressured, 23) Yes, forced with violence or 43) Yes, raped.

23
24 A positive answer was defined as having experienced sexual violence. Women with more
25
26 than one positive answer were classified according to the most severe level reported. The
27
28 answering options were coded into three levels of severity for the sexual violence: 1) mild
29
30 (pressured), 2) moderate (forced with violence) and 3) severe (raped). ~~and by the timing of~~
31
32 the abuse (previously or recent). Recent meant exposed to sexual violence in the current
33
34 pregnancy or the last 6 or 12 months before pregnancy. Women could also indicate when the
35
36 violence had taken place: 1) during this pregnancy, 2) during the last six month before
37
38 pregnancy or 3) earlier. Approximately 1 700 women who filled out the first version of Q1
39
40 had the answering options earlier and during the last 12 month when assessing time. We
41
42 therefore created the variables previous and recent sexual violence, recent containing sexual
43
44 violence during last 12 month, including the current pregnancy. Among the women who
45
46 participated several times we included the first pregnancy only to ensure that the exposure
47
48 was included only once per woman. More details about the exposure variable can be found in
49
50 our previous studies ~~;~~ [23;24] and in supplementary table 1.
51
52
53
54
55
56
57
58
59
60

Outcome variables

All outcome variables were obtained from the MBRN. Gestational age at birth in days was based on ultrasound at (approximately) gestational week 18. For women with no ultrasound, the gestational age was based on the last menstrual period (1.7%). PTB was defined as a gestational age <37, LBW as a birth weight <2500 grams, and SGA was defined as birth weight below the 10th percentile for the gestational age at birth. SGA was calculated using Norwegian specific fetal growth tables by Skjerven et al.[25]

Adjusting variables

Maternal age, parity, socio-economical status, smoking and body mass index (BMI) were considered as possible confounding factors and were adjusted for. All adjusting variables were taken from the MoBa. In Q1, age was categorised into 5 groups: younger than 20 years, 20–24 years, 25–29 years, 30–34 years or 35 years and older. As a proxy for socio-economic status, we used the woman's education in years (categorised into 4 groups): primary (<12 years), secondary (12 years), higher ≤ 4 years (13–16 years and) and higher >4 years (≥ 17 years). Parity was dichotomised into nulli- and multiparous women. Smoking was categorised as no smoking or smoking, which included both daily and occasional smoking. BMI was grouped into 4 categories: <20, 20-24.9, 25.0-29.9 or ≥ 30.0 kg/m². We also adjusted for mental distress because it is considered to be associated with both the exposure and the outcome.[4,5] Mental distress was measured using 5 items from The Hopkins symptoms checklist (SCL-5) with a cut-off at ≥ 2.0 points, as suggested by Strand [26] and obtained from Q3.

1
2
3 Because of the co-occurrence of different violence types,[7] we examined the effect of
4 physical and emotional abuse as a child or as an adult in the multivariable statistical models.
5
6 Information on adult physical violence was taken from Q1 and consisted of a positive answer
7 to whether women as an adult had experienced being slapped, hit, kicked or otherwise
8 bothered in a physical manner. Child physical violence was taken from Q3 and consisted of a
9 positive answer to the question “Have you experienced physical violence before the age of
10 18?” Emotional abuse as a child (<18) or as an adult (≥18) consisted of a positive answer to
11 either, “Someone has over a long period of time systematically tried to subdue, degrade or
12 humiliate you” or “Someone has threatened to hurt you or someone close to you” or both.
13
14
15
16
17
18
19
20
21
22
23
24

25 Previous preterm birth and inadequate antenatal care are considered to be associated with the
26 exposure and the outcomes.[2, 3, 5] Because a previous preterm birth may be a result of
27 sexual violence prior to the related pregnancy, we did not control for a previous preterm birth.
28
29 In Norway, the majority of women attend antenatal care, a free and a well-integrated part of
30 the public health system; therefore, we did not control for antenatal care. Ethnicity was not
31 considered a relevant covariate confounding factor in our study because the majority of the
32 MoBa participants are ethnic Norwegian.
33
34
35
36
37
38
39
40
41
42

43 **Statistical Analysis**

44 Characteristics were presented as percentages within the entire sample and the different
45 outcomes. Linear regression was performed to assess differences in birth weight and
46 gestational age for children born to women with and without a history of mild, moderate and
47 severe sexual violence. The association between sexual violence and PTB, LBW and SGA
48 was estimated with crude and adjusted odds ratios using logistic regression analysis. Sexual
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 violence was analysed as a categorical variable: 1=mild sexual violence, 2= moderate sexual
4 violence and 3=severe sexual violence with no sexual violence as the reference group. All
5
6
7 analyses were adjusted for maternal age, parity, education, smoking, BMI and mental distress
8
9 in the first step. Birth weight was additionally adjusted for gestational age. We further
10
11 adjusted for other types of violence in the second step. We initially tested the correlation
12
13 between other types of violence and sexual violence because of co-occurrence, and all
14
15 Pearson's correlation coefficients were below the generally accepted cut-off of <0.4 for use as
16
17 a covariate in the regression analyses.[27] Post protocol, we stratified the sample into
18
19 spontaneous start of birth and provider-initiated start of birth (induced start of birth or elective
20
21 caesarean section) for gestational age because a provider-initiated start could influence the
22
23 time point of birth. Information on how the birth started was taken from MBRN. We
24
25 additionally performed a sensitivity analysis in which we examined the association between
26
27 sexual violence and SGA and LBW among women who had a spontaneous birth at term (≥ 37
28
29 weeks) because we wanted to examine the effect of violence in a group of women who were
30
31 considered to be low risk according to gestational age and start of birth. When we examined
32
33 the timing of the sexual violence, we compared women who were exposed to recent (~~sexual~~
34
35 violence (within the last ~~violence during pregnancy or the last 6 or~~ 12 months) and those
36
37 exposed to previous sexual violence to non-abused women. We also examined the timing
38
39 among women reporting recent and previous severe sexual violence (rape) for all outcomes.
40
41 The prevalence of missing data was generally low with 2.5% for BMI, 3.7% for education,
42
43 and 0.7% for smoking during pregnancy. Because of this, no imputing methods for missing
44
45 data were used,[28] except for the missing data for the SCL-5 (3.2%), which were replaced by
46
47 the series mean. The results of the logistic regression analyses remained approximately the
48
49 same when performed with the complete exclusion of missing data compared with using the
50
51 imputed missing data for SCL-5.
52
53
54
55
56
57
58
59
60

The comparison group for all analyses was women not reporting any sexual violence. All analyses were performed with the statistical package SPSS for WINDOWS (SPSS Inc., Chicago, IL, USA) version 18. P-values <0.05 were considered statistically significant.

RESULTS

The mean gestational age at birth in the total sample was 279.6 days (standard deviation 11.9 days), and the mean birth weight was 3592.7 grams (standard deviation 547.1 g). Table 1 displays the characteristics in the total sample and by the different outcomes.

Table 1. Background characteristics in the total sample and by preterm birth (PTB): gestational age <week 37, low birth weight (LBW): weight <2500 g and small for gestational age (SGA) weight below the 10th percentile by gestational age at birth in the Mother and Child Cohort

	Total	PTB	LBW	SGA
	N=76 870	N=3620	N=2107	N=6257
	%	%	%	%
Age				
<20	1.5	2.2	2.6	1.9
20–24	12.2	12.7	13.8	13.2
25–29	36.2	34.6	34.8	36.4
30–34	35.7	33.7	33.2	34.2
≥35	14.4	16.7	15.6	14.3
Education				
Primary	2.4	3.1	3.9	2.7
Secondary	34.8	38.5	38.8	35.0
Higher ≤4 years	37.8	35.6	37.3	36.1
Higher >4 years	21.3	19.2	20.5	22.2
Missing	3.7	3.6	3.7	3.9
Parity				
Nulliparous	54.9	61.7	68.2	70.7
Multiparous	45.1	38.3	31.8	29.3
Smoking				
No	90.9	89.3	85.8	85.6
Yes	8.5	10.1	13.6	13.6
Missing	0.7	0.6	0.6	0.7

1					
2					
3	Pre-pregnancy body mass				
4	index				
5	<20	12.4	12.8	15.6	19.1
6					
7	20–24.9	54.9	49.9	50.8	55.6
8					
9	25–29.9	21.0	22.0	19.0	15.2
10					
11	≥30	9.2	12.6	11.8	7.6
12					
13	Missing	2.5	2.7	2.9	2.4
14	Mental distress				
15	No	93.3	92.3	91.3	92.3
16					
17	Yes	6.8	7.7	8.7	7.7
18	Adult physical violence				
19	No	85.5	83.9	83.6	84.7
20					
21	Yes	14.5	16.1	16.4	15.3
22	Child physical violence				
23	No	94.5	94.4	94.1	94.3
24					
25	Yes	5.5	5.6	5.9	5.5
26	Adult emotional abuse				
27	No	83.6	82.1	81.3	82.5
28					
29	Yes	16.4	17.9	18.7	17.5
30	Child emotional abuse				
31	No	86.2	85.1	84.7	85.5
32					
33	Yes	13.8	14.9	15.3	14.5

34

35

36 The prevalence of adverse neonatal outcomes was generally highest in the youngest (<20 y)

37 and the oldest age groups (≥35 y) among smokers and women with primary school education.

38

39 A BMI ≥30 was associated with PTB and LBW and BMI <20 with SGA. Women who

40 reported mental distress also reported more PTB, LBW and SGA.

41

42

43

44

45

46

47 Among the 76 870 women enrolled, 9263 (12.1%) reported a history of mild sexual violence,

48 2102 (2.8%) moderate and 2746 (3.5%) severe. Women with a history of sexual violence

49 were significantly younger and they were more likely to have primary school education.

50

51 Additionally, these women more frequently reported smoking, a BMI ≥30 and mental distress.

52

53

54

55

56 These women more often experienced other types of violence both as children and adults

57 (data not provided in tables).

58

59

60

A lower gestational age at birth was observed for newborns from women who reported moderate and severe sexual violence with approximately two days when birth was provider-initiated (Table 2).

Table 2. Differences in gestational age for spontaneous and provider-initiated start of birth and birth weight for non-exposed women and women exposed to mild, moderate and severe sexual violence

	N= 76 870 (%)	Mean	Crude estimate β (95% CI)	Adjusted estimate β (95% CI)*
Gestational age				
No sexual violence	62 699 (81.6)	279.7 days	1	1
<i>Mild sexual violence</i>	9263 (12.1)			
Spontaneous start	7323 (11.5)	279.9 days	0.09 (-0.16; 0.33)	0.22 (-0.04; -0.48)
Provider-initiated start	1940 (12.5)	280.3 days	0.36 (-0.37; 1.08)	0.64 (-0.11; 1.39)
<i>Moderate sexual violence</i>	2162 (2.8)			
Spontaneous start	1670 (2.7)	279.3 days	-0.40 (-0.87; 0.06)	-0.28 (-0.78; 0.21)
Provider-initiated start	492 (3.2)	277.6 days	-2.13 (-3.41; -0.84)	-2.02 (-3.39; -0.67)
<i>Severe sexual violence</i>	2746 (3.6)			
Spontaneous start	2048 (3.3)	278.7 days	-0.91 (-1.37; -0.44)	-0.65 (-1.15; -0.16)
Provider-initiated start	698 (4.5)	277.5 days	-2.24 (-3.47; -1.01)	-1.92 (-3.22; -0.62)
Birth weight				
No sexual violence	62 699 (81.6)	3594 g	1	1
Mild sexual violence	9263 (12.1)	3597 g	2.03 (-9.26; 13.32)	0.04 (-9.16; 9.23)**
Moderate sexual violence	2162 (2.8)	3582 g	-13.61 (-34.74; 7.51)	6.11 (-11.30; 23.52)**
Severe sexual violence	2746 (3.6)	3556 g	-38.33 (-59.17; -17.49)	-0.76 (-18.05; 16.53)**

*Adjusted for maternal age, parity, education, smoking, body mass index and mental distress

**Additional adjustment for gestational age

Among women with a spontaneous start of birth, the gestational age was approximately one half of a day shorter when women reported severe sexual violence. These findings were significant in an adjusted analysis. A crude analysis showed that women who reported a history of severe violence delivered on average 38.3 g lighter children, a difference that disappeared when controlling for gestational age, mother's age, parity, education, smoking, BMI and mental distress. There were no differences regarding birth weight between women with a history of mild or moderate sexual violence compared with non-abused women.

Results from the logistic regression analysis are presented in Table 3.

Table 3. Odds of preterm birth (PTB), low birth weight (LBW) and small for gestational age (SGA) with 95% confidence intervals (CI) according to the different levels of sexual violence

	N (%)	Prevalence (%)	Crude odds ratio (95% CI)	Adjusted odds ratio (95% CI)*
PTB		3620 (4.7)		
No sexual violence	62 699 (81.6)	2931 (4.7)	1	1
Mild sexual violence	9263 (12.1)	412 (4.4)	0.95 (0.85-1.06)	0.93 (0.83-1.03)
Moderate sexual violence	2162 (2.8)	115 (5.3)	1.15 (0.95-1.39)	1.14 (0.93-1.39)
Severe sexual violence	2746 (3.6)	162 (5.9)	1.28 (1.08-1.51)	1.15 (0.97-1.37)
LBW		2107 (2.7)		
No sexual violence	62 699 (81.6)	1681 (2.7)	1	1
Mild sexual violence	9263 (12.1)	257 (2.8)	1.04 (0.91-1.18)	0.98 (0.85-1.12)
Moderate sexual violence	2162 (2.8)	75 (3.5)	1.30 (1.03-1.65)	1.19 (0.93-1.53)
Severe sexual violence	2746 (3.6)	94 (3.4)	1.29 (1.04-1.59)	1.07 (0.85-1.34)
SGA		6257 (8.1)		
No sexual violence	62 699 (81.6)	5061 (8.1)	1	1
Mild sexual violence	9263 (12.1)	768 (8.3)	1.03 (0.95-1.12)	1.00 (0.91-1.08)
Moderate sexual violence	2162 (2.8)	178 (8.3)	1.02 (0.87-1.19)	0.95 (0.80-1.12)
Severe sexual violence	2746 (3.6)	250 (9.1)	1.14 (1.00-1.30)	1.05 (0.91-1.21)

*Adjusted for maternal age, parity, education, smoking, body mass index and mental distress

Women who reported severe sexual violence had higher odds of PTB, LBW and SGA in a crude analysis, an association that was attenuated and no longer significant when adjusted for maternal age, parity, education, smoking, BMI and mental distress. Other types of violence, both as a child and adult, had small attenuating effects on the odds ratios and were not included in the final models.

The sensitivity analysis, in which we examined the association between a history of sexual violence and SGA and LBW in a sub-sample of women who had a spontaneous term birth, showed the same pattern as in the total sample reported in Table 3. Women who reported severe sexual violence had higher odds of LBW and SGA in a crude analysis but not in the adjusted analysis (Data not provided in tables).

1
2
3 A crude analysis was used to examine if the timing of the violence was associated with
4
5 adverse outcome. Women who reported recent sexual violence had a higher risk for LBW
6
7 (OR 1.60 95% CI 1.04-2.17) compared with non-abused women. The association was no
8
9 longer significant in the adjusted analysis. In our study, 684 (0.9%) women reported recent
10
11 sexual violence (mild, moderate and severe) and 13 487 (17.5%) previous sexual violence
12
13 (Supplementary Table S24). There was no association between recent severe sexual violence
14
15 (rape) and adverse neonatal outcome (Supplementary Table S32). There were 66 (0.1%)
16
17 women who reported ~~recent rape during pregnancy or the last 6 or in 12 months in~~ this study.
18
19
20
21
22
23

24 DISCUSSION

25 Main outcome

26
27 We found that moderate and severe sexual violence were associated with a reduction in
28
29 gestational age at birth. The largest effect was observed when birth was provider-initiated
30
31 among women exposed to moderate or severe violence. These women had an approximately
32
33 two-day reduction in gestational age. There was no significant association between sexual
34
35 violence and PTB, LBW or SGA in the adjusted analysis.
36
37
38
39
40
41

42 Strength

43
44 This study, based on information from a large population-based study, the Norwegian Mother
45
46 and Child Cohort study (MoBa), which is linked to the Medical Birth Registry (MBRN), gave
47
48 a unique opportunity to assess the association between sexual violence and outcome for
49
50 newborns. The validity of the data in ~~the~~ MoBa has in earlier research been described as
51
52 sufficient for large-scale epidemiologic studies.[29, 30] Our study was strengthened by the
53
54 fact that the information on the different outcome variables was collected prospectively from
55
56 the quality-assessed MBRN.[31] The outcomes in this study are part of a complex
57
58
59
60

phenomenon that has several different risk factors.[2] The setting in this study, with small social and health inequalities, may therefore be suitable to isolate the effect of sexual violence on adverse neonatal outcomes.

Limitations

There are also limitations to our study. The participation rate of 40.6% in MoBa is low, and MoBa suffers to some extent from selection bias. The women included in the study are older, have more education, smoke less and are less likely to be of a non-Norwegian origin than the Norwegian population. Although it is likely that there is a socioeconomic gradient that influence prevalence estimates, a recent study by Nilsen et al. has found that this may affect the prevalence estimates, but there was no evidence that the exposure-outcome associations in the MoBa study were affected by selection bias.[32] This socioeconomic gradient may also Furthermore, as preterm birth is associated with ethnicity, the ethnic homogeneous sample in MoBa may limit the generalisability of our findings. The lack of a validated instrument for measuring the exposure is a limitation to this study, and violence measured in modules as part of a larger questionnaire, as that in MoBa, may achieve a lower disclosure rate.[5] However, a similar prevalence to that found in our study was reported in a Nordic study examining sexual violence and health.[33] The exposure was measured in gestational week 17, and sexual violence during pregnancy after this is therefore not included. In addition, we have no information on the context and frequency of the violence or information regarding the perpetrator.

To our knowledge, no studies have examined the influence of lifetime sexual violence reported during pregnancy on the gestational age at birth for newborns. There were minor differences in the gestational age between abused and non-abused women in this study, and

1
2
3 the clinical importance of our findings for the health of the newborn is most likely limited.
4
5 However, the difference between the provider-initiated and spontaneous initiation of birth
6
7 may be of interest. Shorter provider-initiated pregnancies may suggest an increase in elective
8
9 inductions and elective caesarean sections for those exposed to violence. This is supported by
10
11 others [34, 35] and in our previous study on sexual violence and maternal outcome.[36]
12
13 Studies have emphasised the meaning of control for abused women when giving birth,[37]
14
15 and choosing a planned start of birth may help the abused women remain in control.
16
17
18
19

20
21 Unlike the current meta-analysis (~~not yet published~~), mentioned in a WHO report (~~not yet~~
22
23 ~~published~~);[5] we did not find sexual violence to be associated with PTB and LBW in
24
25 adjusted analysis. To our knowledge, no study has found an association between violence and
26
27 SGA. The studies included in the meta-analysis ~~were limited to investigated both~~ sexual
28
29 and/or physical intimate partner violence, ~~in contrast to our study, which investigated lifetime~~
30
31 ~~sexual violence~~. However, our findings are supported by a Canadian population based study
32
33 with a sample of 6 421 pregnant women [17] and a prospective cohort study including 1 555
34
35 women from the US [15]. The exposure in these studies was physical and sexual violence
36
37 prior to pregnancy and in pregnancy without being limited to an intimate partner. [15, 17]
38
39 These studies showed no association between violence and PTB or LBW.
40
41
42
43
44

45 The nature of the exposure measured in the MoBa study makes it difficult to directly compare
46
47 our findings to others, mainly because we examined lifetime sexual violence by any
48
49 perpetrator, not limited to intimate partner. Sexual violence is considered to be traumatic for
50
51 the victim regardless if the perpetrator is a partner or not.[38] When an intimate partner is the
52
53 perpetrator, sexual violence may in addition be accompanied by controlling behaviour and
54
55 include both physical and emotional abuse.[38] Rape by strangers is usually a single violent
56
57
58
59
60

1
2
3 event with a higher risk of physical injury. Both forms of violence are associated with adverse
4 health effects, [5, 38] but the effect may differ. -Studies on the health effects of non-partner
5 sexual violence are limited with less robust data than for intimate partner violence.[5]
6
7 Unfortunately, we were not able to examine the effect of the perpetrator in this study because
8 the MoBa study does not provide this information. However, research suggests that a
9 substantial proportion of sexual violence occurs within an intimate relationship.[38] Our data
10 from the MoBa does not differentiate between intimate partner violence and sexual abuse by
11 non-partnersIt, -but it is not unlikely that the question about severe sexual violence (rape)
12 primarily reflects non-partner sexual abuse. -and that mild sexual violence (pressured to
13 sexual acts) may be a more psychological exposure. Our crude analyses showed different
14 results, with a significantly higher OR for adverse neonatal outcome among women who
15 reported severe sexual violence, thus supporting the idea that the different levels of violence
16 are different exposures. The use of a more comprehensive instrument when measuring the
17 exposure, with multiple response options regarding context, frequency and perpetrator would
18 have clarified this further and provided more comprehensive knowledge about the nature of
19 the violence. Nevertheless, we had the opportunity to control for both emotional and physical
20 abuse in preliminary analysis and this did not change the ORs.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 =

44
45
46
47 Several pathways between sexual violence and adverse neonatal outcomes are suggested, both
48 direct and indirect.[5] The direct pathway of violence during pregnancy can lead to a preterm
49 birth, and examples of indirect pathways include more health-risk behaviours, depression and
50 stress/anxiety.[5, 40] Both experienced violence and living in an abusive environment can
51 cause increased stress levels, which could be on the pathway between abuse and adverse
52
53
54
55
56
57
58
59
60

1
2
3 neonatal outcome. Maternal exposure to stress can influence the hypothalamic-pituitary axis
4 (HPA) hormones, and it is suggested that changes in these hormones may cause negative
5 outcomes, such as a reduction in gestational age and foetal growth restriction.[4, 41]
6
7

8
9 The prevalence of abuse during pregnancy is small in our study and may have decreased the
10 power to detect an association between violence during pregnancy and adverse neonatal
11

12 outcomes, an association that is supported by other studies.[10, 14, 4239, 4340] Living in an
13

14 abusive relationship may have stopped women from disclosing the violence. In our study,
15

16 sexual violence was assessed approximately in gestational week 17, and events of violence
17

18 after that ~~may~~ have been missed. Some studies suggest that the risk of sexual violence may
19

20 increase with the length of the pregnancy for women that are exposed.[41]- The studies that
21

22 report the highest prevalence of violence have measured this several times during pregnancy.
23

24 [6] Studies have reported a violence prevalence during pregnancy of between 3 to 19%.[6,
25

26 442], including both ~~A direct comparison to our study is difficult because these numbers~~
27

28 estimate both physical and sexual violence. ~~abuse during pregnancy; they may suggest that~~
29

30 sexual violence during pregnancy is underreported in our study. Underreporting among the
31

32 non-exposed may have caused a misclassification that has diminished the associations
33

34 between sexual violence and neonatal outcomes in our study. Since the exposure was
35

36 collected before the outcome, it is unlikely that misclassification was related to the outcome,
37

38 thus resulting in a nondifferential misclassification that has biased the result towards the null.
39

40 Living in an abusive relationship may also stop women from disclosing the violence.
41

42 However, our prevalence of recent sexual (0.9%) violence is similar to the prevalence of 1%
43

44 reported in a survey that assessed intimate partner violence in a national population based
45

46 study on violence among a representative sample of Norwegian women.[453] This number
47

48 reflects the prevalence of sexual violence reported during the last year. r, not especially
49

50 among pregnant women.
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5 Several pathways between sexual violence and adverse neonatal outcomes are suggested, both
6 direct and indirect.[5] The direct pathway of violence during pregnancy can cause immediate
7 complications as bleeding, rupture of membranes and PTB, and examples of indirect
8 pathways include more health-risk behaviours, depression and stress/anxiety.[5, 44] Both
9 experienced violence and living in an abusive environment can cause increased stress levels,
10 which could be on the pathway between abuse and adverse neonatal outcome. Maternal
11 exposure to stress can influence the hypothalamic pituitary axis (HPA) hormones, and it is
12 suggested that changes in these hormones may cause negative outcomes, such as a reduction
13 in gestational age and fetal growth restriction.[4, 45] It has been proposed that mental distress
14 and symptoms of depression are on the causal pathway between violence and adverse health
15 outcome, yet it has also been suggested that women with mental health difficulties are more
16 likely to be victims of violence.[46] Because the relationship may be bi-directional, we chose
17 to control for mental distress in our study. Similarly, the health risk behaviours smoking and
18 BMI may be on the pathway between sexual violence and neonatal outcome. Nevertheless, we
19 kept these covariates in the regression analysis because they are especially related to birth
20 weight and PTB.[3, 47]
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 **Conclusion**

46
47 Overall, our findings provide no evidence for an association between lifetime sexual violence
48 and adverse neonatal outcomes. A small significant effect on the gestational age at birth was
49 detected, but the clinical importance of this is most likely limited for the health of the
50 newborn. PTB, LBW and SGA all have complex origins with multiple possible pathways.[1,
51
52
53
54
55
56
57
58
59
60
2] Although we did not find an association between sexual violence and PTB, LBW or SGA

1
2
3 in adjusted analyses, crude analyses in our study suggested that sexual violence may be a risk
4
5 factor for adverse neonatal outcomes for some women; however, for the majority of women,
6
7 the relationship was confounded by other risk factors. It is possible that these factors were
8
9 the result of prior exposure to violence, but this could not be assessed in this study. Antenatal
10
11 care is one of the most important entry points in which women seek health care without
12
13 necessarily disclosing ongoing exposure to violence or a history of sexual violence. It is
14
15 recommended that caregivers and clinicians ask women about exposure to violence when
16
17 assessing conditions that may be complicated by violence,[486] and more knowledge about
18
19 this conditions is needed. Additionally, antenatal care may offer opportunities for women to
20
21 receive help, both if they are exposed to violence and in providing assistance to change
22
23 behavioural factors contributing to adverse outcomes.
24
25
26
27
28

29 **Competing interests**

30
31 None.
32
33
34

35 **Contributions to authorship**

36
37 LH participated in the conception and design of the study, performed the analysis and drafted
38
39 the manuscript. BS participated in the conception of the study, advised on the statistical
40
41 analyses and drafting of the manuscript. SV advised on the statistical analyses and the drafting
42
43 of the manuscript. ML participated in the conception and design of the study, advised on the
44
45 statistical analyses and participated in drafting the manuscript. All authors read and approved
46
47 the final version.
48
49
50
51
52

53 **Funding**

54
55 The Norwegian Mother and Child Cohort Study is supported by the Norwegian Ministry of
56
57 Health and the Ministry of Education and Research, NIH/NIEHS (contract no N01-ES-
58
59
60

1
2
3 75558), NIH/NINDS (grant no.1 UO1 NS 047537-01 and grant no.2 UO1 NS 047537-06A1),
4
5 and the Norwegian Research Council/FUGE (grant no. 151918/S10). The first author of this
6
7 article is supported by [South-Eastern Norway Regional Health Authority \(grant no.](#)
8
9 [2012010\)The Research Council of Norway \(grant no. 204292/H10\)](#), and the second author is
10
11 supported by the [The Research Council of Norway \(grant no. 204292/H10\)](#), [South Eastern](#)
12
13 [Norway Regional Health Authority \(grant no. 2012010\)](#).
14
15
16
17
18

19 Acknowledgement

20 We are grateful to all the participating families in Norway who take part in this ongoing
21
22 cohort study.
23
24
25
26
27

28 Reference List

- 29 (1) Green NS, Damus K, Simpson JL, et al. Research agenda for preterm birth:
30 recommendations from the March of Dimes. *Am J Obstet Gynecol* 2005;193:626-35.
- 31 (2) March of Dimes, PMNCH, Save The Children, WHO. Born Too Soon: The Global
32 Action Report on Preterm Birth. Geneva: World Health Organization 2012
- 33 (3) Wilcox AJ. On the importance-and the unimportance-of birth weight. *Int J Epidemiol*
34 2001;30:1233-41.
- 35 (4) Lazinski MJ, Shea AK, Steiner M. Effects of maternal prenatal stress on offspring
36 development: a commentary. *Arch Womens Ment Health* 2008;11:363-75.
- 37 (5) Garcia-Moreno C, Pallitto C, Devries K, et al. Global and regional estimates of
38 violence against women: prevalence and health effects of intimate partner violence
39 and non-partner sexual violence. Geneva: World Health Organization; 2013.
- 40 (6) Campbell J, Garcia-Moreno C, Sharps P. Abuse During Pregnancy in Industrialized
41 and Developing Countries. *Violence Against Women* 2004;10:770-89.
- 42 (7) Garcia-Moreno C, Jansen HA, Ellsberg M, et al. Prevalence of intimate partner
43 violence: findings from the WHO multi-country study on women's health and
44 domestic violence. *Lancet* 2006;368:1260-9.
- 45 (8) Sanchez SE, Alva AV, Diez CG, et al. Risk of spontaneous preterm birth in relation to
46 maternal exposure to intimate partner violence during pregnancy in Peru. *Matern*
47 *Child Health J* 2013;17:485-92.
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 (9) Leeners B, Stiller R, Block E, et al. Pregnancy complications in women with
4 childhood sexual abuse experiences. *J Psychosom Res* 2010;69:503-10.
5
6 (10) Nunes MA, Camey S, Ferri CP, et al. Violence during pregnancy and newborn
7 outcomes: a cohort study in a disadvantaged population in Brazil. *Eur J Public Health*
8 2011;21(1):92-7.
9
10 (11) Rodrigues T, Rocha L, Barros H. Physical abuse during pregnancy and preterm
11 delivery. *Am J Obstet Gynecol* 2008;198:171-6.
12
13 (12) Noll JG, Schulkin J, Trickett PK, et al. Differential pathways to preterm delivery for
14 sexually abused and comparison women. *J Pediatr Psychol* 2007;32:1238-48.
15
16 (13) Silverman JG, Decker MR, Reed E, et al. A. Intimate partner violence victimization
17 prior to and during pregnancy among women residing in 26 U.S. states: associations
18 with maternal and neonatal health. *Am J Obstet Gynecol* 2006;195:140-8.
19
20 (14) Murphy CC, Schei B, Myhr TL, et al. Abuse: a risk factor for low birth weight? A
21 systematic review and meta-analysis. *CMAJ* 2001;164:1567-72.
22
23 (15) Fried LE, Cabral H, Amaro H, et al. Lifetime and during pregnancy experience of
24 violence and the risk of low birth weight and preterm birth. *J Midwifery Womens*
25 *Health* 2008 ;53:522-8.
26
27 (16) Janssen PA, Holt VL, Sugg NK, et al. Intimate partner violence and adverse
28 pregnancy outcomes: a population-based study. *Am J Obstet Gynecol* 2003;188:1341-
29 7.
30
31 (17) Urquia ML, O'Campo PJ, Heaman MI, et al. Experiences of violence before and
32 during pregnancy and adverse pregnancy outcomes: an analysis of the Canadian
33 Maternity Experiences Survey. *BMC Pregnancy Childbirth* 2011;11:42.
34
35 (18) Altarac M, Strobino D. Abuse during pregnancy and stress because of abuse during
36 pregnancy and birthweight. *J Am Med Womens Assoc* 2002;57:208-14.
37
38 (19) Cokkinides VE, Coker AL, Sanderson M, et al. Physical violence during pregnancy:
39 maternal complications and birth outcomes. *Obstet Gynecol* 1999;93:661-6.
40
41 (20) Grimstad H, Schei B, Backe B, et al. Physical abuse and low birthweight: a case-
42 control study. *Br J Obstet Gynaecol* 1997;104:1281-7.
43
44 (21) Kaye DK, Mirembe FM, Bantebya G, et al. Domestic violence during pregnancy and
45 risk of low birthweight and maternal complications: a prospective cohort study at
46 Mulago Hospital, Uganda. *Trop Med Int Health* 2006;11:1576-84.
47
48 (22) Magnus P, Irgens LM, Haug K, et al. Cohort profile: the Norwegian Mother and Child
49 Cohort Study (MoBa). *Int J Epidemiol* 2006;35:1146-50.
50
51 (23) Lukasse M, Henriksen L, Vangen S, et al. Sexual violence and pregnancy-related
52 physical symptoms. *BMC Pregnancy Childbirth* 2012;12:83.
53
54
55
56
57
58
59
60

- 1
2
3 (24) Henriksen L, Vangen S, Schei B, et al. Sexual violence and antenatal hospitalization. *Birth* 2013;40:281-8.
- 4
5
6 (25) Skjaerven R, Gjessing HK, Bakketeig LS. et al. Birthweight by gestational age in
7 Norway. *Acta Obstet Gynecol Scand* 2000;79:440-9.
- 8
9 (26) Strand BH, Dalgard OS, Tambs K, et al. Measuring the mental health status of the
10 Norwegian population: a comparison of the instruments SCL-25, SCL-10, SCL-5 and
11 MHI-5 (SF-36). *Nord J Psychiatry* 2003;57:113-8.
- 12
13 (27) Field A. Discovering statistics using SPSS. Los Angeles: SAGE; 2009.
- 14
15 (28) Veierød MB, Lydersen S, Laake P, et at. Medical statistics: in clinical and
16 epidemiological research. Oslo: Gyldendal akademisk; 2012.
- 17
18 (29) Kvalvik LG, Nilsen RM, Skjaerven R, et al. Self-reported smoking status and plasma
19 cotinine concentrations among pregnant women in the Norwegian Mother and Child
20 Cohort Study. *Pediatr Res* 2012 ;72:101-7.
- 21
22 (30) Furu K, Karlstad O, Skurtveit S, et al. High validity of mother-reported use of
23 antiasthmatics among children: a comparison with a population-based prescription
24 database. *J Clin Epidemiol* 2011;64:878-84.
- 25
26 (31) Irgens LM. The Medical Birth Registry of Norway. Epidemiological research and
27 surveillance throughout 30 years. *Acta Obstet Gynecol Scand* 2000;7:435-9.
- 28
29 (32) Nilsen RM, Vollset SE, Gjessing HK, et al. Self-selection and bias in a large
30 prospective pregnancy cohort in Norway. *Paediatr Perinat Epidemiol* 2009 23:597-
31 608.
- 32
33 (33) Hilden M, Schei B, Swahnberg K, et al. A history of sexual abuse and health: a Nordic
34 multicentre study. *BJOG* 2004;111:1121-7.
- 35
36 (34) Nerum H, Halvorsen L, Straume B, et al. Different labour outcomes in primiparous
37 women that have been subjected to childhood sexual abuse or rape in adulthood: a
38 case-control study in a clinical cohort. *BJOG* 2013;120:487-495.
- 39
40 (35) Lukasse M, Vangen S, Oian P, et al. Childhood abuse and caesarean section among
41 primiparous women in the Norwegian Mother and Child Cohort Study. *BJOG*
42 2010;117:1153-7.
- 43
44 (36) Henriksen L, Schei B, Vangen S, et al. Sexual violence and mode of delivery: a
45 population-based cohort study. *BJOG*. ~~In press 2014.~~ [2014 121\(10\):1237-1244](#)
- 46
47 (37) Montgomery E. Feeling Safe: A Metasynthesis of the Maternity Care Needs of
48 Women Who Were Sexually Abused in Childhood. *Birth* 2013;40:88-95.
- 49
50
51
52
53 (38) [Abrahams N, Devries K, Watts C, Pallitto C, Petzold M, Shamu S, et al. Worldwide](#)
54 [prevalence of non-partner sexual violence: a systematic review. *Lancet*.](#)
55 [2014;383\(9929\):1648-54. Epub 2014/02/18.](#)
56
57
58
59
60

Campbell J, Torres S, Ryan J, et al. Physical and nonphysical partner abuse and other risk factors for low birth weight among full term and preterm babies: a multiethnic case control study. *Am J Epidemiol* 1999;150:714-26.

(39) Webster J, Chandler J, Battistutta D. Pregnancy outcomes and health care use: effects of abuse. *Am J Obstet Gynecol* 1996;174:760-7

(4039) Faramarzi M, Esmaelzadeh S, Mosavi S. Prevalence, maternal complications and birth outcome of physical, sexual and emotional domestic violence during pregnancy. *Acta Med Iran* 2005;43:115-22. Leeners B, Neumaier-Wagner P, Quarg AF, et al. Childhood sexual abuse (CSA) experiences: an underestimated factor in perinatal care. *Acta Obstet Gynecol Scand* 2006;85:971-6.

(4140) Valladares E, Ellsberg M, Pena R, et al. Physical partner abuse during pregnancy: a risk factor for low birth weight in Nicaragua. *Obstet Gynecol* 2002;100:700-5.

(41) Shah PS, Shah J. Maternal exposure to domestic violence and pregnancy and birth outcomes: a systematic review and meta-analyses. *J Womens Health (Larchmt)*. 2010;19:2017-31. Valladares E, Pena R, Ellsberg M, et al. Neuroendocrine response to violence during pregnancy--impact on duration of pregnancy and fetal growth. *Acta Obstet Gynecol Scand* 2009;88:818-23.

(42) Sharps PW, Laughon K, Giangrande SK. Intimate partner violence and the childbearing year: maternal and infant health consequences. *Trauma Violence Abuse* 2007;8:105-16. Faramarzi M, Esmaelzadeh S, Mosavi S. Prevalence, maternal complications and birth outcome of physical, sexual and emotional domestic violence during pregnancy. *Acta Med Iran* 2005;43:115-22.

(43) Neroien AI, Schei B. Partner violence and health: results from the first national study on violence against women in Norway. *Scand J Public Health* 2008;36:161-8. Valladares E, Ellsberg M, Pena R, et al. Physical partner abuse during pregnancy: a risk factor for low birth weight in Nicaragua. *Obstet Gynecol* 2002;100:700-5.

(44) Leeners B, Neumaier-Wagner P, Quarg AF, et al. Childhood sexual abuse (CSA) experiences: an underestimated factor in perinatal care. *Acta Obstet Gynecol Scand* 2006;85:971-6.

Sharps PW, Laughon K, Giangrande SK. Intimate partner violence and the childbearing year: maternal and infant health consequences. *Trauma Violence Abuse* 2007;8:105-16. (45) Valladares E, Pena R, Ellsberg M, et al. Neuroendocrine response to violence during pregnancy--impact on duration of pregnancy and fetal growth. *Acta Obstet Gynecol Scand* 2009;88:818-23. Neroien AI, Schei B. Partner violence and health: results from the first national study on violence against women in Norway. *Scand J Public Health* 2008;36:161-8.

(46) Devries KM, Mak JY, Bacchus LJ, Child JC, Falder G, Petzold M, et al. Intimate partner violence and incident depressive symptoms and suicide attempts: a systematic review of longitudinal studies. *PLoS Med* 2013;10:e1001439. World health organisation. Responding to intimate partner violence and sexual violence against women: WHO clinical and policy guidelines. Geneva: WHO; 2013


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

(47) McDonald SD, Han Z, Mulla S, Beyene J. Overweight and obesity in mothers and risk of preterm birth and low birth weight infants: systematic review and meta-analyses. *BMJ* 2010;341:3428.

(48) World health organisation. Responding to intimate partner violence and sexual violence against women: *WHO clinical and policy guidelines*. Geneva: WHO; 2013.

For peer review only

Figure 1. Inclusion and exclusion process



90x107mm (300 x 300 DPI)

Table S1. Questions on sexual violence in The Norwegian Mother and Child Cohort Study (MoBa)

The first version, answered by 1712 women (version A):

Have you ever in been pressured or forced to sexual relations?

(Fill in one or several boxes.)

	Last 12 months	Earlier
No, never.....	<input type="checkbox"/>	<input type="checkbox"/>
Yes, pressured.....	<input type="checkbox"/>	<input type="checkbox"/>
Yes, forced with violence.....	<input type="checkbox"/>	<input type="checkbox"/>
Yes, raped.....	<input type="checkbox"/>	<input type="checkbox"/>

The questions used in the other versions:

Have you ever in been pressured or forced to sexual relations?

(Fill in one or several boxes.)

	During this pregnancy	Last 6 months before pregnancy	Earlier
No, never.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Yes, pressured.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Yes, forced with violence.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Yes, raped.....	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Table S2: Odds of preterm birth (PTB), low birth weight (LBW) and small for gestational age (SGA) with 95% confidence intervals (CI) according to previous and recent sexual violence

	N (%)	Prevalence (%)	Crude odds ratio (95% CI)	Adjusted odds ratio (95% CI)
PTB		3620 (4.7)		
No sexual violence	62 699 (81.6)	2931 (4.7)	1	1
Previous sexual violence	13487 (17.5)	650 (4.8)	1.03 (0.95-1.13)	0.99 (0.90-1.09)
Recent sexual violence	684 (0.9)	39 (5.7)	1.23 (0.89-1.71)	1.17 (0.83-1.65)
LBW		2107 (2.7)		
No sexual violence	62 699 (81.6)	1681 (2.7)	1	1
Previous sexual violence	13 487 (17.5)	397 (2.9)	1.10 (0.99-1.23)	1.0 (0.86-1.12)
Recent sexual violence	684 (0.9)	29 (4.2)	1.60 (1.04-2.33)	1.45 (0.97-2.17)
SGA		6257 (8.1)		
No sexual violence	62 699 (81.6)	5061 (8.1)	1	1
Previous sexual violence	13 487 (17.5)	1136 (8.4)	1.05 (0.98-1.12)	0.98 (0.92-1.06)
Recent sexual violence	684 (0.9)	60 (8.8)	1.10 (0.84-1.43)	1.18 (0.90-1.56)

*Adjusted for maternal age, parity, education, smoking, body mass index and mental distress

Table S3: Odds of preterm birth (PTB), low birth weight (LBW) and small for gestational age (SGA) with 95% confidence intervals (CI) according to previous and recent severe violence (rape)

	N (%)	Prevalence (%)	Crude odds ratio (95% CI)	Adjusted odds ratio (95% CI)*
PTB		3620 (4.7)		
No rape	74 124 (96.4)	3458 (4.7)	1	1
Previous rape	2680 (3.5)	157 (5.9)	1.27 (1.08-1.50)	1.14 (0.97-1.36)
Recent rape	66 (0.1)	5 (7.6)	1.68 (0.67-4.17)	1.72 (0.68-4.33)
LBW		2107 (2.7)		
No rape	74 124 (96.4)	2013(2.7)	1	1
Previous rape	2680 (3.5)	91 (3.4)	1.26 (1.02-1.56)	1.06 (0.84-1.33)
Recent rape	66 (0.1)	3 (4.5)	1.71 (0.54-5.44)	1.55 (0.48-5.02)
SGA		6257 (8.1)		
No rape	74 124 (96.4)	6007 (8.1)	1	1
Previous rape	2680 (3.5)	247 (9.2)	1.15 (1.01-1.32)	1.06 (0.92-1.23)
Recent rape	66 (0.1)	3 (4.5)	0.54 (0.17-1.72)	0.55 (0.17-1.79)

*Adjusted for maternal age, parity, education, smoking, body mass index and mental distress

STROBE 2007 (v4) Statement—Checklist of items that should be included in reports of *cohort studies*

Section/Topic	Item #	Recommendation	Reported on page #
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	page 1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	page 2
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	page 3 and
Objectives	3	State specific objectives, including any prespecified hypotheses	page 4
Methods			
Study design	4	Present key elements of study design early in the paper	From page 4
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection in addition reference is given for further reading of detailed information on MoBa study (referece 22)	Page 4
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up	NA
		(b) For matched studies, give matching criteria and number of exposed and unexposed	NA
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	Presented on pages 5-7
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	Page 5-7
Bias	9	Describe any efforts to address potential sources of bias controlling for confounding and covariates	pages 6 and 7.
Study size	10	Explain how the study size was arrived at	figure 1 and page5
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	Page 6
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	page 7 and 8
		(b) Describe any methods used to examine subgroups and interactions	page 8
		(c) Explain how missing data were addressed	page 8
		(d) If applicable, explain how loss to follow-up was addressed	NA
		(e) Describe any sensitivity analyses	page 8
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed All listed in Figure 1	
		(b) Give reasons for non-participation at each stage	Page 5

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

		(c) Consider use of a flow diagram Figure 1	
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	Table 1: page 9 and 10
		(b) Indicate number of participants with missing data for each variable of interest	Table 1
		(c) Summarise follow-up time (eg, average and total amount)	NA
Outcome data	15*	Report numbers of outcome events or summary measures over time	Table 1
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	Page 6-7, 11 - 12 and Table 2 and 3
		(b) Report <i>category</i> boundaries when continuous variables were categorized	Page 6 and Table 1
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	page 12-13 Supplementary Tables 1 and 2
Discussion			
Key results	18	Summarise key results with reference to study objectives	page 13
Limitations			
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	Page 13-16
Generalisability	21	Discuss the generalisability (external validity) of the study results	Page 14
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	Page 17

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.