

Experimental Pulmonary Inflammatory Injury in the Monkey

Susan D. Revak, Charles L. Rice, Ingrid U. Schraufstatter, Wayne A. Halsey, Jr., Benjamin P. Bohl, Robert M. Clancy, and Charles G. Cochrane

Department of Immunology, Scripps Clinic and Research Foundation, La Jolla, California 92037; and Department of Surgery, Harborview Medical Center, Seattle, Washington 98104

Abstract

Inflammatory pulmonary injury was induced in *Macacca mulatta* rhesus monkeys by the intrabronchial instillation of the formylated peptide norleu-leu-phe (FNLP) or phorbol myristate acetate (PMA). Indicators of pulmonary injury included an increase in mean protein content of bronchoalveolar lavage (BAL) fluid from 0.51 mg/ml in untreated animals to 3.74 mg/ml and 6.64 mg/ml in FNLP- and PMA-treated animals, respectively, the appearance of a diffuse pulmonary infiltrate in chest roentgenograms, and histologic evidence of a predominantly neutrophilic leukocytic infiltration.

Concomitant with the appearance of pulmonary injury was the generation of proteases and oxidants in the BAL fluids. Neutrophil elastase, bound to α_1 -protease inhibitor (α_1 -PI), was found to increase from 0.47 μ g/ml in untreated monkeys to 0.99 μ g/ml in FNLP-treated animals and 1.23 μ g/ml in monkeys receiving PMA. Radioiodinated human prekallikrein, instilled for 2 min into the inflammatory site and retrieved by lavaging, was found to have undergone proteolytic cleavage; this cleavage was not consistently inhibitable with the inclusion of antibody to elastase. BAL fluids were shown to contain an amidolytic activity when tested on the synthetic substrate *H*-D-pro-phe-arg-pNA. This activity was partially inhibitable with known inhibitors of active Hageman factor and kallikrein. β -Glucuronidase levels in the BAL fluids increased from 0.85 U/ml to 4.36 U/ml and 8.25 U/ml in FNLP- and PMA-treated animals, respectively. Myeloperoxidase (MPO) levels also increased from 1.37 OD U/ml \cdot min to 16.59 and 30.47 OD U/ml \cdot min in the same groups of animals.

Oxidant generation was also assessed in several different ways. The specific activity of the oxidant-sensitive inhibitor α_1 -PI recovered in the BAL fluid decreased from 0.80 in control samples to 0.57 and 0.65 in FNLP- and PMA-treated animals. That this inactivation was due to oxidant injury of the molecule was confirmed by the return to full activity of four out of five BAL samples after their incubation with the reducing agent dithiothreitol in the presence of methionine sulfoxide peptide reductase. The specific activity of catalase in the BAL fluids of animals given 3-amino, 1,2,4 triazole (AT) 1 h before lavaging showed drops from 0.97 in untreated monkeys to 0.04 in FNLP-treated and 0.49 in PMA-treated monkeys. MPO levels also fell in the AT-treated injured animals from 16.59 to 0.85 Δ OD/

min \cdot ml in FNLP animals in the absence and presence of AT, and 30.47 to 0.60 Δ OD/min \cdot ml in PMA-treated animals. Inhibition of MPO by AT was shown in vitro to be H₂O₂ dependent. Total glutathione levels in the BAL fluids did not change appreciably after FNLP or PMA treatment.

These studies present substantial evidence of the generation of both proteases and oxidants during the establishment of acute pulmonary inflammatory injury in an experimental primate model.

Introduction

Both proteases and oxidants may play important roles in the development of pulmonary inflammatory injury. Both have been detected in vivo in human disease and in models of pulmonary inflammation in experimental animals (1–13). Recent studies have revealed the presence of neutrophilic elastase in bronchoalveolar lavage (BAL)¹ fluids from patients with respiratory distress syndrome and other inflammatory conditions (3, 6, 10). Neutrophilic acid proteases and other leukocytic enzymes were found in BAL fluids from rabbits undergoing experimentally induced pulmonary inflammation (13).

Oxidant activity in situ has been measured indirectly. In human beings with pulmonary inflammation, evidence for the generation of oxidants in the bronchoalveolar region has been provided by analysis of α_1 -proteinase inhibitor (α_1 -PI). α_1 -PI is known to lose inhibitory activity for proteases when a methionyl residue in the active site is converted to methionyl sulfoxide (14–16). In vitro studies have shown that human α_1 -PI can become inactivated in the presence of halide and phorbol myristate acetate (PMA)-stimulated human neutrophils, presumably through the oxidative action of the myeloperoxidase (MPO)–hydrogen peroxide system (17). In the respiratory distress syndrome of adults and premature infants, α_1 -PI in BAL fluids was found to be in great part inactive (3, 6, 9, 10), and the activity of the α_1 -PI could be restored by treatment with methionyl sulfoxide peptide reductase in the presence of dithiothreitol (DTT) (9). Oxidative inactivation of the elastase inhibitory capacity of α_1 -PI may also play a critical role in the pathogenesis of emphysema (18, 19).

Oxidant generation and oxidant-mediated injury have been documented in experimental animal systems as well. In rabbits, extracellular catalase in inflamed lungs was inhibited by 3-amino, 1,2,4 triazole (AT), which can occur only when H₂O₂ is present

Dr. Rice is with the Harborview Medical Center, Seattle, WA.

This is Publication No. 3691-IMM from the Research Institute of the Scripps Clinic.

Received for publication 6 November 1984 and in revised form 1 April 1985.

J. Clin. Invest.

© The American Society for Clinical Investigation, Inc.

0021-9738/85/09/1182/11 \$1.00

Volume 76, September 1985, 1182–1192

1. *Abbreviations used in this paper:* ACE, angiotensin converting enzyme; α_1 -PI, α_1 -protease inhibitor; AT, 3-amino, 1,2,4 triazole; BAL, bronchoalveolar lavage; CT, chloramine T; DTT, dithiothreitol; ELISA, enzyme-linked immunosorbent assay; FNLP, formyl-norleu-leu-phe; HNE, human neutrophil elastase; HSA, human serum albumin; MNE, monkey neutrophil elastase; MPO, myeloperoxidase; MS, methionine sulfoxide; NMP, normal monkey plasma; PK, prekallikrein; PMA, phorbol myristate acetate; SBTI, soybean trypsin inhibitor; SDS, sodium dodecyl sulfate.

(20–23), and this inhibition was prevented by the use of antioxidant or by neutrophil depletion (13). A marked fall in intracellular glutathione, a major defender against oxidants, was also found in these animals, and was likewise preventable by antioxidant or neutrophil depletion (13). In rats, inactivators of H_2O_2 , scavengers of oxygen free radicals, and chelation of Fe^{3+} (to prevent formation of hydroxyl radicals), all inhibited the development of pulmonary edema and injury of endothelial cells (8, 11, 12). In other studies, perfusion of stimulated normal human neutrophils into the vasculature of in vitro perfused rabbit lungs produced vascular injury, while similar treatment with neutrophils of patients with chronic granulomatous disease were without effect (7).

Although information regarding the biochemical aspects of the inflammatory process may be obtained from studies in rodents and rabbits, many biochemical facets of their plasma proteins and inflammatory cells are dissimilar to those of human beings. We have therefore conducted studies on the development of inflammatory pulmonary disease in subhuman primates. In particular, we have measured the presence or generation in situ of nonleukocytic and leukocytic proteases such as neutrophil-elastase, leukocytic oxidants, arachidonate metabolites, angiotensin-converting enzyme, and components of the complement and contact system. The relationship of the concentrations or presence of these mediators in the lung and inflammatory injury will be assessed. The primate model also provides a system for future studies on the use of therapeutic agents for blocking or controlling specific components of the inflammatory process.

Methods

Proteins and antibodies

Human prekallikrein (PK) was a gift from Dr. Yoshio Hojima of our laboratory and was radiolabeled by the chloramine T (CT) method (24) to a specific activity of $10 \mu\text{Ci}/\mu\text{g}$. Antisera to human neutrophilic elastase was prepared in our laboratory as described previously (6) and was found to crossreact with elastase from rhesus monkey neutrophil granules. α_1 -PI was isolated from monkey plasma using the methods developed for human α_1 -PI purification (9) and antisera was produced in a goat by biweekly multiple intradermal injections of $60 \mu\text{g}$ protein in minced acrylamide gel plus complete Freund's adjuvant (Difco Laboratories, Detroit, MI). Antibody produced after 10 wk showed a single coalescent line in double immunodiffusion tests with monkey plasma and purified monkey α_1 -PI.

Human catalase was prepared by a method based on that of Stansell and Deutsch (25). Human erythrocytes sedimented through gelatin were washed three times with 0.9 percent NaCl at room temperature. Washed cells were stored frozen at -20°C until used (1 wk). 900 ml of packed cells was lysed by adding an equal volume of 0.2% saponin in cold water and stirring at 4°C for 1 h. The lysate was dialyzed against 3 mM sodium potassium phosphate, pH 7.0 ("starting buffer"), and then added to 750 ml of DEAE-Sephacel beads (Pharmacia, Uppsala, Sweden). After shaking for 1 h at 4°C , the mixture was poured into a scintered glass funnel and washed until the red color was reduced to light pink. The gel was then packed in a 5×28 -cm column and washing with starting buffer continued until $OD_{290} = 0.050$. Proteins were then eluted with a gradient of 2 liters each of starting buffer and starting buffer + 0.2 M NaCl. Catalase was eluted from 6 to 8 mmho. The region was pooled and brought to 50% saturation with solid ammonium sulfate. Proteins precipitating at this concentration were discarded and the supernatant brought to 70% saturation with ammonium sulfate. The proteins precipitated at this concentration were dialyzed against 0.01 M sodium acetate, pH 5.0, and then applied to a column containing 230 ml of SP-Sephadex C-50. The column was washed at a flow rate of 90 ml/h and then eluted with a gradient of 2 liters each of 0.01 M acetate, pH 5.0 (0.9 mmho), and 0.01

M acetate plus 0.4 M NaCl (35 mmho). Three overlapping peaks of catalase activity were found eluting from 9.4 to 23 mmho. Catalase in the three peaks could not be distinguished by electrophoresis or by gel filtration. In the presence of mercaptoethanol, all showed >95% homogeneity in a band at $\sim 60,000$ mol wt. Analysis on G-150 gel filtration resulted in a mol wt of 110,000. A total of 137 mg of human catalase was recovered. Antibody to human catalase was produced in rabbits by the weekly subcutaneous injection into multiple sites of $100 \mu\text{g}$ of purified protein in complete Freund's adjuvant. Monospecific antibody, produced by the fourth week, was found to crossreact with catalase from monkey erythrocytes.

Reagents

Formyl-norleu-leu-phe (FNLP) (Vega, Tucson, AZ) was dissolved 50 mg/ml in dimethyl sulfoxide and stored at -20°C until used. PMA (Midland Corp., Brewster, NJ) was dissolved 10 mg/ml in dimethyl sulfoxide and stored in the dark at -70°C until used. AT (Sigma Chemical Co., St. Louis, MO) was dissolved in saline at a concentration of 100 mg/ml just before use. Human serum albumin (HSA) was from Armour Pharmaceutical Company ("Albuminar-25", Kankakee, IL).

Experimental protocol

Rhesus monkeys, *Macacca mulatta*, 6–10 kg were used in these studies. All animals were cared for according to the principles of the Guide for the Care and Use of Laboratory Animals,² and were under the continuous care of a veterinarian. Anesthesia was induced with ketamine hydrochloride (Parke-Davis, Morris Plains, NJ) 8–10 mg/kg intravenously, and maintained with 3–5 mg/kg every 2 h until the completion of the experiment. The animals were intubated with a cuffed 6 Fr. orotracheal tube.

Preliminary experiments were performed in both rabbits (13) and monkeys to determine what inciting agents would yield reproducible, nonlethal, localized inflammatory injury. Two leukocytic stimuli, FNLP and PMA, were chosen. Both have been shown in numerous studies to be chemoattractants, to cause aggregation of polymorphonuclear leukocytes (PMNs) and release of oxidants and proteases, and to cause lung injury when used in vivo (7, 13, 26–33). Optimal doses and times of lavage sampling were likewise determined in preliminary experiments. $200 \mu\text{g}/\text{kg}$ FNLP, instilled intrabronchially, was found to cause a moderate infiltration of PMNs into the lung after 5 h. Because <10% of the formyl peptide is present in lung tissue at that time (13), a second dose of FNLP was given at 5 h to induce the release of oxidants and proteases from the accumulated cells that would be measured in lavage samples taken 1 h later. Intrabronchial instillation of $10 \mu\text{g}/\text{kg}$ PMA was found to cause a more rapid and severe leukocytic infiltration. This second stimulus was used to determine whether a different response of mediator systems could be detected. Fig. 1 shows a typical infiltration of leukocytes and the interstitial edema seen after PMA challenge.

Standard anterior–posterior chest radiographs were obtained and reviewed before beginning the experiment. Blood was drawn from the femoral artery into citrate for cell counts, hematocrit, and plasma samples. 5 ml of blood was centrifuged and 2 ml of plasma was frozen immediately. Two pulmonary lavages were performed on each animal as described below. The recovered BAL fluids, referred to as the "untreated" samples throughout this paper, served as controls for each animal. Pulmonary injury was then induced by one of two means: FNLP, $200 \mu\text{g}/2$ ml per kg in saline or PMA, $10 \mu\text{g}/2$ ml per kg, in saline containing 0.01% Evan's Blue dye plus 0.01% HSA was instilled intrabronchially above the carina with the animals held in an upright position. After 4.5 h, x-rays were again taken of the animals that received FNLP. AT in saline, in a concentration of 1 g/10 ml per kg or saline alone (10 ml/kg) was administered intraperitoneally. A second intrabronchial instillation of FNLP (same dose as first time) was then made, incorporating 0.01%

2. Department of Health, Education and Welfare Publication (NIH) 78-23, revised 1978, Office of Science and Health Reports, DRR/NIH, Bethesda, MD 20205.




Figure 1. Photomicrograph of the lung of a rhesus monkey obtained 5 h after instillation of PMA. Accumulation of neutrophils in alveolar septae and spaces are apparent along with amorphous material with tinctorial characteristics of edema fluid. $\times 156$.

Evan's Blue-HSA. 1 h later (5.5 h after the first FNLP instillation) terminal lavages and blood samples were taken. PMA-challenged animals were x-rayed and given AT or saline intraperitoneally 3 h after injury. Terminal samples of blood and BAL fluids were taken 1 h later. Control animals were treated identically to the FNLP animals except that FNLP was omitted from the instillation fluid and neither AT nor saline was given intraperitoneally.

Lavage procedure (untreated, e.g., preinjury, samples): 2 min before lavage, 2 ml of saline containing 100 $\mu\text{g/ml}$ HSA and 7.1 μCi ^{125}I -PK were instilled via a soft-tipped cannula passed through the endotracheal tube into the lower respiratory tract and advanced until wedged. The cannula was left in place for 2 min and then 15 ml of saline was instilled and immediately recovered. Because of the low volumes of recovery in normal, nonedematous lungs, a second 15-ml lavage was often done in the same site. The volume recovered was recorded and 0.5-ml samples were added to 0.5 ml of methanol or 0.5 ml isopropanol and frozen immediately on dry ice for later analysis of leukotrienes B_4 and C_4/D_4 , respectively. The remainder of the lavage fluid was centrifuged in 1.5-ml capacity polypropylene tubes for 5 s in a Beckman Microfuge B (Beckman Instruments, Inc., Palo Alto, CA) adding sequential 1.5-ml aliquots to the same tube until all the sample had been centrifuged. The supernatant BAL fluids were frozen immediately on dry ice. To lyse and remove any contaminating erythrocytes, the cell pellet was resuspended in 25 μl of Hanks' buffer. (If two lavages had been done in the same site, the two cell pellets were combined at this point.) 1 ml of cold distilled water was added and the sample mixed continually for 30 s while being kept on wet ice. Isotonicity was then restored with 333 μl of 3.5% NaCl. The tubes were centrifuged for 5 s; then the supernatant was removed and discarded. 500 μl of distilled water was added to the cell pellet and the sample was frozen on dry ice for later analysis. The terminal lavage procedure differed only as follows: After the initial lavage was taken (2 min after ^{125}I -PK instillation), a second cannula was inserted through

the endotracheal tube into a different lobe; the first cannula was then removed. 2 ml of saline containing 100 $\mu\text{g/ml}$ HSA, 7.1 μCi ^{125}I -PK and 55 μl of immunopurified anti-human neutrophil elastase (HNE) were instilled. 2 min later a 15-ml saline lavage was performed. The two samples thus obtained were processed as above, only the washed cell pellets as well as the supernatants were kept separate. Animals given PMA had their terminal lavages performed with 7.5 ml of saline instead of 15 ml.

Pathophysiologic changes in the lungs were difficult to assess in the monkeys because, in order to retain the animal for further experiments, injury was limited to a segment of one lobe. Hence it was not possible to carry out analyses of lung weight or functional assays such as changes in PO_2 and PCO_2 . Although these latter were attempted, changes were minimal and varied greatly with the time after administration of ketamine. Resistance and compliance assays were not attempted owing to the limited zone of injury. To assess the presence of inflammation, we therefore performed open biopsies at 3 and 6 h on the PMA- and FNLP-treated monkeys, respectively, and at 24 h in both groups. Untreated control monkeys were also biopsied. In addition, radiographs were taken at the same times noted above, as well as measurement of total protein and cells in the BAL fluid taken at the same times. It was also interesting that the volume of lavage fluid aspirated after instillation of saline solution was greater in injured lungs as noted below. Nevertheless, one of the drawbacks of these studies in living primates is the lack of quantitative estimates of lung damage.

Protein. Protein contents of plasma, BAL fluid, and cell pellets were measured by the method of Lowry et al. (34).

Enzyme assays. Catalase activity in BAL samples was measured by the method of Aebi (35) by following the disappearance of H_2O_2 spectrophotometrically. MPO was measured as follows: Wells of a flat-bottom 96-well microtiter plate were filled with 50 μl of sample in 0.1 M citrate buffer, pH 4.2. 50 μl of citrate buffer containing 1 mM 2,2'-azino-di-(3-ethylbenzthiazoline sulfonic acid) (ABTS, Sigma Chemical Co., St. Louis,

MO) and 0.03% H₂O₂ was added. Absorbance at 414 nm was read at two time points within the first 15 min on a Titertek Multiskan photometer (Flow Laboratories, McLean, VA) and the data were expressed as the change in optical density per minute per milliliter of sample. β -Glucuronidase was assayed by a modification of a previously described method (36): In 1.5-ml polypropylene tubes, 20 μ l of sample or standards (0.05–0.6 U standard β -glucuronidase, Calbiochem-Behring Corp., La Jolla, CA) were added to 30 μ l of 0.1 M acetate buffer, pH 4.5. 50 μ l of phenolphthalein glucuronide (Sigma Chemical Co.), 1.3 mg/ml in acetate buffer, was added and the tubes were capped and incubated overnight at 37°C. The reaction was stopped and developed with the addition of 100 μ l of 0.5 M glycine-NaOH buffer, pH 10.7. The samples were transferred to a 96-well flat-bottom microtiter plate and absorbance at 540 nm read on a Titertek Multiskan (Flow Laboratories). A linear relationship existed between the absorbance and the amount of β -glucuronidase added. Amidolytic activity was measured using a final concentration of 100 nM *H*-D-pro-phe-arg-pNA (S2302, Kabi Diagnostica, Stockholm, Sweden) in 0.1 M tris buffer pH 8.0. Samples were incubated with the substrate for 30 min at 37°C and the reaction stopped with 1 N acetic acid. Absorbance at 405 nm was read and the data were expressed as the change in absorbance per hour per 50 μ l of samples. Corn inhibitor, used in these studies as a specific inhibitor of active Hageman factor (37) was a gift from Dr. Yoshio Hojima of this laboratory. Soybean trypsin inhibitor (SBTI, Worthington Biochemical Corp., Freehold, NJ) was used as a kallikrein inhibitor.

Immunologic assays

Levels of monkey α_1 -PI protein were determined by the radial immunodiffusion method of Mancini et al. (38) using antisera prepared as described above. Catalase levels were determined by an enzyme-linked immunosorbent assay (ELISA) utilizing an antibody to human catalase that exhibited good crossreactivity with the monkey protein, performed as follows: 96-well flat-bottom polystyrene microtiter plates (Nunc Immuno Plate II, Thomas Scientific, Philadelphia, PA) were used for these assays. 50 μ l of immunopurified rabbit anti human catalase F(Ab')₂ antibodies absorbed with Protein A-Sepharose beads (Pharmacia, Uppsala, Sweden) to remove Fc fragments and residual intact IgG and diluted to 20 μ g/ml in phosphate-buffered saline (PBS) were placed into each well. The plate was covered and incubated overnight at 4°C on an orbital rocker (Nutator, McGaw Park, IL) to coat the wells. The plate was emptied, and each well washed with 150 μ l of 2% bovine serum albumin (BSA, Sigma Chemical Co.) in PBS followed by four washes of PBS alone. 50 μ l of standards ranging from 0 to 0.6 μ g/ml purified human catalase or 50 μ l of unknown samples was added to each well. The plate was incubated on the Nutator for at least 1 h and washed with BSA and PBS as above. 50 μ l of immunopurified IgG rabbit anti-human catalase diluted in 0.1% BSA in PBS was added to each well. After a minimum incubation period of 20 min at room temperature, the plate was again washed with BSA and PBS. 50 μ l of a 1:200 dilution in 2% BSA/PBS of Protein A-glucose oxidase (made by the addition of 10 mg of glucose oxidase to 4 mg of protein A) was then added to each well. After a 30-min incubation on the Nutator at room temperature, the wells were emptied and washed eight times with PBS. To each well was then added 100 μ l of a substrate solution consisting of 4 mg of ophenylene diamine, 9 ml citrate-phosphate buffer, pH 5, 73.4 μ g of horseradish peroxidase and 1.1 ml of 18% glucose. The plate was incubated, uncovered, in the dark for up to 2 h at room temperature until a dark yellow color developed in the most concentrated standards. The reaction was stopped by the addition of 100 μ l of 5 N H₂SO₄ to each well. The plate was read on the Titertek Multiskan using a filter of 492 nm. A linear relationship existed between the concentration of catalase and the absorbance.

Concentrations of monkey neutrophil elastase (MNE) complexed with α_1 -PI were also determined by an ELISA assay. Standards were made by taking a known quantity of monkey elastase and complexing it completely with monkey α_1 -PI. Data were expressed as μ g/ml of complexed MNE. Plates, incubations, volumes, washes, and reading of the plate are as described above. The reagents used for the MNE- α_1 -PI assay

were (a) immunopurified rabbit anti human neutrophil elastase F(Ab')₂ absorbed with insolubilized Protein-A and diluted to 10 μ g/ml with PBS, (b) standard complexes ranging from 20 to 150 ng/ml MNE or BAL fluids to be tested, (c) immunopurified goat anti-human α_1 -PI, (d) rabbit anti-goat Fc fragment (Cappel Laboratories, West Chester, PA) diluted 1/50, (e) Protein A-glucose oxidase, (f) substrate solution, (g) 5 N H₂SO₄ to terminate the reaction. A linear relationship existed between the absorbance at 492 nm and the log of the amount of complexed monkey elastase.

α_1 -PI activity. α_1 -PI activity in the monkey plasmas and BAL fluids was measured by inhibition of porcine pancreatic elastase (Sigma Chemical Co.) (39). A pool of normal monkey plasma (NMP) containing 1.3 mg/ml α_1 -PI was used as a standard. Reactivation of oxidized α_1 -PI in the presence of the reducing agent dithiothreitol (DTT, Sigma, St. Louis, MO) and methionine sulfoxide (MS) peptide reductase (kindly provided by Dr. Nathan Brot, Hoffmann-La Roche, Inc., Nutley, NJ) was performed as described previously (9), based on the method of Abrams et al. (40). In the current studies, a single incubation time of 2 h at 37°C was used, with triplicate samples containing DTT, DTT plus MS reductase, or sample + buffer being analyzed.

Glutathione. Total glutathione (reduced and oxidized) was determined by following spectrophotometrically the disappearance of NADPH in the presence of glutathione reductase and 5,5'-dithiobis (2-nitrobenzoic acid) (41).

Cleavage of ¹²⁵I-PK. To measure in situ protease activity, ¹²⁵I-PK was instilled 2 min before lavage as described above under *Experimental protocol*. Aliquots of BAL fluids were counted and sample volumes containing equal quantities of ¹²⁵I were applied to 9% polyacrylamide gels containing 0.1% sodium dodecyl sulfate (SDS) (42). All samples were run both with and without reduction with β -mercaptoethanol. After electrophoresis, the gels were fixed, stained, and dried, and visualization of the position of the radiolabel obtained by autoradiography using X-Omat AR film (Eastman Kodak XAR-5, Rochester, NY). Quantitation of the cleavage of the ¹²⁵I-PK is described in the Results section.

Leukotrienes. Analyses for leukotrienes LTB₄, LTC₄, and LTD₄ were performed as described (43).

Angiotensin-converting enzyme (ACE). A fluorometric assay (44) was used, employing porcine ACE (Calbiochem-Behring Corp.) as a standard. Data are expressed as nanomoles L-histadyl-L-leucine converted per minute per milliliter of sample.

In vitro inhibition of MPO by AT in the presence of H₂O₂. To determine whether inhibition of MPO activity by AT was H₂O₂ dependent, a system similar to that described by Cohen and Hochstein (45) was developed which allowed a slow and continuous diffusion of H₂O₂ into a mixture of MPO plus AT. 250 μ l of 30% H₂O₂ (Fisher Scientific Co., Tustin, CA) was placed into a 1.5-ml capacity conical, plastic micro tube (Sarstedt, Princeton, NJ) which had had its cap removed. The H₂O₂ tube was placed into a 20-ml capacity polyethylene scintillation vial (American Scientific Products, Irvine, CA). 49.5 mU/500 μ l of purified human MPO (Behring Diagnostics, La Jolla, CA) in 1 mg/ml BSA in Tris-buffered saline was added to the scintillation vial along with 50 μ l of either water or 250 mM AT. The vial was capped and allowed to incubate for 2 h in a shaking 37° water bath. The H₂O₂ containing tube was then removed and the solution assayed for its MPO activity as described above. The system was found to result in the diffusion of \sim 1 μ mol of H₂O₂ (measured by the method of Hyslop and Sklar [46]) into the 500 μ l sample per hour.

Cell counts were performed by standard clinical methods.

Statistical analyses were performed by the use of Student's two-tailed *t* tests.

Results

Assessment of injury produced by instillation of formylated peptide (FNLP) and phorbol ester (PMA)

Several methods were used to determine the presence of inflammatory injury. The protein content of each BAL fluid was mea-

sured as an index of transudation of protein from the vascular space into the pulmonary tissue. 27 lavage samples taken from 16 monkeys before the administration of the injurious agent had a mean protein level of 0.51 mg/ml. 1 h after the second instillation of FNLP, the mean protein content was 3.74 mg/ml. Protein in lavages taken 3 h after treatment with PMA had a mean level of 6.64 mg/ml (see Table I). These statistically significant ($P < 0.001$) increases were not seen in the control animals that had a mean 6-h protein level of 0.85 mg/ml ($P = 0.3$). A qualitative indicator of pulmonary edema was found in the recovery volumes of lavage fluid. Expressed as milliliter of fluid recovered per milliliter instilled, the mean values were 0.10 ml for the preinjury samples, 0.19 ml after FNLP treatment, 0.68 ml after PMA treatment (Table I), and 0.07 ml in control monkeys.

In order to insure that the lavage was obtained in areas of induced inflammation, Evan's Blue dye was included in the instillation of PMA or the second dose of FNLP. Blue dye was recovered to varying degrees in almost all BAL samples indicating that the areas lavaged had been exposed to the injurious agent.

Additional evaluation of the extent of injury was obtained by x-ray studies of the lungs before and after exposure to the inflammatory agents. All preexperiment chest radiographs were normal. Films taken 4.5 h after FNLP treatment or 3 h after PMA instillation, revealed patchy alveolar infiltrates, confined to a single lobe (Fig. 2). The remainder of the lung was clear. Additional x-rays obtained 24 h later demonstrated the persistence of the infiltrate, and in some animals, an increase in density, consistent with early lobar consolidation.

In some animals open lung biopsies were taken 1 h after the second instillation of FNLP and 3–4 h after administration of PMA. The tissues revealed patchy accumulation of neutrophils in the alveolar septae and alveolar spaces. This was accompanied by modest amounts of protein-rich edema fluid (Fig. 1). These changes were considerably more severe in the monkeys treated with PMA.

Peripheral white blood cell counts, starting from an average of $6.0 \times 10^3/\text{mm}^3$, rose in most cases, to an average of $17.1 \times 10^3/\text{mm}^3$ 5.5 h after FNLP treatment and to $12.5 \times 10^3/\text{mm}^3$ 4 h after PMA challenge.

Determination of protease activity in BAL fluid

Assays of elastase activity on the synthetic substrate methoxysuc-ala-ala-pro-val-pNA of the BAL fluids were negative. To determine if neutrophil elastase was inactivated by binding to α_1 -PI, an ELISA technique was developed to measure the amount




Figure 2. Chest radiographs of monkey before (left) and 3 h after (right) intrabronchial instillation of PMA. A patchy infiltrate can be seen in the right lower lobe after treatment.

of elastase- α_1 -PI complex in each lavage sample (see Methods). The mean value obtained in 13 BAL fluids of untreated monkeys was $0.47 \mu\text{g}/\text{ml}$. 11 BAL samples taken at the end of the experimental period from animals treated with FNLP had an average elastase level of $0.99 \mu\text{g}/\text{ml}$. BAL fluid taken similarly from PMA-treated animals had an average of $1.23 \mu\text{g}/\text{ml}$ elastase complexed to α_1 -PI (Table II). To confirm that all the elastase present in the samples was complexed to α_1 -PI, duplicate ELISA assays were performed with additional monkey α_1 -PI added to each sample before assay. No significant increases were observed, indicating that all active elastase was already bound to α_1 -PI.

As assays of enzymatic activity in lavage fluids may not accurately reflect proteolytic activity in the bronchoalveolar space, we attempted to assess the presence of proteolytic activity in situ. To accomplish this, we instilled ^{125}I -labeled PK as a probe of protease activity into the bronchoalveolar space 2 min before the lavage procedure. ^{125}I -PK is cleaved by active Hageman factor, neutrophil elastase and other proteolytic enzymes. The ^{125}I -PK recovered in the lavage fluid was subsequently analyzed on SDS-polyacrylamide gels for cleavage of the protein. The extent of cleavage was rated with a numerical value of 0–4, with 0 reflecting no cleavage; 1, up to 30% of the PK cleaved; 2, ~30–60% cleaved; 3, ~60–90% cleaved; and 4, indicating complete disappearance of the native molecule. Representative samples are shown in Fig. 3. 11 of 12 samples taken before injury showed 0 cleavage. After injury with FNLP or PMA, cleavage of PK was found to be variable. The average of FNLP-injured lung lavages was 1.5 while PMA-injured lung lavages had a mean cleavage value of 1.0 (Table II).

An attempt was made to assign specificity of the protease activity by including anti-elastase antibody with the radiolabeled PK in lavage samples taken from a different site in each animal. Although it could be shown in vitro that antibody to elastase could effectively block cleavage of PK induced by elastase, no consistent blockage was seen in vivo. As was the case in the absence of antibody, cleavage of PK in the presence of antibody was variable. (See representative samples in Fig. 3.) Again, rated on a scale of 0–4, the average cleavage value of lavages performed in the presence of antibody to elastase in FNLP-treated animals was 0.75 and in PMA-injured animals was 1.1 (Table II). Cleavage patterns of PK as analyzed on SDS-polyacrylamide gels run in the presence or absence of β -mercaptoethanol differed from those obtained with purified neutrophil elastase in vitro but were

Table I. Evidence of Pulmonary Injury

Stimulus	BAL fluid protein	Lavage recovered per ml instilled
	mg/ml	ml
Untreated	0.51 ± 0.14 ($n = 27$)	0.10 ± 0.01 ($n = 23$)
Control	0.85 ± 0.24 ($n = 8$)	0.07 ± 0.01 ($n = 8$)
FNLP	3.74 ± 1.12 ($n = 16$)	0.19 ± 0.03 ($n = 12$)
PMA	6.64 ± 2.12 ($n = 10$)	0.68 ± 0.11 ($n = 10$)

Data are expressed in this, and all subsequent tables, as mean \pm standard error of the mean.

Table II. Protease Activity in BAL Fluids

	MNE complexed to α_1 -PI $\mu\text{g/ml BAL}$	In situ cleavage of PK*		Amidolytic activity \S $\Delta A_{405}/\text{h}$	β -Glucuronidase activity U/ml
		-Ab \ddagger	+Ab \ddagger		
Untreated	0.47 \pm 0.09 (n = 13)	0.04 \pm 0.04 (n = 12)	—	—	0.85 \pm 0.14 (n = 9)
FNLP	0.98 \pm 0.48 (n = 11)	1.50 \pm 0.57 (n = 8)	0.75 \pm 0.75 (n = 4)	0.07 \pm 0.02 (n = 11)	4.36 \pm 1.23 (n = 15)
PMA	1.22 \pm 0.73 (n = 10)	1.00 \pm 0.63 (n = 5)	1.1 \pm 0.25 (n = 5)	0.22 \pm 0.08 (n = 8)	8.25 \pm 3.05 (n = 10)

* 0 = no cleavage; 4 = complete cleavage (see text). \ddagger Antibody to elastase. \S On substrate *H*-D-pro-phe-arg-pNA.

indistinguishable from patterns obtained by treatment of ^{125}I -PK with active Hageman factor.

50- μl samples of all the terminal lavage samples were assayed for amidolytic activity on the synthetic substrate *H*-D-pro-phe-arg pNA. Amidolytic activity of samples from FNLP- and PMA-treated animals is presented in Table II. To determine whether this activity was due to active Hageman factor, lavage samples were preincubated with corn inhibitor, a known specific inhibitor of Hageman factor activity (37). The amount of corn inhibitor used was sufficient to block an amount of HFa that would produce equal cleavage of the synthetic substrate as did BAL fluid. Partial inhibition was seen in most of the cases studied. Kallikrein is also capable of cleaving this synthetic substrate and an attempt was made to assess its potential presence in the lavage samples by the addition of SBTI, an inhibitor of kallikrein, to the samples prior to assay. The amount added was sufficient to block kallikrein in concentrations giving amidolysis equal to that seen with the BAL fluids. Again only partial inhibition was seen. Addition of corn inhibitor and SBTI together failed to block completely the amidolytic activity. It therefore appeared that a portion of the proteolytic activity was attributable to active Hageman factor and to an SBTI inhibitable protease, possibly kallikrein. An additional protease was also present.

The appearance of β -glucuronidase and myeloperoxidase in

the BAL fluids of untreated and FNLP- and PMA-treated animals was measured. β -glucuronidase was found to rise from 0.85 to 4.36 U/ml ($P < 0.04$) and 8.25 U/ml ($P < 0.04$), respectively (Table II). MPO, specific to neutrophils, showed a striking increase from 1.37 OD U/ml \cdot min in untreated to 16.59 ($P < 0.1$) and 30.47 ($P < 0.001$) OD U/ml \cdot min in FNLP- and PMA-treated samples (Table III). 50% of the control animals, which received saline and dilute DMSO, but no FNLP or PMA, showed elevated levels of β -glucuronidase activity after 5 h. MPO activity was mildly elevated in one out of eight control samples.

Assessment of oxidant activity in pulmonary inflammation

A diminished specific activity of α_1 -PI in BAL fluids has been used as an indicator of oxidant generation in vivo with inactive α_1 -PI at its native molecular weight in BAL fluids from human patients with respiratory distress syndrome being reactivated by exposure to reducing agents in the presence of MS peptide reductase (9). In the current experiments the specific activity of α_1 -PI was determined in the lavage samples taken before instillation of FNLP or PMA and in the BAL samples taken at the termination of the experiment. The specific activity of the α_1 -PI fell from a mean of 0.80 in control samples to 0.57 and 0.65 with FNLP or PMA injury, respectively ($P < 0.06$ for both) (Table IV). In samples of four control animals, one showed no inhibition of α_1 -PI activity, two showed moderate inhibition (specific activities of 0.58 and 0.69) and one showed marked inhibition (specific activity of 0.32). An additional comparison was made utilizing plasma samples taken prior to or at the end of the injury period and analyzing the α_1 -PI specific activities in them. These values, also shown in Table IV, were 0.91, 0.89, and 0.86 for untreated, FNLP-treated, and PMA-treated animals,




Figure 3. SDS-polyacrylamide gel of ^{125}I -PK instilled into monkeys. All samples were reduced with β -mercaptoethanol prior to electrophoresis. Control gels on left show ^{125}I -PK alone (A) or with the addition of β -active Hageman factor (B) or HNE (C). Gel D is the pattern obtained with 11 of the 12 untreated monkeys. The remaining gels are representative of post-injury samples in the absence (E-H) or presence (I, J) of antibody (Ab) to HNE. "Cleavage value" is described in text.

Table III. Activities of Catalase and MPO in BAL Fluids

	Catalase		MPO
	Total $\mu\text{g/ml}$	Specific activity	
Untreated (n = 19)	0.37 \pm 0.08	0.97 \pm 0.16	1.37 \pm 0.55
FNLP (n = 9)	0.70 \pm 0.22	0.57 \pm 0.10	16.59 \pm 12.91
FNLP + AT (n = 6)	0.34 \pm 0.05	0.04 \pm 0.02	0.85 \pm 0.38
PMA (n = 6)	1.29 \pm 0.26	1.08 \pm 0.20	30.47 \pm 6.23
PMA + AT (n = 4)	1.44 \pm 1.17	0.49 \pm 0.25	0.60 \pm 0.37

Table IV. Specific Activity of α_1 -PI

	BAL fluid		Plasma	
	Total	Specific activity	Total	Specific activity
	$\mu\text{g/ml}$		$\mu\text{g/ml}$	
Untreated	12.9±4.3 (n = 11)	0.80±0.04 (n = 11)	1043±37.5 (n = 3)	0.91±0.06 (n = 3)
FNLFP	31.9±10.4 (n = 12)	0.57±0.06 (n = 12)	1504.7±80.3 (n = 8)	0.89±0.04 (n = 8)
PMA	112.4±32.8 (n = 9)	0.65±0.06 (n = 9)	1088.4±70.4 (n = 4)	0.86±0.02 (n = 4)

respectively. These data suggest inactivation of α_1 -PI is occurring in the lung specifically.

To determine whether the observed inactivation of α_1 -PI was due to oxidative injury, we measured the increase in elastase inhibitory capacity of several BAL samples having specific activities of <0.80 after their incubation with the reducing agent DTT in the presence or absence of MS reductase. The results, presented in Table V, show complete reactivation of α_1 -PI in four out of five samples and partial restoration of activity in the fifth case. It is of interest to note that DTT alone resulted in significant, but not always complete, reactivation of the α_1 -PI in all cases.

The inactivation of catalase in sites of pulmonary inflammation in animals treated with aminotriazole has also been used as an indicator of oxidant (H_2O_2) generation in vivo (13). In order to obtain values of specific activity, we measured total catalase in the lavage fluids immunologically using an ELISA assay and catalase activity by disappearance of H_2O_2 spectrophotometrically. Before injury, the average catalase specific activity of 19 lavage samples tested was 0.97 with the mean concentration being 0.37 $\mu\text{g/ml}$. The average specific activities of the terminal lavage samples is given in Table III. With both FNLFP and PMA-induced injury, a significant reduction (P

Table V. Reactivation of α_1 -PI by Exposure to DTT±MS Reductase

Sample no.	BAL 1*	BAL 2	BAL 3	BAL 4	BAL 5	Oxidized NMP
‡ Sample alone (initial activity)	1.5	29.5	12.9	22.2	6.5	51.2
‡ Sample + DTT	3.4	40.0	14.6	35.5	8.3	197.0
‡ Sample + DTT + MS reductase	6.3	49.1	27.0	34.8	9.9	715.8
§ Total α_1 -PI protein in sample	12.5	52.2	24.5	28.8	10.6	1300.0

All values are micrograms/milliliter.

* PMA injured, all others FNLFP.

‡ Activity of α_1 -PI as determined by porcine elastase inhibition.

§ Measured by radial immunodiffusion.

Table VI. Inhibition of MPO by AT in the Presence of H_2O_2

	mU/ml
MPO	503
MPO + AT	484
MPO + AT + H_2O_2	8
MPO + H_2O_2	210

< 0.001 and $P < 0.1$) was seen in monkeys given aminotriazole. Specific activity for FNLFP-treated animals fell from 0.57 to 0.04 and for PMA-treated animals from 1.08 to 0.49. It is interesting to note that even in the absence of AT, there was a drop ($P < 0.1$) in specific activity of those animals receiving FNLFP.

The inhibitory effect of AT in the presence of H_2O_2 extended to activity of myeloperoxidase as well. As shown in Table III, MPO activity fell from 16.59 to 0.85 OD U/min · ml in the case of FNLFP-injured BAL fluids when AT was added ($P = 0.3$) and from 30.47 to 0.60 OD U/min · ml with PMA injury ($P < 0.02$). To determine whether this inhibition was dependent upon H_2O_2 , studies were conducted in vitro which allowed for the slow diffusion of H_2O_2 into a solution of myeloperoxidase and aminotriazole (see Methods). As shown in Table VI, >98% of the MPO activity was inhibited by AT in the presence of H_2O_2 as compared to <4% inhibition in its absence. An inhibition of 58% was seen using H_2O_2 diffusion in the absence of AT, but this effect did not occur when crude cell lysates were used as a source of MPO. These studies indicate that MPO inhibition by AT, like that of catalase, is dependent upon the presence of H_2O_2 . The study validates the use of the inhibition of MPO by AT as an assay of H_2O_2 generation in the whole animal.

The level of the antioxidant glutathione was measured in the cell pellets recovered from the lavage samples to assess the possibility that increase in oxidant formation might be reflected by a decrease in intracellular antioxidants as is the case in our previous studies in rabbits (13). The results, shown in Table VII, suggest that this was not the case. Normalized to mg of cell protein, total glutathione levels were found to be variable, resulting in mean values not differing significantly ($P > 0.4$) from the untreated value of 37.5 nM/mg. Treatment of the animals with AT did not appear to affect the glutathione levels ($P > 0.4$).

Leukotrienes. Assays were performed for detection of the leukotrienes LTB_4 , LTC_4 , LTD_4 in 11 lavage samples. None were detected. The limits of sensitivity of the method would

Table VII. Glutathione Content of Cellular Component of Lavage Fluids

	nmol/mg cellular protein
Untreated (n = 14)	37.5±7.0
FNLFP (n = 10)	25.9±5.7
FNLFP + AT (n = 6)	33.7±5.0
PMA (n = 6)	31.8±8.0
PMA + AT (n = 3)	31.1±16.7

have allowed for the detection of 10 pmol/ml of LTB₄ and 40 pmol/ml of LTC₄ or LTD₄. Recovery of standards added to several lavage fluids, at the moment the lavage fluid was obtained, was from 74 to 100%. In one sample, from an animal receiving FNLP, an unknown peak eluted from high performance liquid chromatography in the position corresponding to 15-hydroperoxyeicosa tetraenoic acid. The rate of metabolism or loss of leukotrienes in the tissues is not known.

ACE. The ACE level was compared in plasma samples taken at the beginning and end of the experiment in several of the monkeys. No significant differences were seen, with starting levels averaging 158.7 nmol L-histidyl-L-leucine converted per min per ml of plasma and final values averaging 163.5.

Discussion

In these studies we utilized the rhesus monkey to establish an animal model of pulmonary inflammation and analyze the biochemical effector mechanisms responsible for the injury. Injury was induced by the intrabronchial instillation of the *N*-formylated peptide, FNLP, or the phorbol ester, PMA. Control animals which received intrabronchial instillation of saline and diluted dimethyl sulfoxide were also studied. Plasma and BAL samples taken before and after inflammation developed served as the source of material for biochemical analyses of mediating systems potentially involved in the development of inflammation.

The control animals, receiving only saline and diluted dimethyl sulfoxide, which were included in these studies, showed little evidence of pulmonary inflammation although some biochemical evidence of changes appeared in some of the samples. It is not unreasonable to expect some response from repeated instillation of fluids and lavaging and the variability of the responses suggests that some animals are more sensitive than others. Several investigators have used repeated saline lavages as a method of inducing pulmonary inflammation (47, 48), although the volumes used were much greater than those used in the current studies. It is not our intent in these studies to prove that the inflammatory response invoked is entirely due to FNLP or PMA but rather to establish a model of pulmonary inflammation induced by a defined, reproducible, stimulus and mode of introduction into the animal.

The data show that the protocol used did induce moderate to severe pulmonary inflammation in the experimental animals (Table I). The amount of protein in the BAL fluids, as well as the quantity of fluid recovered, increased substantially. Mortality after PMA injury was 33% by 5 h. None of the monkeys receiving FNLP died. X-rays of the lung fields revealed inflammatory infiltrates (Fig. 2), and histologic examination of open lung biopsies confirmed the presence of leukocytic infiltration (Fig. 1), which consisted of predominantly neutrophils at 4–6 h after instillation of FNLP or PMA, and mononuclear cells at 24 hours.

Analyses of the BAL fluid revealed the presence of both protease and oxidant activity. Studies to identify neutrophil elastase were prompted by the earlier findings of this enzyme in BAL fluids of patients with inflammatory disease of the lung (3, 6, 10) and of the presence of leukocytic proteases in BAL fluids of rabbits (13). The immunologic assay described in this paper enabled us to detect from 0.04–5.5 µg/ml of neutrophil elastase, all of which was complexed to and inactivated by α₁-PI. The concentration of α₁-PI present in the BAL fluids was approxi-

mately that which would be present due to the transudation of plasma proteins from the vascular space. It was possible that elastase, once secreted, remained bound to the cells and connective tissue lining the alveoli and was not removed by our lavaging procedure. This possibility is suggested by the relatively low levels of neutrophil elastase present in BAL fluids of stimulated monkeys compared with levels of myeloperoxidase and β-glucuronidase which are also constituents of the azurophil granules of neutrophils. It was for these reasons that we attempted to show protease activity in situ by instilling and retrieving ¹²⁵I-labeled PK as a probe. The ¹²⁵I-PK was instilled into two different lobes of each monkey, one in the presence of antibody to elastase (which could be shown in vitro to block to cleavage of ¹²⁵I-PK by elastase) and one in the absence of antibody. Although good cleavage was seen in some samples, no statistical difference could be seen when antibody to neutrophil elastase was present (Fig. 3 and Table II). Furthermore, the pattern of cleavage fragments of the ¹²⁵I-PK obtained was not the same as that seen with elastase in vitro. This was especially evident in samples run in the absence of the reducing agent β-mercaptoethanol (data not shown). Other enzymes, such as active Hageman factor (49–51) and a mast-cell derived protease (52) are also known to cleave PK. The failure of the specific inhibitors from corn and soybean to completely inhibit the amidolytic activity of the BAL fluids suggests as well that multiple proteases are probably present.

The inability to detect active neutrophil elastase does not preclude its potential importance in mediating tissue injury in the lung. As noted in the previous paragraph, active elastase could be present, but bound to structures in the lung and less amenable to inhibition by α₁-PI. In addition, neutrophils may well release elastase while adherent to connective tissue components of the lung in a position protected from α₁-PI or the ¹²⁵I-PK probe (a relatively large molecule that may not penetrate beneath adherent neutrophils). It should also be noted that in these short term experiments, i.e., <6 h of inflammation as opposed to >24 h in adult respiratory distress syndrome patients as an example, the α₁-PI remains relatively active. It is altogether possible that oxidative inactivation of the α₁-PI in the vicinity of leukocytes is greater than in more distant areas. This concept is supported by the complete oxidative inactivation of neutrophilic MPO seen in the presence of aminotriazole. Elastase at the point of release from leukocytes could therefore be protected from inactivation by α₁-PI by this means as well.

Another way to facilitate the detection of elastase activity would be to block the α₁-PI activity in vivo. This was done in six monkeys in the current studies (unpublished results) following published procedures (16, 53) in which the animals were given 50 mg/kg CT intravenously daily for 7 d before injury with FNLP. On the day preceding and the day of the experiment, two additional CT injections were given since activity of the α₁-PI in the monkey plasma before these final injections of CT was >70% of normal. While levels of α₁-PI activity then did approach 10% of normal, the surprising observation was made that the neutrophils in these monkeys had lost most of their elastase activity. CT could be shown in vitro to have no direct effect on neutrophil elastase. Why CT treatment should affect elastase levels in neutrophils in vivo is unclear.

Three methods were employed to monitor the production of oxidants in vivo. The first, measurement of the specific activity of α₁-PI, is based on the finding in recent years that a variety of biologically produced substances can cause oxidative inactivation

of α_1 -PI (9, 14, 15, 19, 39). Stimulated neutrophils have been shown to release such oxidants (54–57) and MPO, in the presence of H_2O_2 and halide has also been shown to cause inactivation of α_1 -PI (58). Studies showing the direct inactivation of α_1 -PI in the presence of PMA-stimulated human neutrophils and halide have been reported (17). In the studies reported here, significant inactivation of α_1 -PI did occur, with mean specific activities dropping from 0.80 in untreated animals to 0.57 and 0.65 in FNLP- and PMA-treated animals, respectively. The specific activity of α_1 -PI in plasma of treated and untreated monkeys ranged from 0.86 to 0.91, suggesting that inactivation was occurring at the site of inflammation in the lung but not systemically. It is possible for α_1 -PI to become inactivated by means other than oxidation. Thiol proteases can cleave the molecule (59) and complexing with neutrophil elastase results in “inactive” α_1 -PI. While elastase- α_1 -PI complexes can be demonstrated in the present studies, the maximum quantity of α_1 -PI which might be involved is less than 4 percent. That the inactivation of α_1 -PI observed in these studies was, in fact, due to oxidation, was demonstrated by our ability to recover the activity by exposure of the partially inactive BAL samples to the reducing agent dithiothreitol in the presence of MS reductase. Full activity was recovered in four out of five cases (Table V) and partial recovery was observed in the fifth. Contrary to our previous studies using human α_1 -PI (9), partial reactivation was found to occur using DTT alone. This may represent a species difference or perhaps a true difference in the sensitivity of the oxidized α_1 -PI to reactivation. Normal monkey plasma oxidized with N-chlorosuccinimide showed 15% reactivation with DTT, which increased to 55% when the MS reductase was present.

A second method of oxidant detection was the inactivation of catalase (23), and, as shown in this paper, myeloperoxidase, which occurs in animals pretreated with AT when H_2O_2 is produced. In these studies, striking decreases in catalase specific activity were seen when AT-treated animals were injured with either FNLP or PMA (Table III). Because inhibition of catalase in AT-treated animals only occurs when H_2O_2 is present, these diminished specific activities constitute good evidence that oxidant is being produced. The more striking decrease in catalase specific activity in monkeys treated with FNLP + AT as opposed to PMA + AT could be a reflection of (a) the shorter exposure of the PMA-treated monkeys to the stimulus (3 h as opposed to 5–6 h); (b) a greater loss of catalase into the extracellular milieu, presumably as a result of greater cell lysis; or (c) the double exposure of FNLP monkeys to the stimulant, the second exposure being timed to react with leukocytes that enter the lung tissues at 4–5 h after the first exposure. It is interesting to note that even in the absence of AT, FNLP-treated animals showed a drop in catalase specific activity from 0.97 to 0.57. The reasons for this are unclear, but it is possible that the catalase was inactivated by oxidants. It also appeared in these studies that levels of myeloperoxidase were greatly affected by pretreatment of the monkeys with AT (Table III). Although it would be reasonable to assume that the mechanism of inhibition might be similar to that of the catalase- H_2O_2 -AT interaction, this had never been shown for MPO. We therefore conducted in vitro studies in which H_2O_2 was allowed to diffuse slowly into mixtures of MPO and AT and compared the residual MPO activity with that obtained with similar samples in which H_2O_2 was absent. The results (Table VI) clearly show the requirement for H_2O_2 for this inhibition.

The third method of oxidant detection that we used was measurement of intracellular glutathione levels. Participation of the glutathione oxidation-reduction cycle in antioxidant reactions would be reflected in decreased levels of glutathione intracellularly. Such decreases were not seen in these studies (Table VII). In similar studies done in rabbits (13), decreases in glutathione in lavaged alveolar cells were seen, although the decrease was not as dramatic as it was in whole lung tissue.

The establishment of a primate model for the study of pulmonary inflammatory injury is an important adjunct to studies usually performed in lower mammals. The ability to measure the generation in vivo of both proteases and oxidants as they appear during the development of pulmonary injury is essential to further studies and to the successful monitoring of the effects of various potential therapeutic agents.

Acknowledgments

The authors wish to acknowledge the expert assistance of Dr. Blaine Anderson, Dr. Ramarao Shankararao, and Dr. Laksman Sehgal, Ms. Hansa Sehgal, and Ms. Augustine Hunt-Coleman for their expert care and handling of the monkeys used in these studies. We thank also Dr. Manfred Schmitt for his help in the development of the ELISA assays, Dr. Ernest Beutler for his suggestion of a method for slow diffusion of H_2O_2 into a solution, Ms. Zenaida Oades for her help in performing H_2O_2 concentration determinations, Dr. Roger Spragg for his assistance in the statistical analyses, Dr. John Peters for his helpful discussions, and Mrs. Monica Bartlett for preparation of the manuscript.

These studies were supported by U. S. Public Health Service grants A117354 and 2P50-23584, the Council for Tobacco Research (764I), and the Office of Naval Research (105-837).

References

1. Janoff, A., B. Sloan, G. Weinbaum, V. Damiano, R. A. Sandhaus, J. Elias, and P. Kimbel. 1977. Experimental emphysema produced by purified human neutrophil elastase. Tissue localization of the instilled protease. *Am. Rev. Respir. Dis.* 115:461–478.
2. Cochrane, C. G., and S. D. Revak. 1980. The participation of high molecular weight kininogen in hypotensive, shock and intravascular coagulation. *Clin. Immunol. Immunopathol.* 15:367–374.
3. Lee, C. T., A. M. Fein, M. Lippmann, H. Holtzman, P. Kimbel, and G. Weinbaum. 1981. Elastolytic activity in pulmonary lavage fluid from patients with adult respiratory distress syndrome. *N. Engl. J. Med.* 304:192–196.
4. O’Brodivich, H. M., S. A. Stalcup, L. M. Pang, J. S. Lipset, and R. B. Mellins. 1981. Bradykinin production and increased pulmonary endothelial permeability during acute respiratory failure in unanesthetized sheep. *J. Clin. Invest.* 67:514–522.
5. Johnson, K. J., J. C. Fantone III, J. Kaplan, and P. A. Ward. 1981. In vivo damage of rat lungs by oxygen metabolites. *J. Clin. Invest.* 67:983–993.
6. McGuire, W. W., R. G. Spragg, A. B. Cohen, and C. G. Cochrane. 1982. Studies on the pathogenesis of the adult respiratory distress syndrome. *J. Clin. Invest.* 69:543–553.
7. Shasby, D. M., K. M. Vanbenthuyzen, R. M. Tate, S. S. Shasby, I. McMurtry, and J. E. Repine. 1982. Granulocytes mediate acute edematous lung injury in rabbits and in isolated rabbit lungs perfused with phorbol myristate acetate: Role of oxygen radicals. *Am. Rev. Respir. Dis.* 125:443–447.
8. Till, G. O., K. J. Johnson, R. Kunkel, and P. A. Ward. 1982. Intravascular activation of complement and acute lung injury. Dependency on neutrophils and toxic oxygen metabolites. *J. Clin. Invest.* 69:1126–1135.

9. Cochrane, C. G., R. G. Spragg, and S. D. Revak. 1983. Studies on the pathogenesis of the adult respiratory distress syndrome: Evidence of oxidant activity in bronchoalveolar lavage fluid. *J. Clin. Invest.* 71: 754-761.
10. Merritt, T. A., C. G. Cochrane, K. Holcomb, B. Bohl, M. Hallman, D. Strayer, K. K. Edwards III, and L. Gluck. 1983. Elastase and α_1 -proteinase inhibitor activity in tracheal aspirates. Role of inflammation in the pathogenesis of bronchopulmonary dysplasia. *J. Clin. Invest.* 72: 656-666.
11. Ward, P. A., G. O. Till, R. Kunkel, and C. Beauchamp. 1983. Evidence for role of hydroxyl radical in complement and neutrophil-dependent tissue injury. *J. Clin. Invest.* 72:789-801.
12. Fligel, S. E. G., P. A. Ward, K. J. Johnson, G. O. Till. 1984. Evidence for role of hydroxyl radicals in immune complex induced vasculitis. *Am. J. Pathol.* 115:375-382.
13. Schraufstatter, I. U., S. D. Revak, and C. G. Cochrane. 1984. Proteases and oxidants in experimental pulmonary inflammatory injury. *J. Clin. Invest.* 73:1175-1184.
14. Johnson, D., and J. Travis. 1978. Structural evidence for methionine at the reaction site of α -1-proteinase inhibitor. *J. Biol. Chem.* 253:7142-7144.
15. Johnson, D., and J. Travis. 1979. The oxidative inactivation of human α -1-proteinase inhibitor. Further evidence for methionine at the reactive center. *J. Biol. Chem.* 254:4022-4026.
16. Cohen, A. B. 1979. The effects *in vivo* and *in vitro* of oxidative damage to purified α_1 -antitrypsin and to the enzyme-inhibiting activity of plasma. *Am. Rev. Resp. Dis.* 119:953-960.
17. Zaslów, M. C., R. A. Clark, P. J. Stone, J. D. Calor, G. L. Snider, and C. Franzblau. 1983. Human neutrophil elastase does not bind to alpha₁-protease inhibitor that has been exposed to activated human neutrophils. *Am. Rev. Resp. Dis.* 128:434-439.
18. Carp, H., and A. Janoff. 1978. Possible mechanisms of emphysema in smokers. *Am. Rev. of Resp. Dis.* 118:617-621.
19. Carp, H., F. Miller, J. R. Hoidal, and A. Janoff. 1982. Potential mechanism of emphysema: α 1-proteinase inhibitor recovered from lungs of cigarette smokers contains oxidized methionine and has decreased elastase inhibitory capacity. *Proc. Natl. Acad. Sci. USA.* 79:2041-2045.
20. Hein, W. G., D. Appleman, and H. T. Pyfrom. 1956. Effects of 3-amino-1,2,4-triazole (AT) on catalase and other components. *Am. J. Physiol.* 186:19-26.
21. Margoliash, E., and A. Novogrodsky. 1958. A study of the inhibition of catalase by 3-amino-1:2:4-triazole. *Biochem. J.* 68:468-475.
22. Margoliash, E., A. Novogrodsky, and A. Schejter. 1960. Irreversible reaction of 3-amino-1:2:4 triazole and related inhibitors with the protein of catalase. *Biochem. J.* 74:339-348.
23. Sinet, P. M., R. E. Heikkela, and G. Cohen. 1980. Hydrogen peroxide production by rat brain *in vivo*. *J. Neurochem.* 34:1421-1428.
24. McConahey, P. J., and F. J. Dixon. 1966. A method for trace iodination of proteins for immunologic studies. *Int. Arch. Allergy Appl. Immunol.* 29:185-193.
25. Stansell, M. J., and H. F. Deutsch. 1965. Preparation of crystalline erythrocyte and catalase from human erythrocytes. *J. Biol. Chem.* 240:4299-4305.
26. Showell, H. J., R. J. Freer, S. H. Zigmond, E. Schiffmann, S. Aswanikumar, B. Corcoran, and E. L. Becker. 1976. The structure-activity relations of synthetic peptides as chemotactic factors and inducers of lysosomal enzyme secretion for neutrophils. *J. Exp. Med.* 143:1154-1169.
27. O'Flaherty, J. T., H. J. Showell, E. L. Becker, and P. A. Ward. 1978. Substances which aggregate neutrophils. *Am. J. Pathol.* 92:155-166.
28. Zigmond, S. H. 1978. Chemotaxis by polymorphonuclear leukocytes. *J. Cell. Biol.* 77:269-287.
29. Freer, R. J., A. R. Day, J. A. Radding, E. Schiffmann, S. Aswanikumar, H. J. Showell, and E. L. Becker. 1980. Further studies on the structural requirements for synthetic peptide chemoattractants. *Biochemistry.* 19:2404-2410.
30. O'Flaherty, J. T., S. Cousart, A. S. Lineberger, E. Bond, D. A. Bass, D. Phil, L. R. De Chatelet, E. S. Leake, and C. G. McCall. 1980. Phorbol myristate acetate. *In vivo* effects upon neutrophils, platelets, and lung. *Am. J. Pathol.* 101:79-92.
31. Weissmann, G., J. E. Smolen, and H. M. Korchak. 1980. Release of inflammatory mediators from stimulated neutrophils. *N. Engl. J. Med.* 303:27-34.
32. Johnson, K. J., and P. A. Ward. 1982. Acute and progressive lung injury after contact with phorbol myristate acetate. *Am. J. Pathol.* 107:29-35.
33. Shasby, D. M., K. M. Vanbenthuyzen, R. M. Tate, S. S. Shasby, I. McMurtry, and J. E. Repine. 1982. Granulocytes mediate acute edematous lung injury in rabbits and in isolated rabbit lungs perfused with phorbol myristate acetate: role of oxygen radicals. *Am. Rev. Resp. Dis.* 125:443-447.
34. Lowry, O. H., N. J. Rosebrough, A. L. Farr, and R. J. Randall. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* 193:265-275.
35. Aebi, H. 1974. Catalase. *In Methods of Enzymatic Analysis.* H. U. Bergmeyer, editor. Vol. 2. Academic Press, Inc., New York, 673-684.
36. Fishman, W. H. 1965. *In Methods of Enzymatic Analysis.* H. U. Bergmeyer, editor. Academic Press, Inc., New York, 869-874.
37. Hojima, Y., J. V. Pierce, and J. J. Pisano. 1980. Hageman factor fragment inhibitor in corn seeds: purification and characterization. *Thromb. Res.* 20:149-162.
38. Mancini, G. A., A. O. Carbonas, and J. F. Heremans. 1965. Immunological quantitation of antigen by single radial immunodiffusion. *Immunochemistry.* 2:235-254.
39. Pannell, R., D. Johnson, and J. Travis. 1974. Isolation and properties of human plasma α -1-proteinase inhibitor. *Biochemistry.* 13:5439-5445.
40. Abrams, W. R., G. Weinbaum, L. Weissbach, H. Weissbach, and N. Brot. 1981. Enzymatic reduction of oxidized α -1-proteinase inhibitor restores biological activity. *Proc. Natl. Acad. Sci. USA.* 78:7483-7486.
41. Brehe, J. E., and H. B. Burch. 1976. Enzymatic assay for glutathione. *Anal. Biochem.* 74:189-197.
42. Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T₄. *Nature (Lond.)* 227:680-685.
43. Clancy, R. M., and T. E. Hugli. 1983. The extraction of leukotrienes (LTC₄, LTD₄ and LTE₄) from tissue fluids: the metabolism of these mediators during IgE-dependent hypersensitivity reactions in lung. *Anal. Biochem.* 133:30-39.
44. Friedlund, J., and E. Silverstein. 1976. A sensitive fluorimetric assay for angiotensin-converting enzyme. *Am. J. Clin. Pathol.* 66:416-424.
45. Cohen, G., and P. Hochstein. 1961. Glucose-6-phosphate dehydrogenase and detoxification of hydrogen peroxide in human erythrocytes. *Science (Wash. DC)* 134:1756-1757.
46. Hyslop, P. A., and L. A. Sklar. 1984. A quantitative fluorimetric assay for the determination of oxidant production by polymorphonuclear leukocytes: its use in the simultaneous fluorimetric assay of cellular activation processes. *Anal. Biochem.* 141:280-286.
47. Kazmierowski, J. A., J. I. Gallin, and H. Y. Reynolds. 1977. Mechanism for the inflammatory response in primate lungs. Demonstration and partial characterization of an alveolar macrophage-derived chemotactic factor with preferential activity for polymorphonuclear leukocytes. *J. Clin. Invest.* 59:273-281.
48. Damiano, V. V., A. Cohen, A. L. Tsang, G. Batra, and R. Petersen. 1980. A morphologic study of the influx of neutrophils into dog lung alveoli after lavage with sterile saline. *Am. J. Pathol.* 100:349-354.
49. Kaplan, A. P., and K. F. Austen. 1970. A prealbumin activator of prekallikrein. *J. Immunol.* 105:802-811.
50. Wuepper, K. D., and C. G. Cochrane. 1972. Plasma prekallikrein: Isolation, characterization, and mechanism of activation. *J. Exp. Med.* 135:1-20.

51. Bouma, B. N., L. A. Miles, G. Beretta, and J. H. Griffin. 1980. Human plasma prekallikrein. Studies of its activation by activated Factor XII and of its inactivation by diisopropyl phosphofluoridate. *Biochemistry*. 19:1151-1160.
52. Meier, H. L., A. P. Kaplan, L. M. Lichtenstein, S. Revak, C. G. Cochrane, and H. H. Newball. 1983. Anaphylactic release of a prekallikrein activator from human lung in vitro. *J. Clin. Invest.* 72:574-581.
53. Abrams, W. R., A. B. Cohen, V. V. Damiano, A. E. Eliraz, P. Kimbel, D. R. Meranze, and G. Weinbaum. 1981. A model of decreased functional α -1-proteinase inhibitor. *J. Clin. Invest.* 68:1132-1139.
54. Babior, B. M., R. Kipnes, and J. Curmutte. 1973. Biological defense mechanisms. The production by leukocytes of superoxide, a potential bactericidal agent. *J. Clin. Invest.* 52:741-744.
55. Goldstein, I. M., D. Roos, H. B. Kaplan, and G. Weissmann. 1975. Complement and immunoglobulins stimulate superoxide production by human leukocytes independently of phagocytosis. *J. Clin. Invest.* 56:1155-1163.
56. Klebanoff, S. J., and H. Rosen. 1978. Ethylene formation by polymorphonuclear leukocytes. Role of myeloperoxidase. *J. Exp. Med.* 148:490-506.
57. Carp, H., and A. Janoff. 1979. In vitro suppression of serum elastase-inhibitory capacity by reactive oxygen species generated by phagocytosing polymorphonuclear leukocytes. *J. Clin. Invest.* 63:793-797.
58. Matheson, N. R., P. S. Wong, and J. Travis. 1979. Enzymatic inactivation of alpha-1-proteinase inhibitor by neutrophil myeloperoxidase. *Biochem. Biophys. Res. Commun.* 88:402-409.
59. Johnson, D., and J. Travis. 1977. Inactivation of human alpha-1-proteinase inhibitor by thiol proteinases. *Biochem. J.* 163:639-641.