

## **SUPPLEMENTAL MATERIAL**

**Supplemental Table 1.** CMR measurements of study cohorts at 3 month follow-up examination.

All Subjects (n=65)	No Prior Chemo (n=42)	Prior Chemo (n=23)	No Prior vs Prior Chemo: p-value
<b><i>CMR Measurements at 3 Month Follow-up</i></b>			
Heart Rate, bpm	77±15	77±14	75±17
LVEDV, ml	113±32	115±31	112±33
LVEDV indexed, ml/m <sup>2</sup>	59±16	58±17	61±13
LVESV, ml	51±18	50±16	54±21
LVESV indexed, ml/m <sup>2</sup>	27±9	25±9	29±9
Stroke Volume, ml	62±18	65±19	57±15
Stroke Volume indexed, ml/m <sup>2</sup>	33±8	34±8	32±7
T2w Relative Enhancement	1.4±0.3	1.4±0.3	1.3±0.3
Myocardial Edema, %	5.6±11	7.0±13	3.2±6.9
LGE-SI	15.9±6.8	15.6±6.8	16.5±7.0
LVEF, %	54±7	56±6	52±7
Values reported as mean ± standard deviation. <i>p</i> <0.05 considered statistically significant (*).			
<i>p</i> -values reported in last column from t-test of differences in means for continuous variables and Fisher's exact / chi-square tests for categorical differences between subgroups of prior chemotherapy.			

