

Supplementary table 2. Genotype data on Hardy-Weinberg equilibrium and minor allele frequency on the combined cohort.

| rs-nr | Position | Observed | Predicted | Hardy-Weinberg | Successful | Minor allele | Major:Minor |
|------------|-----------|----------------|----------------|---------------------|-------------|--------------|---------------------|
| | | heterozygosity | heterozygosity | equilibrium p value | genotyped % | frequency | allele [#] |
| rs2808632 | 159687373 | 0.472 | 0.434 | 0.163 | 100 | 0.318 | T:G |
| rs1800947 | 159683438 | 0.117 | 0.110 | 0.653 | 100 | 0.059 | C:G |
| rs1130864 | 159683091 | 0.421 | 0.424 | 0.966 | 100 | 0.305 | G:A |
| rs1205 | 159682233 | 0.446 | 0.434 | 0.751 | 99.4 | 0.318 | C:T |
| rs3093077 | 159679636 | 0.098 | 0.094 | 0.921 | 99.7 | 0.049 | A:C |
| rs876538 | 159675717 | 0.376 | 0.350 | 0.260 | 99.4 | 0.226 | C:T |
| rs11265257 | 159668984 | 0.472 | 0.476 | 0.932 | 100 | 0.391 | C:T |

#Nucleotides are given according to the plus strand.