

Table 9: Comparison of patients who had SD-OCT and those with TD-OCT at 2 years (N=1023)

	Patients with SD-OCT for ORT evaluation (n=384)	Patients with TD-OCT only (n=639)	P value*
Baseline Characteristics			
Age, Mean (SE)	78.6 (0.39)	79.0 (0.30)	0.40
VA in Study eye, Mean (SE)	61.4 (0.68)	60.8 (0.53)	0.45
total lesion size, Mean (SE)	2.5 (0.13)	2.4 (0.10)	0.61
GA in Study eye, n(%)	30 (7.8%)	39 (6.1%)	0.31
Diabetes, n(%)	67 (17.4%)	108 (16.9%)	0.86
blocked fluorescence (Lesion component) , n(%)	49 (12.8%)	93 (14.6%)	0.46
SHRM, n(%)	281 (74.5%)	492 (77.7%)	0.25
2-Year Outcomes			
VA in Study eye, Mean (SE)	67.2 (0.93)	67.7 (0.72)	0.65
Total lesion size, Mean (SE)	3.3 (0.16)	3.2 (0.13)	0.61
GA in study eye, n(%)	77 (20.1%)	139 (22.2%)	0.48
blocked fluorescence (pathology in foveal center) , n(%)	9 (2.3%)	18 (2.8%)	0.69
SHRM, n(%)	175 (45.8%)	293 (47.0%)	0.74

*P values are two group ttest for comparison of means, from Fisher's exact test for comparison of proportions.