

SUPPLEMENTARY TABLE S1. MOBILIZATION EFFECT ON SPLEEN SIZE, SPLENIC INFARCTS, WHITE BLOOD CELL COUNT, AND RETICULOCYTES

	<i>Spleen ratio</i>	<i>Frequency of splenic infarcts</i>	<i>White blood cells</i>		<i>Reticulocytes (%)</i>	
			<i>Nonsplenectomized</i>	<i>Splenectomized</i>	<i>Nonsplenectomized</i>	<i>Splenectomized</i>
Untreated	21.6 ± 1.00	—	3860 ± 1189	12,947 ± 4426	24.61 ± 1.74	15.82 ± 1.07 ^d
G-CSF	30.44 ± 1.13 ^a	8.82%	23,943 ± 2714 ^a	39,842 ± 7716 ^{ac}	27.11 ± 3.56	11.42 ± 2.00 ^d
Plerixafor	25.37 ± 1.18	3.57%	15,200 ± 2834 ^b	19,138 ± 5862 ^c	24.14 ± 1.84	13.65 ± 1.37 ^d
G-CSF+ plerixafor	30.54 ± 1.51 ^a	5.56%	20,225 ± 2849 ^a	86,044 ± 16,890 ^a	21.83 ± 1.50	12.74 ± 2.09 ^d

^a*p* < 0.05 vs. untreated, ^b*p* = 0.08 vs. untreated, ^c*p* < 0.05 vs. G-CSF+ plerixafor, ^d*p* < 0.05 vs. nonsplenectomized.