

ONLINE SUPPLEMENT.**Table A.** Hospital PCI Appropriateness for Non-Acute Indications by Hospital Quartile of Asymptomatic Patients at Angiography among Hospitals Reporting at Twice as Many Elective Angiography than Elective PCI

Hospital Characteristics	All Hospitals (n=375)	Hospital Quartile of Asymptomatic Patients				P value
		1 – Lowest (n=93)	2 (n=94)	3 (n=94)	4 – Highest (n=94)	
Median proportion of asymptomatic, % (IQR)	25.1 (16.2, 36.2)	12.4 (9.1, 13.7)	20.6 (18.1, 22.3)	30.7 (27.7, 33.4)	43.0 (39.1, 50.0)	
Range of asymptomatic, %	1.6-73.6	1.6-16.0	16.2-25.1	25.1-36.0	36.2-73.6	
PCI Appropriateness, median (IQR) %						
Inappropriate	22.7 (16.8, 30.5)	15.2 (10.0, 21.1)	21.2 (17.0, 27.1)	24.8 (19.8, 29.9)	29.8 (22.7, 35.7)	<.001
Uncertain	41.4 (34.5, 48.5)	45.4 (35.3, 52.8)	45.5 (38.5, 50.7)	40.2 (34.3, 47.0)	37.6 (32.6, 43.7)	<.001
Appropriate	33.1 (23.9, 42.1)	38.4 (25.8, 50.7)	31.4 (23.5, 41.2)	31.9 (25.0, 40.1)	32.9 (22.6, 39.2)	<.01

Abbreviations: IQR, interquartile range