

ONLINE SUPPLEMENT

A risk factor-based predictive model of outcomes in carotid endarterectomy: the National Surgical Quality Improvement

Program 2005-2010

Kimon Bekelis, M.D.^{1,*}, Samuel F. Bakhoum, Ph.D.^{2,*}, Atman Desai, M.D.¹, Todd A. MacKenzey, Ph.D.³, Philip Goodney, M.D.⁴,

Nicos Labropoulos, Ph.D., D.I.C.⁵

¹Section of Neurosurgery, Dartmouth-Hitchcock Medical Center, Lebanon, NH

²Geisel School of Medicine at Dartmouth, Hanover, NH

³Department of Medicine, Dartmouth-Hitchcock Medical Center, Lebanon, NH

⁴Section of Vascular Surgery, Dartmouth-Hitchcock Medical Center, Lebanon, NH

⁵Section of Vascular Surgery, SUNY Stony Brook, Stony Brook, NY

*These authors contributed equally to this work and co-primary authors

Running Head: NSQIP: predictive model of CEA outcomes

Subject codes: [44], [49], [76]

Key words: carotid stenosis, carotid endarterectomy, risk prediction, NSQIP

Figures: 3, **Tables:** 2, **Supplemental material:** 1

Word count: 4,500

Corresponding Author:

Kimon Bekelis, M.D.

Dartmouth-Hitchcock Medical Center

One Medical Center Dr.

Lebanon, NH 03756

kbekelis@gmail.com

T: 603-650-5110

F: 603-650-4547

Supplemental Figures

Figure 1. Diagrams of receiver operating characteristic (ROC) curves demonstrating the sensitivity against the 1-specificity for all the predictive models for the 30-day postoperative risk of stroke, MI, death, or their combination (*from left to right*).







Figure 2. Multivariate analysis of the 30-day risk of stroke (A), myocardial infarction (B), death (C), or their combination after carotid endarterectomy (D). The odds ratios are represented with forest plots on the right side of each table.




A



B



C



D