

Original Article: Association of urine α 1-microglobulin with kidney function decline and mortality in HIV-infected women

Authors: Vasantha Jotwani^{1,2}, Rebecca Scherzer^{1,2}, Alison Abraham³, Michelle M Estrella⁴, Michael Bennett⁵, Mardge H Cohen⁶, Marek Nowicki⁷, Anjali Sharma⁸, Mary Young⁹, Phyllis C Tien¹, Joachim H Ix^{10, 11}, Mark J Sarnak¹², Chirag R Parikh^{13,14}, Michael G Shlipak^{1,2}

¹Department of Medicine, San Francisco VA Medical Center, San Francisco, USA

²Department of Epidemiology and Biostatistics, University of California, San Francisco, USA

³Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore, USA

⁴Department of Medicine, Johns Hopkins School of Medicine, Baltimore, USA

⁵Division of Nephrology and Hypertension, Cincinnati Children's Hospital Medical Center, Cincinnati, USA

⁶Departments of Medicine, Stroger Hospital and Rush University, Chicago, USA

⁷Department of Medicine, University of Southern California, Los Angeles, USA

⁸Division of Infectious Diseases, Department of Medicine, SUNY Downstate Medical Center, Brooklyn, USA

⁹Division of Infectious Diseases and Travel Medicine, Department of Medicine, Georgetown University Medical Center, Washington, D.C., USA

¹⁰Division of Nephrology-Hypertension, Department of Medicine, University of California, San Diego, USA

¹¹Nephrology Section, Veterans Affairs San Diego Healthcare System, USA

¹²Division of Nephrology, Tufts Medical Center, Boston, USA

¹³Section of Nephrology, Department of Medicine, Yale University, New Haven, USA

¹⁴Program of Applied Translational Research, Yale University, New Haven, USA