

Adjudin protects rodent cochlear hair cells against gentamicin ototoxicity via SIRT3-ROS pathway

Yizhou Quan^{a,b#}, Li Xia^{c#}, Jiaxiang Shao^{a,b}, Shankai Yin^c, C. Yan Cheng^d, Weiliang Xia^{a,b*}, Wei-Qiang Gao^{a,b*}

^aState Key Laboratory of Oncogenes and Related Genes, Renji-MedX Stem Cell Research Center, Ren Ji Hospital, School of Medicine, Shanghai Jiao Tong University, Shanghai, China

^bSchool of Biomedical Engineering & Med-X Research Institute, Shanghai Jiao Tong University, Shanghai, China

^cDepartment of Otolaryngology, Affiliated Sixth People's Hospital of Shanghai Jiao Tong University, Otolaryngology Institute of Shanghai Jiao Tong University, Shanghai, China

^dThe Mary M. Wohlford Laboratory for Male Contraceptive Research, Center for Biomedical Research, Population Council, New York, New York, USA

[#]The authors contributed equally to this work

^{*}Corresponding authors:

Weiliang Xia, Ph.D., Room 211, Med-X Research Institute, Shanghai Jiao Tong University, 1954 Huashan Road, Shanghai 200030, China. Tel: +86-21-62933291; Fax: +86-21-62932302; Email: wxia@sjtu.edu.cn

Or Wei-Qiang Gao, Ph.D., Stem Cell Research Center, RenJiHospital, Shanghai Jiao Tong University School of Medicine, 1630 Dongfang Road, Shanghai 200127, China. Tel: +86-21-62932049; Email: weiqgao@yahoo.com




Fig.S1 Whole mount cochlear tissues infected with GFP or RFP -lenti-virus. Bar = 100 μ m.




Fig.S2 Quantification of outer hair cell loss in the group that received gentamicin (40 mg/kg) or adjudin (75 mg/kg) before gentamicin (40 mg/kg) application and was observed 7 days post the injection. $p > 0.05$, two-way ANOVA of the parameters between various regimens, followed by post-hoc tests.