

Supplemental Table 1. Demographic and clinical characteristics of all patients with a surveillance record available, patients with a surveillance record who seroconverted to a SFGR, and patients with a surveillance record who did not seroconvert.

	All patients (n=53)	Seroconverters (n=12)	Non-seroconverters (n=41)
DEMOGRAPHIC INFORMATION			
Age – median (range)	50 (1 – 80)	49 (1 – 70)	51 (7 – 80)
Gender			
Male	29/47 (61.7%)	7/10 (70.0%)	22/37 (59.5%)
Female	18/47 (38.3%)	3/10 (30.0%)	15/37 (40.5%)
Race			
White	45/53 (84.9%)	10/12 (83.3%)	35/41 (85.4%)
Black	3/53 (5.7%)	0/12 (0%)	3/41 (7.3%)
Other	2/53 (3.8%)	2/12 (16.7%)	0/41 (0%)
Hispanic Ethnicity	3/42 (7.1%)	1/11 (9.1%)	2/31 (6.4%)
CLINICAL INFORMATION			
Symptoms/clinical findings			
Fever	27/46 (58.7%)	8/11 (72.7%)	19/35 (54.3%)
Headache	14/36 (38.9%)	4/10 (40.0%)	10/26 (38.5%)
Myalgias	24/40 (60.0%)	6/10 (60.0%)	18/30 (60.0%)
Skin rash	16/41 (39.0%)	3/11 (27.3%)	13/30 (43.3%)
Acute renal failure	2/29 (6.9%)	1/5 (20.0%)	1/24 (4.2%)
Thrombocytopenia	7/28 (25.0%)	3/6 (50.0%)	4/22 (18.2%)
Leukopenia	5/26 (19.2%)	3/5 (60.0%)	2/21 (9.5%)
Anemia	4/28 (14.3%)	2/6 (33.3%)	2/22 (9.1%)
Elevated liver enzymes	8/28 (28.6%)	2/5 (40.0%)	6/23 (26.1%)
Took antibiotics for illness	36/45 (80.0%)	9/9 (100%)	27/36 (75.0%)
Patient hospitalized	7/48 (14.6%)	2/12 (16.7%)	5/36 (13.9%)
Died from illness	0/20 (0%)	0/5 (0%)	0/15 (0%)