

Supplemental Table for:
Prognostic Factors of Survival in a Randomized Phase 3 Trial (MPACT) of Weekly nab-Paclitaxel Plus Gemcitabine vs
Gemcitabine Alone in Patients With Metastatic Pancreatic Cancer
Josep Tabernero et al.

Table S1. Selected baseline characteristics.

Variable	<i>nab-P + Gem</i> (n = 431)	Gem (n = 430)	All Patients (N = 861)	
Age	Median years (min, max) ≥ 65 years, n (%)	62 (27, 86) 177 (41)	63 (32, 88) 188 (44)	63 (27, 88) 365 (42)
Sex, n (%)	Male	245 (57)	257 (60)	502 (58)
Region, n (%)	North America Australia Eastern Europe Western Europe	268 (62) 61 (14) 64 (15) 38 (9)	271 (63) 59 (14) 62 (14) 38 (9)	539 (63) 120 (14) 126 (15) 76 (9)
KPS score, n (%)	70 80 90 100 70–80 90–100	30 (7) 149 (35) 179 (42) 69 (16) 179 (42) 248 (58)	33 (8) 128 (30) 199 (46) 69 (16) 161 (38) 268 (62)	63 (7) 277 (32) 378 (44) 138 (16) 340 (40) 516 (60)
Pancreatic primary tumor location, n (%)	Head Body Tail	191 (44) 132 (31) 105 (24)	180 (42) 136 (32) 110 (26)	371 (43) 268 (31) 215 (25)
Number of metastatic sites, n (%)	1 2 ≥ 3	33 (8) 202 (47) 196 (45)	21 (5) 206 (48) 203 (47)	54 (6) 408 (47) 399 (46)
CA19-9, n (%)	Normal > ULN but < 59 × ULN ≥ 59 × ULN	60 (14) 122 (28) 197 (46)	56 (13) 120 (28) 195 (45)	116 (13) 242 (28) 392 (46)

CA19-9, carbohydrate antigen 19-9; Gem, gemcitabine; KPS, Karnofsky performance status; min, minimum; max, maximum; *nab-P*, *nab*-paclitaxel; ULN, upper limit of normal.