

Supplementary figures

Handedness and the X chromosome: The role of androgen receptor

CAG-repeat length

Larissa Arning*⁺¹, Sebastian Ocklenburg⁺², Stefanie Schulz², Vanessa Ness², Wanda M. Gerding¹, Jan G. Hengstler³, Michael Falkenstein³, Jörg T. Epplen¹, Onur Güntürkün², Christian Beste⁴

Affiliations

¹Department of Human Genetics, Ruhr-University, 44780 Bochum, Germany

²Institute of Cognitive Neuroscience, Biopsychology, Department of Psychology, Ruhr-University, 44780 Bochum, Germany

³Leibniz Research Centre for Working Environment and Human Factors (IfADo), 44139 Dortmund, Germany

⁴Cognitive Neurophysiology, Department of Child and Adolescent Psychiatry, Faculty of Medicine of the TU Dresden, 01309 Dresden, Germany

Figure Captions for supplementary figures

Supplementary figure S1: LQ distribution for female and male participants.

Supplementary figure S2: Handedness Direction 1 phenotypes (left-handed, right-handed) in relation to CAG-repeat length for male (A) and female participants (B: short allele; C: long allele). Error bars show standard error.

Supplementary figure S3: Handedness Consistency phenotypes (consistent, inconsistent) in relation to CAG-repeat length for male (A) and female participants (B: short allele; C: long allele). Error bars show standard error.

Figure Captions for supplementary figures





