

Supplemental Table 1.

Mean National Eye Institute Visual Functioning Questionnaire-25 subscale scores for patients with strabismus, diabetic retinopathy, age-related macular degeneration (AMD), glaucoma, cataract, cytomegalovirus (CMV) retinitis, and low vision. Strabismic patients were recruited for the Quality of Life in Adults with Strabismus study, and other patients were enrolled in the National Eye Institute-Visual Functioning Questionnaire field test group. Dark gray shading indicates subscale scores that were significantly better than patients with strabismus; light gray shading denotes subscale scores that were not significantly different from strabismic patients; and white shading indicates subscale scores that were significantly worse than patients with strabismus. The threshold for significance is set to 0.05.

	Strabismus (n = 41)	Diabetic retinopathy (n = 123)	AMD (n = 108)	Glaucoma (n = 77)	Cataract (n = 93)	CMV retinitis (n = 37)	Low vision (n = 90)
General health	68 ± 22	46 ± 25	65 ± 25	62 ± 25	55 ± 25	45 ± 24	57 ± 27
General vision	59 ± 20	62 ± 21	53 ± 20	71 ± 17	60 ± 17	76 ± 14	38 ± 18
Ocular pain	65 ± 26	88 ± 17	87 ± 17	89 ± 14	86 ± 19	90 ± 16	85 ± 20
Near activities	61 ± 20	63 ± 30	54 ± 27	79 ± 23	73 ± 21	84 ± 20	36 ± 23
Distance activities	67 ± 21	66 ± 30	56 ± 29	77 ± 25	73 ± 22	84 ± 18	38 ± 26
VS* social functioning	81 ± 22	81 ± 26	73 ± 29	89 ± 20	87 ± 19	96 ± 9	50 ± 31
VS* mental health	53 ± 28	66 ± 29	58 ± 27	81 ± 20	77 ± 22	74 ± 22	46 ± 27
VS* role difficulties	56 ± 29	69 ± 31	61 ± 31	84 ± 23	76 ± 22	78 ± 24	44 ± 29
VS* dependency	82 ± 22	77 ± 30	72 ± 30	92 ± 19	88 ± 20	89 ± 12	51 ± 31
Driving	67 ± 23	55 ± 40	39 ± 36	75 ± 28	63 ± 30	80 ± 28	10 ± 23
Color vision	94 ± 13	90 ± 22	85 ± 25	93 ± 17	90 ± 20	98 ± 9	71 ± 31
Peripheral vision	68 ± 25	78 ± 29	77 ± 27	76 ± 27	87 ± 21	78 ± 21	59 ± 32

\*Vision specific