


Residues in most favoured regions [A,B,L]	273	93.8%
Residues in additional allowed regions [a,b,l,p]	18	6.2%
Residues in generously allowed regions [-a,-b,-l,-p]	0	0.0%
Residues in disallowed regions	0	0.0%

Number of non-glycine and non-proline residues	291	100.0%
Number of end-residues (excl. Gly and Pro)	2	
Number of glycine residues (shown as triangles)	20	
Number of proline residues	13	

Total number of residues	326	

Wild kir6.2 model (PM0079770)



Residues in most favoured regions [A,B,L]	267	92.1%
Residues in additional allowed regions [a,b,l,p]	19	6.6%
Residues in generously allowed regions [-a,-b,-l,-p]	1	0.3%
Residues in disallowed regions	3	1.0%

Number of non-glycine and non-proline residues	290	100.0%
Number of end-residues (excl. Gly and Pro)	2	
Number of glycine residues (shown as triangles)	20	
Number of proline residues	13	

Total number of residues	325	

Protein Model Portal_kir6.2_third model

Supplementary Fig. 5. The backbone folding patterns plotted for current kir6.2 model (A) and third existing model of kir6.2 (B). Each black dot spot corresponds to one amino acid residue of kir6.2 chain. On the plot most favoured regions, additional allowed regions, generously allowed regions and disallowed regions are shown in red, yellow, light yellow and white color respectively.