

Appendix B: Functioning family planning (FP) service delivery points, by country and by FP method

Table B1. Burkina Faso: facilities with essential components to provide short- and long-acting FP methods (n=28)

	Hospital (n=3)	Camp health center (n=4)	Non-camp health centers (n=21)
ORAL CONTRACEPTIVE PILLS			
OCPs provided in last 3 months (self-reported)	3 (100%)	4 (100%)	20 (95.2%)
Staff trained to provide short-acting FP methods	3 (100%)	3 (100%) ND* (1)	20 (95.2%)
FP counseling available	3 (100%)	4 (100%)	20 (95.2%)
BP cuff	3 (100%)	4 (100%)	19 (90.5%)
Stethoscope	3 (100%)	4 (100%)	21 (100%)
Daily oral contraceptive pills	3 (100%)	4 (100%)	20 (95.2%)
Functioning OCP service delivery point	3 (100%)	3 (100%) ND* (1)	17 (81%)
INJECTABLES			
Injectables provided in last 3 months (self-reported)	3 (100%)	4 (100%)	20 (95.2%)
Staff trained to provide short-acting FP methods	3 (100%)	3 (100%) ND* (1)	20 (95.2%)
FP counseling available	3 (100%)	4 (100%)	20 (95.2%)
BP cuff	3 (100%)	4 (100%)	19 (90.5%)
Stethoscope	3 (100%)	4 (100%)	21 (100%)
Needles and syringes	3 (100%)	4 (100%)	21 (100%)
Injectable contraceptive (progestin-only)	3 (100%)	4 (100%)	20 (95.2%)
Functioning injectable service delivery point	3 (100%)	3 (100%) ND* (1)	17 (81%)
IUD			
IUD inserted in last 3 months (self-reported)	3 (100%)	1 (25%)	4 (19%)
Staff trained to provide long-acting methods	3 (100%)	3 (100%) ND* (1)	17 (85%) ND* (1)
FP counseling available	3 (100%)	4 (100%)	20 (95.2%)
Sterile gloves	3 (100%)	4 (100%)	21 (100%)
Speculum	3 (100%)	4 (100%)	20 (95.2%)
Uterine sound	3 (100%)	3 (75%)	5 (23.8%)
Uterine tenaculum	3 (100%)	1 (25%)	6 (30%) ND* (1)
Sponge forceps	3 (100%)	2 (50%)	13 (61.9%)
Antiseptics	3 (100%)	4 (100%)	21 (100%)
IUD (copper or levonorgestrel)	3 (100%)	1 (25%)	10 (47.6%)
Functioning IUD service delivery point	3 (100%)	1 (25%)	1 (4.8%)
IMPLANT			
Implant inserted in last 3 months (self-reported)	3 (100%)	4 (100%)	19 (90.5%)
Staff trained to provide long-acting methods	3 (100%)	3 (100%) ND* (1)	17 (85%) ND* (1)
FP counseling available	3 (100%)	4 (100%)	20 (95.2%)
Sponge forceps	3 (100%)	2 (50%)	13 (61.9%)
Scalpel handle (No. 3) and blade	3 (100%)	3 (75%)	16 (76.2%)
Needles and syringes	3 (100%)	4 (100%)	21 (100%)
Antiseptics	3 (100%)	4 (100%)	21 (100%)
Implant	3 (100%)	4 (100%)	20 (95.2%)
Functioning implant service delivery point	3 (100%)	1 (25%)	8 (40%) ND* (1)

*No data

Table B2. DRC: facilities with essential components to provide short- and long-acting FP methods (n=26)

	Hospital (n=1)	Health center (n=25)
ORAL CONTRACEPTIVE PILLS		
OCPs provided in last 3 months (self-reported)	1	18 (72%)
Staff trained to provide short-acting FP methods	1	16 (66.7%) ND* (1)
FP counseling available	1	19 (76%)
BP cuff	1	20 (83.3%) ND* (1)
Stethoscope	1	22 (91.7%) ND* (1)
Daily oral contraceptive pills	1	15 (62.5%) ND* (1)
Functioning OCP service delivery point	1	12 (48%)
INJECTABLES		
Injectables provided in last 3 months (self-reported)	1	18 (72%) ND* (1)
Staff trained to provide short-acting FP methods	1	16 (66.7%) ND* (1)
FP counseling available	1	19 (76%)
BP cuff	1	20 (83.3%) ND* (1)
Stethoscope	1	22 (91.7%) ND* (1)
Needles and syringes	1	23 (95.8%) ND* (1)
Injectable contraceptive (progestin-only)	1	13 (54.2%) ND* (1)
Functioning injectable service delivery point	1	10 (40%)
IUD		
IUD inserted in last 3 months (self-reported)	1	12 (48%)
Staff trained to provide long-acting methods	1	15 (62.5%) ND* (1)
FP counseling available	1	19 (76%)
Sterile gloves	1	19 (79.2%) ND* (1)
Speculum	1	14 (58.3%) ND* (1)
Uterine sound	1	10 (41.7%) ND* (1)
Uterine tenaculum	1	11 (45.8%) ND* (1)
Sponge forceps	1	11 (45.8%) ND* (1)
Antiseptics	1	23 (95.8%) ND* (1)
IUD (copper or levonorgestrel)	1	11 (45.8%) ND* (1)
Functioning IUD service delivery point	1	9 (36%)
IMPLANT		
Implant inserted in last 3 months (self-reported)	1	15 (60%)
Staff trained to provide long-acting methods	1	15 (62.5%) ND* (1)
FP counseling available	1	19 (76%)
Sponge forceps	1	11 (45.8%) ND* (1)
Scalpel handle (No. 3) and blade	1	11 (45.8%) ND* (1)
Needles and syringes	1	23 (95.8%) ND* (1)
Antiseptics	1	23 (95.8%) ND* (1)
Implant	1	13 (54.2%) ND* (1)
Functioning implant service delivery point	1	5 (20%)

*No data

Table B3. South Sudan: facilities with essential components to provide short- and long-acting FP methods (n=9)

	Hospital (n=1)	Health center (n=8)
ORAL CONTRACEPTIVE PILLS		
OCPs provided in last 3 months (self-reported)	0	5
Staff trained to provide short-acting FP methods	1	4 ND* (3)
FP counseling available	1	6
BP cuff	1	8
Stethoscope	1	8
Daily oral contraceptive pills	ND*	3 ND* (1)
Functioning OCP service delivery point	0	1
INJECTABLES		
Injectables provided in last 3 months (self-reported)	1	4
Staff trained to provide short-acting FP methods	1	4 ND* (3)
FP counseling available	1	6
BP cuff	1	8
Stethoscope	1	8
Needles and syringes	1	7
Injectable contraceptive (progestin-only)	ND*	3 ND* (1)
Functioning injectable service delivery point	ND	1
IUD		
IUD inserted in last 3 months (self-reported)	0	0
Staff trained to provide long-acting methods	0	3
FP counseling available	1	6
Sterile gloves	1	8
Speculum	1	6
Uterine sound	1	4 ND* (1)
Uterine tenaculum	1	4 ND* (1)
Sponge forceps	1	4 ND* (1)
Antiseptics	1	8
IUD (copper or levonorgestrel)	ND*	0 ND* (1)
Functioning IUD service delivery point	0	0
IMPLANT		
Implant inserted in last 3 months (self-reported)	0	1
Staff trained to provide long-acting methods	0	3
FP counseling available	1	6
Sponge forceps	1	4 ND* (1)
Scalpel handle (No. 3) and blade	0	5 ND* (2)
Needles and syringes	1	7
Antiseptics	1	8
Implant	ND*	2 ND* (1)
Functioning implant service delivery point	0	0

*No data