

Appendix G: Facilities with essential supplies for diagnosis and treatment of sexually transmitted infections (STIs) and prevention of mother-to-child transmission of HIV (PMTCT), by country

Table G1. Burkina Faso: Facilities with essential supplies for diagnosis and treatment of STIs and for PMTCT (n=28)

	Hospital (n=3)	Camp health center (n=4)	Non-camp health center (n=21)
DIAGNOSIS & TREATMENT OF STIs			
Performed syndromic or laboratory diagnosis and treatment of STIs in last 3 months	2 (67%)	4 (100%)	21 (100%)
Gentamycin	3 (100%)	3 (75%)	21 (100%)
Ceftriaxone	3 (100%)	4 (100%)	21 (100%)
Injectable metronidazole	3 (100%)	2 (50%)	21 (100%)
Facilities with essential supplies for diagnosis and treatment of STIs	2 (67%)	1 (25%)	21 (100%)
PROVISION OF PMTCT			
Administered ARVs to HIV+ mothers in maternity in the last 3 months	3 (100%)	1 (33.3%) ND* (1)	21 (100%)
Administered ARVs to newborns born to HIV+ mothers in maternity in the last 3 months	3 (100%)	1 (25%)	21 (100%)
ARVs for the mother	3 (100%)	3 (75%)	19 (90.5%)
ARVs for the infant	3 (100%)	3 (75%)	18 (90%) ND* (1)
Facilities with essential supplies to provide PMTCT	3 (100%)	1 (25%)	18 (90%) ND* (1)

*No data

Table G2. DRC: Facilities with essential supplies for diagnosis and treatment of STIs and for PMTCT (n=26)

	Hospital (n=1)	Health center (n=25)
DIAGNOSIS & TREATMENT OF STIs		
Performed syndromic or laboratory diagnosis and treatment of STIs in last 3 months	1 (100%)	22 (88%)
Gentamycin	1 (100%)	6 (25%) ND* (1)
Ceftriaxone	1 (100%)	3 (13%) ND* (2)
Injectable metronidazole	1 (100%)	3 (13%) ND* (2)
Facilities with essential supplies for diagnosis and treatment of STIs	1 (100%)	2 (9%) ND* (2)
PROVISION OF PMTCT		
Administered ARVs to HIV+ mothers in maternity in the last 3 months	1	0
Administered ARVs to newborns born to HIV+ mothers in maternity in the last 3 months	1	1 (4%)
ARVs for the mother	1	1 (4%) ND* (1)
ARVs for the infant	1	0 ND* (1)
Facilities with essential supplies to provide PMTCT	1	0

*No data

Table G3. South Sudan: Facilities with essential supplies for diagnosis and treatment of STIs and for PMTCT (n=9)

	Hospital (n=1)	Health center (n=8)
DIAGNOSIS & TREATMENT OF STIs		
Performed syndromic or laboratory diagnosis and treatment of STIs in last 3 months	1	7
Gentamycin	0	6 ND* (1)
Ceftriaxone	0	6 ND* (1)
Injectable metronidazole	0	4 ND* (1)
Facilities with essential supplies for diagnosis and treatment of STIs	0	3
PROVISION OF PMTCT		
Administered ARVs to HIV+ mothers in maternity in the last 3 months	0	0
Administered ARVs to newborns born to HIV+ mothers in maternity in the last 3 months	0	0
ARVs for the mother	ND*	0 ND* (1)
ARVs for the infant	ND*	0 ND* (1)
Facilities with essential supplies to provide PMTCT	0	0

*No data