

**Table e-1. Score of the modified Rankin Scale at 90 days in randomized groups by baseline systolic blood pressure**

	Intensive	Guideline
<b>&lt;160 mmHg</b>		
0: No symptoms at all	24 (13%)	14 (8%)
1: No substantive disability despite symptoms	40 (21%)	45 (26%)
2: Slight disability	38 (20%)	28 (16%)
3: Moderate disability requiring some help	29 (16%)	34 (20%)
4: Moderate-severe disability requiring assistance with daily living	29 (16%)	23 (13%)
5: Severe disability, bed-bound and incontinent	10 (5%)	11 (6%)
6: Death by 90 days	17 (9%)	17 (10%)
<b>160-169 mmHg</b>		
0: No symptoms at all	29 (10%)	19 (7%)
1: No substantive disability despite symptoms	62 (21%)	55 (21%)
2: Slight disability	56 (19%)	50 (19%)
3: Moderate disability requiring some help	47 (16%)	43 (16%)
4: Moderate-severe disability requiring assistance with daily living	51 (17%)	53 (20%)
5: Severe disability, bed-bound and incontinent	14 (5%)	18 (7%)
6: Death by 90 days	36 (12%)	25 (10%)
<b>170-179 mmHg</b>		
0: No symptoms at all	16 (6%)	22 (7%)
1: No substantive disability despite symptoms	51 (19%)	61 (20%)
2: Slight disability	56 (21%)	50 (16%)
3: Moderate disability requiring some help	42 (16%)	54 (17%)
4: Moderate-severe disability requiring assistance with daily living	47 (18%)	55 (18%)
5: Severe disability, bed-bound and incontinent	17 (6%)	31 (10%)
6: Death by 90 days	33 (13%)	36 (12%)
<b>180-189 mmHg</b>		
0: No symptoms at all	12 (5%)	23 (9%)
1: No substantive disability despite symptoms	56 (22%)	39 (15%)
2: Slight disability	44 (17%)	44 (17%)
3: Moderate disability requiring some help	46 (18%)	41 (16%)
4: Moderate-severe disability requiring assistance with daily living	47 (19%)	57 (22%)
5: Severe disability, bed-bound and incontinent	18 (7%)	21 (8%)
6: Death by 90 days	29 (12%)	34 (13%)
<b>≥190 mmHg</b>		
0: No symptoms at all	31 (8%)	29 (7%)
1: No substantive disability despite symptoms	83 (22%)	54 (13%)
2: Slight disability	65 (17%)	94 (23%)
3: Moderate disability requiring some help	56 (15%)	62 (15%)
4: Moderate-severe disability requiring assistance with daily living	76 (20%)	80 (20%)
5: Severe disability, bed-bound and incontinent	24 (6%)	32 (8%)
6: Death by 90 days	51 (13%)	58 (14%)
<b>OVERALL</b>		
0: No symptoms at all	112 (8%)	107 (8%)

1: No substantive disability despite symptoms	292 (21%)	254 (18%)
2: Slight disability	259 (19%)	266 (19%)
3: Moderate disability requiring some help	220 (16%)	234 (17%)
4: Moderate-severe disability requiring assistance with daily living	250 (18%)	268 (19%)
5: Severe disability, bed-bound and incontinent	83 (6%)	113 (8%)
6: Death by 90 days	166 (12%)	170 (12%)

---