

Supplementary Table 1 List of primers used for qPCR and their corresponding annealing temperatures. The cycling conditions were: 95°C for 10 min, followed by 40 cycles of 95°C for 15 sec, T_m for 20 sec and 62°C for 15 sec.

GENE		SEQUENCE (5'-3')	Annealing temperature
cyc D1	Forward	CCGTCCATGCGGAAGATC	62°C
	Reverse	ATGGCCAGCGGGAAGAC	
c-Myc	Forward	TCAAGAGGTGCCACGTCTCC	62°C
	Reverse	TCTTGGCAGCAGGATAGTCCTT	
survivin	Forward	AGAACTGGCCCTTCTTGGGGTC	60°C
	Reverse	CTTTTTATGTTCTCTATGGGGTC	
hTERT	Forward	AAGTTCCTGCACTGGCTGAT	60°C
	Reverse	CACGACGTAGTCCATGTTC	
HPV16 E6	Forward	TGCAATGTTTCAGGACCCA	60°C
	Reverse	CATGTATAGTTGTTTGCAGCTCTGT	
IL-6	Forward	TACCCCCAGGAGAAGATTCC	60°C
	Reverse	GCCATCTTTGGAAGGTTTCAG	
IL-8	Forward	AGGTGCAGTTTTGCCAAGGA	60°C
	Reverse	TTTCTGTGTTGGCGCAGTGT	
β-actin	Forward	TCATGAAGATCCTCACCGAG	58°C
	Reverse	TTGCCAATGGTGATGACCTG	