Electrocardiogram (EKG) findings in stage B versus stage C of the study population

| Abnormality* | Stage B | | Stage C | |
|---|---|---|---|---|
| | N | Percent | N | Percent |
| Atrial fibrillation or flutter | 1 | 5.0 | 4 | 14.8 |
| Junctional rhythm | 0 | 0.0 | 3 | 11.1 |
| Multiple pre-mature ventricular beats | 0 | 0.0 | 3 | 11.1 |
| Grade I atrioventricular block | 1 | 5.0 | 2 | 7.4 |
| Grade II atrioventricular block | 0 | 0.0 | 0 | 0.0 |
| 2:1 Atrioventricular block | 0 | 0.0 | 0 | 0.0 |
| Grade III atrioventricular block | 0 | 0.0 | 1 | 3.7 |
| Right bundle branch block | 14 | 70.0 | 12 | 44.4 |
| Incomplete right bundle branch block | 1 | 5.0 | 0 | 0.0 |
| Left bundle branch block | 1 | 5.0 | 0 | 0.0 |
| Left anterior fascicular block | 12 | 60.0 | 14 | 51.9 |
| Left posterior fascicular block | 0 | 0.0 | 1 | 3.7 |
| Bradycardia (< 50 beats/minute) | 0 | 0.0 | 5 | 18.5 |
| Cardiac pacemaker | 0 | 0.0 | 2 | 7.4 |

*Some individuals had more than one abnormality. Individuals in stage A, by definition, had none of these abnormalities.