

**Table S1. *Francisella* Strains Used in This Study, Related to Experimental Procedures**

Strain	Description	Source
Fn	<i>F. tularensis</i> subsp. <i>novicida</i> Utah 112 strain	*
Fn-FLAG-VgrG	Fn expressing FLAG-VgrG; chromosomal <i>vgrG</i> (FTN_1312) replaced by a copy of <i>vgrG</i> fused with the coding sequence for 3xFLAG epitope at its 5' end	This study
Fn-FLAG-VgrG $\Delta iglA$	Fn-FLAG-VgrG with deletion of <i>iglA</i> (FTN_1324)	This study
Fn-IgIA-GFP	Fn expressing IgIA-GFP; chromosomal <i>iglA</i> replaced by a copy of <i>iglA</i> fused with the coding sequence for superfolder GFP at its 3' end	This study
Fn-IgIA-GFP $\Delta iglB$	Fn-IgIA-GFP with deletion of <i>iglB</i> (FTN_1323)	This study
Fn-IgIA-GFP $\Delta iglC$	Fn-IgIA-GFP with deletion of <i>iglC</i> (FTN_1322)	This study
Fn-IgIA/B-split GFP	Fn expressing IgIA/B-split GFP; chromosomal <i>iglA</i> replaced by a copy of <i>iglA</i> fused with the coding sequence for superfolder GFP domains 1-10 (GFP1-10) at its 3' end and chromosomal <i>iglB</i> replaced by a copy of <i>iglB</i> fused with the coding sequence for superfolder GFP domain 11 (GFP11) at its 5' end	This study
Fn-IgIA $\Delta$ 18N/B-split GFP	Fn-IgIA/B-split GFP with deletion of amino acid residues #2-18 of IgIA	This study
Fn-IgIA/B $\Delta$ C25-split GFP	Fn-IgIA/B-split GFP with deletion of the last 25 amino acid residues #482-506 of IgIB	This study
Fn-IgIA/SodB-split GFP	Fn expressing IgIA/SodB-split GFP; chromosomal <i>iglA</i> replaced by a copy of <i>iglA</i> fused with the coding sequence for GFP1-10 at its 3' end and chromosomal <i>sodB</i> (FTN_1642) replaced by a copy of <i>sodB</i> fused with the coding sequence for GFP11 at its 5' end	This study
Fn-IgIA/B-split GFP, FLAG-VgrG	Fn-IgIA/B-split GFP strain expressing FLAG-VgrG; chromosomal <i>vgrG</i> replaced by a copy of <i>vgrG</i> fused with the coding sequence for 3xFLAG epitope at its 5' end	This study
Fn-IgIA $\Delta$ 18N/B-split GFP, FLAG-VgrG	Fn-IgIA $\Delta$ 18N/B-split GFP strain expressing FLAG-VgrG; chromosomal <i>vgrG</i> replaced by a copy of <i>vgrG</i> fused with the coding sequence for 3xFLAG epitope at its 5' end	This study
Fn-IgIA/B $\Delta$ C25-split GFP, FLAG-VgrG	Fn-IgIA/B $\Delta$ C25-split GFP strain expressing FLAG-VgrG; chromosomal <i>vgrG</i> replaced by a copy of <i>vgrG</i> fused with the coding sequence for 3xFLAG epitope at its 5' end	This study

\* Karl Klose, University of Texas San Antonio, San Antonio, TX