

Table S3. Ischemic atherosclerotic events separated into MI, PAD and ischemic stroke: number of events, and event rate by quintiles of baseline urine NGAL concentration (a patient could have more than one event)

| | | Urine NGAL concentration (ng/ml) | | | | |
|-----------------|------------------------------------|----------------------------------|--------------|---------------|---------------|--------|
| | | ≤6.9 | >6.9 - ≤12.9 | >12.9 - ≤22.6 | >22.6 - ≤49.5 | >49.5 |
| MI | No. of events | 31 | 44 | 40 | 41 | 57 |
| | Total number of years of follow-up | 3571.9 | 3464.4 | 3371.7 | 3417.2 | 3165.2 |
| | Event rates (per 100 person-years) | 0.87 | 1.27 | 1.19 | 1.20 | 1.80 |
| PAD | No. of events | 12 | 18 | 10 | 20 | 46 |
| | Total number of years of follow-up | 3627.9 | 3528.8 | 3451.7 | 3493.9 | 3181.9 |
| | Event rates (per 100 person-years) | 0.33 | 0.51 | 0.29 | 0.57 | 1.45 |
| Ischemic stroke | No. of events | 9 | 13 | 19 | 27 | 22 |
| | Total number of years of follow-up | 3626.8 | 3542.0 | 3416.5 | 3454.0 | 3250.3 |
| | Event rates (per 100 person-years) | 0.25 | 0.37 | 0.56 | 0.78 | 0.68 |