Supplementary Table S2. Comparisons of Optimal Cutoffs of Anthropometric Measures, and Adjusted HRs, 95% Confidence Intervals and *p*-Values of High- vs. Low-Risk Groups Determined Based on the Cutoffs Between All Cancers With and Without Breast Cancer

Measures	Stratification	All cancers			All cancers except breast cancer		
		Cutoff	HR	95%CI	Cutoff	HR	95%CI
Weight (kg)	Overall HT ever	89 89	1.18 1.15	(1.14–1.23) (1.09–1.22)	89	1.21	(1.15–1.28)
	HT never	82	1.25	(1.19–1.31)	82	1.22	(1.15-1.30)
BMI (kg/m ²)	Overall HT ever	31 34	1.16 1.09	(1.12–1.20) (1.02–1.16)	34	1.16	(1.11–1.22)
	HT never	30	1.25	(1.19-1.30)	36	1.21	(1.14-1.29)
Wt/Ht (kg/m)	Overall HT ever	53 53	1.19 1.10	(1.15–1.24) (1.04–1.17)	53	1.19	(1.13–1.25)
	HT never	51	1.24	(1.19-1.30)	52	1.25	(1.18-1.34)
WC (cm)	Overall HT ever HT never	94 94 88	1.16 1.11 1.12	(1.12–1.20) (1.06–1.17) (1.07–1.17)	102 94 102	1.15 1.10 1.20	(1.10–1.20) (1.03–1.17) (1.13–1.27)
WC/Ht (cm/cm)	Overall HT ever HT never	0.61 0.56 0.60	1.17 1.10 1.22	(1.13–1.21) (1.05–1.16) (1.16–1.28)	0.61 0.58 0.61	1.16 1.09 1.21	(1.11–1.20) (1.02–1.16) (1.15–1.29)
WHR (cm/cm)	Overall HT ever HT never	0.80 0.80 0.77	1.09 1.04* 1.20	(1.06–1.12) (1.00–1.08) (1.14–1.08)	0.80 0.92 0.77	1.10 1.04* 1.14	(1.06–1.14) (0.99–1.09) (1.08–1.20)

^{*}Not significant with p > 0.05.