

**Supplementary Table 1. Comparison of the percentage of bronchoalveolar lavage fluid neutrophils in 166 non-HIV-infected patients with *Pneumocystis jirovecii* pneumonia who did and did not have concomitant CMV infection.**

	<b>All (n = 166)</b>	<b>Mild-to- moderate<sup>a</sup> (n = 46)</b>	<b>Severe<sup>b</sup> (n = 120)</b>	<b>P</b>
<b>With CMV infection (n = 43)</b>	25.1 (8.2-47.0)	11.0 (3.1-21.3)	38.5 (11.0-60.5)	0.01
<b>Without CMV infection (n = 123)</b>	10.9 (4.3-28.7)	6.6 (2.5-13.8)	17.0 (6.7-.34.4)	0.001

**NOTE.** Data are shown as median % (interquartile range) of neutrophils. CMV, cytomegalovirus.

<sup>a</sup> 11 patients had concomitant CMV infection.

<sup>b</sup> 32 patients had concomitant CMV infection.

**Supplementary Table 2. Comparison of 30-day and 60-day all-cause mortalities among 166 non-HIV-infected patients with *Pneumocystis jirovecii* pneumonia who did and did not have concomitant CMV infection.**

	<b>With CMV infection</b>	<b>Without CMV infection</b>	
	<b>(n = 43)</b>	<b>(n = 123)</b>	<b><i>P</i></b>
<b>30-day all-cause mortality</b>	11 (25.6)	25 (20.3)	0.47
<b>60-day all-cause mortality</b>	15 (34.9)	34 (27.6)	0.37

**NOTE.** Data are shown as no. (%) of patients. CMV, cytomegalovirus.