

Table S1: List of strains used in this study

Strain	Genotype	Reference
Hu29	<i>h- ade6-M210 leu1-32 ura4-D18</i>	[1]
Hu1867	<i>h+ fft3::hph ade6-M210 uraDS/E leu1-32</i>	[1]
Hu1824	<i>h- pht1-myc::KanMx6 leu1-32 ade6-M210 ura4-D18</i>	[3]
Hu1993	<i>h- fft3::hph pht1-myc::KanMx6 leu1-32 ade6-M210 ura4-</i>	[3]
Hu2130	<i>h- Man1-7myc-Dam:NAT (nmt41 promoter)</i>	[4]
Hu2078	<i>h- 7myc-Dam-NLS:NATr (nmt41 promoter)</i>	[4]
Hu2193	<i>fft3::hph Man1-7myc-Dam:Nat (nmt41 promoter)</i>	This study
Hu2206	<i>fft3::hph 7myc-Dam-NLS:NATr (nmt41 promoter)</i>	This study
Hu2417	<i>h- fft3-K418R-myc::hph ade6-M210, ura4-D18, leu1-32</i>	This study
Hu1911	<i>h+ fft3-myc::hph leu1-32 ade6-210 ura4-D18</i>	This study
Hu1271	<i>h+ fft3-TAP Kan-MX6 leu1-32 ade6-210 ura4-D18</i>	This study
Hu2330	<i>h- sfc4-myc::kan ade6-M210 leu1-32 ura4-D18</i>	This study
Hu2343	<i>sfc4-myc::kan fft3-TAP::hph ade6-M210 leu1-32 ura4-D18</i>	This study
Hu2192	<i>h+ nup61::kan, ade6-M216, ura4-D18, leu1-32</i>	Bioneer Deletion Collection
Hu2173	<i>h+ man1::kan, ade6-M216, ura4-D18, leu1-32</i>	Bioneer Deletion Collection
Hu2413	<i>cnd2-GFP::NatR</i>	This study
Hu2414	<i>cnd2-GFP::NatR fft3D::HygR</i>	This study
Hu802	<i>h- clr6-1 (ts)</i>	[5]
Hu2108	<i>clr6-1 (ts) fft3::hph ade6-M210? uraDS/E? leu1-32?</i>	This study
Hu369	<i>h- clr3::kanMX ade6-M210, ura4-D18, leu1-32</i>	[5]
Hu2102	<i>fft3::hph clr3::KanMX adeM210 leu1-32 ura4-</i>	This study
Hu1026	<i>h- hos2::LEU2 leu1-32</i>	[5]
Hu2117	<i>fft3::hph hos2::leu2 leu1-32 ade6M210? uraDS/E?</i>	This study
Hu2424	<i>h+ otr1R(SphI)::ura4+ ura4-DS/E ade6-M210 leu1Δ::nmt1(81x)-dam-myc-nup85-kan</i>	[2]
Hu2422	<i>h+ otr1R(SphI)::ura4+ ura4-DS/E ade6-M210 leu1Δ::nmt1(81x)-dam-myc-kan</i>	[2]
Hu2438	<i>fft3::HygR leu1D::nmt1(81x)-dam-myc-nup85-KanR</i>	This study
Hu2434	<i>fft3::HygR leu1D::nmt1(81x)-dam-myc-KanR</i>	This study
Hu2582	<i>h90 taz1-mCherry::kanR hus5+::ccr1N-GFP::LEU2 leu1-32 lys1-131 ura4-D18 ade6-m210</i>	[6]
Hu2583	<i>h90 taz1-mCherry::kanR hus5+::ccr1N-GFP::LEU2 leu1-32 bqt4D::ura4+ lys1-131 ura4-D18 ade6-m210</i>	[6]
Hu2575	<i>h90 taz1-mCherry::kanR hus5+::ccr1N-GFP::LEU2 leu1-32 fft3D::HygR lys1-131 ura4-D18 ade6-m210</i>	This study
Hu2595	<i>h90 taz1-mCherry::kanR hus5+::ccr1N-GFP::LEU2 leu1-32 bqt4D::ura4+ fft3D::HygR lys1-131 ura4-D18 ade6-m210</i>	This study
Hu2241	<i>h- Nup61-myc::Kan ade6-M210 ura4-D18 leu1-32</i>	This study
Hu2256	<i>h- Nup61-myc::Kan fft3D::Hyg ade6-M210 ura4-D18orDS/E leu1-32</i>	This study
Hu2530	<i>h- Nup61-myc::Kan man1D::Kan ade6-M21X ura4-D18 leu1-32</i>	This study
Hu2566	<i>h- Nup61-myc::Kan bqt4D::ura4+ ade6-M21X ura4-D18 leu+</i>	This study
Hu2568	<i>h- Nup61-myc::Kan bqt4D::ura4+ fft3D::Hyg ade6-M21X ura4-D18orDS/E leu1-32</i>	This study
Hu2446	<i>fft3-K418R-myc::Hyg 7myc-Dam-NLS:NAT (nmt41 promoter)</i>	This study
Hu2448	<i>fft3-K418R-myc::Hyg Man1-7myc-Dam:NAT (nmt41 promoter)</i>	This study
Hu2771	<i>h- pht1-GFP::NAT leu1-32 ura4-</i>	[7]
Hu2791	<i>pht1-GFP::NAT fft3::HygR</i>	This study
Hu2797	<i>pht1-GFP::NAT fft3-K418R-myc::HygR</i>	This study
Hu2779	<i>fft3-myg::HygR ura4+</i>	This study
Hu2803	<i>fft3-myg::HygR ura4::ura4-LTR</i>	This study

References in Supplemental Material

1. Strålfors A, Walfridsson J, Bhuiyan H, Ekwall K (2011) The FUN30 chromatin remodeler, Fft3, protects centromeric and subtelomeric domains from euchromatin formation. *PLoS Genet* 7: e1001334. doi:10.1371/journal.pgen.1001334.
2. Woolcock KJ, Stunnenberg R, Gaidatzis D, Hotz H-R, Emmerth S, et al. (2012) RNAi keeps Atf1-bound stress response genes in check at nuclear pores. *Genes & Development* 26: 683–692. doi:10.1101/gad.186866.112.
3. Buchanan L, Durand-Dubief M, Roguev A, Sakalar C, Wilhelm B, et al. (2009) The Schizosaccharomyces pombe JmjC-protein, Msc1, prevents H2A.Z localization in centromeric and subtelomeric chromatin domains. *PLoS Genet* 5: e1000726. doi:10.1371/journal.pgen.1000726.
4. Steglich B, Filion GJ, van Steensel B, Ekwall K (2012) The inner nuclear membrane proteins Man1 and Ima1 link to two different types of chromatin at the nuclear periphery in *S. pombe*. *Nucleus* 3: 77–87.
5. Wirén M, Silverstein RA, Sinha I, Walfridsson J, Lee H-M, et al. (2005) Genome wide analysis of nucleosome density histone acetylation and HDAC function in fission yeast. *The EMBO Journal* 24: 2906–2918.
6. Chikashige Y, Yamane M, Okamasa K, Tsutsumi C, Kojidani T, et al. (2009) Membrane proteins Bqt3 and -4 anchor telomeres to the nuclear envelope to ensure chromosomal bouquet formation. *The Journal of Cell Biology* 187: 413–427. doi:10.1083/jcb.200902122.
7. Ogiyama Y, Ohno Y, Kubota Y, Ishii K (2013) Epigenetically induced paucity of histone H2A.Z stabilizes fission-yeast ectopic centromeres. *Nature Structural & Molecular Biology* 20: 1397–1406. doi:10.1038/nsmb.2697.