

**S3 Table:** Individual experts' final answers to four questions eliciting their beliefs about the relevance of the MYCYC trial results for informing opinion about 6-month remission rates in the MYPAN trial. Individual and consensus answers were reached after a summary of the design of MYCYC was provided, but before the actual MYCYC results were revealed.

<b>Expert</b>	<b>Q1</b>	<b>Q2</b>	<b>Q3</b>	<b>Q4</b>
1	0.60	0.50	0.65	0.60
2	0.50	0.20	0.50	0.20
3	0.55	0.40	0.50	0.50
4	0.65	0.20	0.35	0.15
5	0.50	0.20	0.40	0.60
6	0.50	0.50	0.50	0.50
7	0.55	0.40	0.55	0.40
8	0.60	0.10	0.60	0.10
9	0.50	0.30	0.30	0.20
10	0.50	0.20	0.50	0.20
11	0.50	0.50	0.50	0.50
12	0.70	0.20	0.60	0.10
13	0.50	0.30	0.30	0.40
14	0.60	0.25	0.25	0.25
15	0.60	0.20	0.60	0.20
<b>Mean</b>	<b>0.56</b>	<b>0.30</b>	<b>0.47</b>	<b>0.33</b>
<b>Median</b>	<b>0.55</b>	<b>0.25</b>	<b>0.50</b>	<b>0.25</b>
<b>Consensus values<sup>†</sup></b>	<b>0.55</b>	<b>0.25</b>	<b>0.5</b>	<b>0.25</b>

**Q1:** What is the chance that the 6-month remission rate on CYC/steroids in the MYCYC patient group exceeds that in the MYPAN patient group?

**Q2:** What is the chance that the 6-month remission rate on CYC/steroids in the MYPAN patient group exceeds that in the MYCYC patient group by more than 10%?

**Q3:** As Q1 above but concerning MMF/steroids instead of CYC/steroids.

**Q4:** As Q2 above but concerning MMF/steroids instead of CYC/steroids.

<sup>†</sup>Once individuals had finalised their answers to Q1-Q4, median values were calculated and it was proposed that these be taken forward as a working summary of the group's consensus opinion. Once the MYCYC data were revealed, median values of answers to Q1-Q4 were used to incorporate these data into the prior distributions for  $\theta$ ,  $p_C$  and  $p_E$  based on expert opinion alone. Through constructive discussions, experts then reached the consensus that densities incorporating MYCYC data had face validity for representing their beliefs (93% agreement; no abstentions permitted).