

**Online Supporting Material**

**SUPPLEMENTAL TABLE 1** Associations between mothers' child-feeding practices and child's sugar-sweetened beverage<sup>1</sup> (SSB) intake among 6 y old stratified by child's weight status — Infant Feeding Practices Year 6 Follow-Up Study, 2012 (n=1350)

Mothers' child-feeding practices <sup>3</sup>	SSB intake during the past month at 6 y of age by child's weight status						P value <sup>a</sup>		
	Underweight/normal weight children			Overweight/obese children					
	0 times/d	>0 to <1 time/d	≥1 time/d	0 times/d	>0 to <1 time/d	≥1 time/d			
	n	% <sup>2</sup>	time/d % <sup>2</sup>	% <sup>2</sup>	time/d % <sup>2</sup>	n	% <sup>2</sup>	time/d % <sup>2</sup>	P value <sup>a</sup>
Make sure that my child does not eat too many sweets or junk foods	1031					319			0.0001
No	113	14.2	54.9	31.0	57.8	45	4.44	37.8	
Yes	918	21.2	63.4	15.4	64.2	274	15.0	20.8	
If I did not guide or regulate my child's eating, my child would eat too much of his/her favorite foods	1031					319			0.003
No	349	25.8	60.7	13.5	57.6	92	16.3	26.1	
Yes	682	17.7	63.3	18.9	65.6	227	12.3	22.0	
Especially careful to make sure my child eats enough	1031					319			0.07
No	367	23.2	62.9	13.9	63.8	127	16.5	19.7	
Yes	664	19.0	62.2	18.8	63.0	192	11.5	25.5	
Encourage your 6-y-old to eat all of the food on his/her plate	1031					319			0.08
No	419	23.6	61.1	15.3	65.5	148	13.5	21.0	
Yes	612	18.3	63.4	18.3	61.4	171	13.5	25.2	

<sup>1</sup> Include regular soda, sweetened drinks such as Kool-Aid™, lemonade, sweet tea, Hi-C™, cranberry cocktail, Gatorade™, and others.

<sup>2</sup> Percentages may not add up to 100% because of rounding.

<sup>3</sup> “No” included “1”, “2”, and “3” on the Likert scale and “Yes” included “4” and “5” on the Likert scale.

<sup>a</sup> Based on  $\chi^2$  tests.