

### Appendix 3 (as supplied by the authors): Characteristics of included studies

<b>Study/Location</b>	Amaro 2006 [1] Italy
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	Baranowski 2003 [3] United States
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	Barkin 2012 [4] United States
<b>Objective</b>	To test the effect of a culturally tailored, family-centered, short-term behavioural intervention on BMI in Latino-American preschool-aged children
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: bilingual research assistant approached individuals in the waiting area of community agencies (e.g., pediatric clinics, community centers); study advertised via: flyers, community organizations; Spanish radio, Spanish newspapers, word-of-mouth</p> <p>Inclusion criteria: parents &gt;18 years; self-defined as Hispanic/Latino; child aged 2-6; not currently enrolled in another healthy lifestyle program; valid telephone number; planning on remaining in the city for the next 6 months</p> <p>Unit of allocation: parent-child dyads</p> <p>Unit of analysis: child</p> <p>Intention to treat: yes</p>
<b>Participants</b>	<p>Sample: 106 dyads [95% of dyads that completed study (71/75) included mothers]</p> <p>Intervention n=54; Control n=52</p> <p>Age (child) mean (SD) (years): Intervention: 4.2 (0.9); Control: 4.1 (0.9)</p> <p>Gender (child) [Female n (%): Intervention n=16 (45.7%); Control n=22 (55%)</p> <p>Loss to follow-up: Intervention n=19; Control n=12</p>
<b>Intervention</b>	<p>Description of intervention: Salud Con La Familia (Heart with the family); 12 weekly 90-minute skills-building sessions for parents and preschool-aged children to improve nutritional habits, increase physical activity, and decrease sedentary activity</p> <p>Description of control: brief school readiness program conducted as alternative to active intervention; met 3 times for 60 minutes over the 12-week study period</p> <p>Duration of intervention: 12 weeks</p> <p>Length of follow-up: immediate post</p>
<b>Study/Location</b>	Beech 2003 [5] United States; Companion paper: Story [6]
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details

<b>Study/Location</b>	Bellows 2013 [7] United States
<b>Objective</b>	To assess the efficacy of an intervention on gross motor skill performance, physical activity, and weight status of preschoolers
<b>Methods</b>	Design: RCT Selection: NR Unit of allocation: child Unit of analysis: child Intention to treat analysis: no
<b>Participants</b>	Sample: 263 Intervention n=132; Control n=131 Age mean (SD) (months): Intervention: 53.0 (6.8); Control: 51.5 (6.6) Gender (Female): 45% Race/Ethnicity: 59% Hispanic SES: all participants considered to have low SES Loss to follow-up: Intervention n=34; Control n=28
<b>Intervention</b>	Description of intervention: “The Food Friends: Get Movin’ With Mighty Moves”: 18 weeks classroom based intervention 4 days/week for 15–20 min/day; 72 lessons that comprised multiple activities (143 total activities) focused on gross motor skill and healthy eating; led by classroom teacher Description of control: NR Duration of intervention: 18 weeks Length of follow-up: immediate post
<b>Study/Location</b>	Black 2010 [8] United States
<b>Objective</b>	To evaluate a 12-session home/community-based health promotion/obesity prevention program (Challenge!) on changes in BMI, body composition, physical activity, and diet
<b>Methods</b>	Design: RCT Selection: one group participated in investigation of growth and development; other group recruited from middle schools; researchers visited classes and described the project and the possibility of receiving a health program with a “personal trainer” Inclusion criteria: aged 11-16 years; reside in nearby low-income communities Unit of allocation: individuals Unit of analysis: individuals Intention to treat: No
<b>Participants</b>	Sample: 235 Intervention n=121; Control n=114 Age mean (SD) (years): Intervention: 13.3 (1.0); Control: 13.3 (1.0)

	<p>Gender [Female n (%): Intervention n=62 (51.2%); Control n=54 (47.4%)</p> <p>Race/Ethnicity: Non-Hispanic blacks: Intervention: 118 (97.5%); Control: 110 (96.5%)</p> <p>Loss to follow-up: Intervention n=30; Control n=21</p>
<b>Intervention</b>	<p>Description of intervention: 12 sessions that included a challenge (e.g., persuade someone to drink water instead of soda), setting a personal goal related to diet or PA (e.g., eat 2 vegetables/day, walk 30 min/day), make and taste healthy snacks and engage in PA with mentors; taste tests (e.g., regular/diet soda), recipes for healthy snacks (e.g., breakfast sundae with yogurt, granola and fruit), and recommendations for PA; parents welcome to participate; mentors left recipes and information for the family</p> <p>Description of control: no mentor, no contact between baseline and follow-up</p> <p>Duration of intervention: 11 months</p> <p>Length of follow-up: 24 months</p>
<b>Study/Location</b>	Bonsergent 2013 [9] France; Companion paper: Briancon [10]
<b>Objective</b>	To evaluate the 2-year effectiveness of three strategies aimed at preventing overweight and obesity among adolescents in a high school setting
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: 24 high schools randomly selected after stratification on department and type of education (general and technological or vocational)</p> <p>Inclusion criteria: high school must be a state administrated establishment</p> <p>Unit of allocation: school</p> <p>Unit of analysis: child</p> <p>Intention to treat: no</p>
<b>Participants</b>	<p>Sample: 5,354</p> <p>Intervention n=2,641; Control n=2,713 (baseline and follow-up data only presented by completers and non-completers)</p> <p>Age mean (SD) (years): Completers: 15.6 (0.7)</p> <p>Gender (Female): completers 57.6%</p> <p>Loss to follow-up: 33.9% overall</p>
<b>Intervention</b>	<p>Description of intervention: dietary and physical activity lectures for 5 hours in Grade 10, 6 hours in Grade 11 and group work to exchange, find and present answers to problems related to eating habits, physical activity and the environment</p> <p>Description of control: non-education strategy</p> <p>Duration of intervention: 24 months</p> <p>Length of follow-up: immediate post</p>
<b>Study/Location</b>	Brandsetter 2012 [11] Germany
<b>Objective</b>	To describe the effects of a school-based intervention for overweight prevention on children's BMI and other measures of fat mass

<b>Methods</b>	<p>Design: RCT</p> <p>Selection: all principals were informed in writing about the study and were asked to invite first-grade teachers to participate</p> <p>Inclusion criteria: NR</p> <p>Unit of allocation: school</p> <p>Unit of analysis: child</p> <p>Intention to treat: no</p>
<b>Participants</b>	<p>Sample: 1,119</p> <p>Intervention n=540; Control n=579</p> <p>Age mean (SD) (years): Intervention: 7.61 (0.42); Control: 7.53 (0.42)</p> <p>Gender (Female): Intervention 44.9%; Control 47.9%</p> <p>Loss to follow-up: Intervention n=51; Control n=42</p>
<b>Intervention</b>	<p>Description of intervention: school-based, within existing curriculum focused on health promoting behaviour change with action alternatives in three areas: drinking sugar-sweetened beverages (drinking water, discovering hidden sugar in drinks), spending time with screen media (leisure activities without TV), and being physically active (learning about local sport and leisure facilities); 1 school year of materials covering: 29 30-60 minute teaching units; 2 short (5-7 minute) blocks of PA exercises a day, 6 family homework lessons; teacher training and parent information materials</p> <p>Description of control: no intervention</p> <p>Duration of intervention: 1 school year</p> <p>Length of follow-up: immediate post</p>
<b>Study/Location</b>	Brown 2013 [12] United States
<b>Objective</b>	To develop a lifestyle change program for Native American youth by modifying the Diabetes Prevention Program (DPP) and to assess implementation indicators and short term behavioural and physiological outcomes of the intervention among a pilot sample
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: Northern Plains Indian youth 10-14 years old living on 2 American Indian reservations in north-central and southwestern Montana</p> <p>Unit of allocation: child</p> <p>Unit of analysis: child</p> <p>Intention to treat: no</p>
<b>Participants</b>	<p>Sample: 76</p> <p>Intervention n=38; Control n=38</p> <p>Age mean (SD) (years): Overall: 11.4 (1.1)</p> <p>Gender (Female): 50%</p> <p>Race/Ethnicity: Native American</p>

	Loss to follow-up: Intervention n=6; Control n=6
<b>Intervention</b>	<p>Description of intervention: general content and behavioural were based on the original DPP lifestyle change model; strategies targeted healthy weight maintenance, lowering fat intake, increasing physical activity; 9 sessions over 3 months; weight goal to slow or reduce BMI growth; emphasis on traditional activities (e.g., berry picking, horseback riding, dancing, hunting, hiking, and camping), use of storytelling and native language to convey information, and participation of elders; hands-on interactive activities</p> <p>Description of control: addressed risks for alcohol and drug use</p> <p>Duration of intervention: 3 months</p> <p>Length of follow-up: immediate post</p>
<b>Study/Location</b>	Burgi 2012 [13] Switzerland; Companion papers: Puder [14], Niederer [15]
<b>Objective</b>	To examine whether a multidimensional lifestyle intervention is equally effective in children of migrant and/or low educational level parents
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: public preschool classes randomly selected in areas with a high migrant population from two different socio-cultural and linguistic regions in Switzerland</p> <p>Inclusion criteria: for preschool classes a &gt;40% prevalence of migrant children and no participation in any other prevention project</p> <p>Unit of allocation: class</p> <p>Unit of analysis: children</p> <p>Intention to treat: yes</p>
<b>Participants</b>	<p>Sample: 652</p> <p>Intervention n=342; Control n=310</p> <p>Age mean (SD) (years): Intervention: 5.2 (0.6); Control: 5.2 (0.6)</p> <p>Gender [Female n (%): Intervention n=167 (49%); Control n=159 (51%)</p> <p>Loss to follow-up: Intervention n=18; Control n=9</p>
<b>Intervention</b>	<p>Description of intervention: children participated in a PA program consisting of four 45 min sessions per week; teachers participated in two 3 hour workshops to learn the content and practical aspects of the intervention and in one informal meeting to exchange experiences; parents participated in three interactive information and discussion evenings about promotion of PA, healthy food, limiting TV use and the importance of sufficient sleep</p> <p>Description of control: Regular school curriculum</p> <p>Duration of intervention: 10 months</p> <p>Length of follow-up: immediate post</p>
<b>Study/Location</b>	Caballero 2003 [16] United States
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details

<b>Study/Location</b>	Campbell 2013 [17] Australia; Companion paper: Campbell [18]
<b>Objective</b>	To assess the effectiveness of a parent-focused intervention on infants' obesity-risk behaviours and BMI
<b>Methods</b>	Design: RCT Selection: 14 local government areas randomly selected from 28 eligible; fifty percent of eligible first-time parents' groups in each area randomly selected (62/103 groups) and approached by research staff during a standard nurse-facilitated group session Inclusion criteria: parent groups eligible if $\geq 8$ parents enrolled or $\geq 6$ parents enrolled in areas of low SES Unit of allocation: parent group Unit of analysis: child Intention to treat: yes
<b>Participants</b>	Sample: 542 Intervention n=271; Control n=271 Age mean (SD) (months): Overall: 3.9 (1.6) Gender (Female): 47.4% Loss to follow-up: Intervention n=30; Control n=32
<b>Intervention</b>	Description of intervention: dietitian-delivered intervention comprised six 2-hour sessions delivered quarterly during the first-time parents' group regular meeting Description of control: 6 newsletters on non obesity-focused themes; parents received usual care from their MCH nurse, who may have provided lifestyle advice. Duration of intervention: 15 months Length of follow-up: immediate post
<b>Study/Location</b>	Crespo 2012 [19] United States; Companion paper: Elder [20]
<b>Objective</b>	To evaluate the impact of a community health advisor intervention to promote healthy eating and physical activity and prevent excess weight gain among Latino children
<b>Methods</b>	Design: RCT Selection: elementary schools within 3 school districts in south San Diego County Unit of allocation: school Unit of analysis: child Intention to treat: yes
<b>Participants</b>	Sample: 808 Intervention 1 n=198; Intervention 2 n=165; Intervention 3 n=218; Control n=227 Age mean (SD) (years): Overall: 5.9 (0.9) Gender (Female): 50% Loss to follow-up: Intervention 1 n=31; Intervention 2 n=20; Intervention 3 n=22;

	Control n=22
<b>Intervention</b>	<p>Description of intervention 1: home/family environmental change – activities delivered by eight promotoras (community health advisors) consisted of home visits, newsletters, recipe cards, goal setting, booster phone calls</p> <p>Description of intervention 2: community-only environmental change – school playground improvements and implementation of salad bars/improvements to salad bars, improvements to community parks, change in classroom practices, physical education equipment, children’s menus at restaurants</p> <p>Description of intervention 3: family-plus-community-environmental change – combination of interventions 1 and 2</p> <p>Description of control: no intervention</p> <p>Duration of intervention: 3 years</p> <p>Length of follow-up: immediate post</p>
<b>Study/Location</b>	Cunha 2013 [21] Brazil
<b>Objective</b>	To evaluate the effectiveness of a school-based intervention involving families and teachers to promote healthy eating habits in adolescents and reduce increase in BMI
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: selected 20 schools with fifth grade classes out of 35 municipal schools; all located in areas not considered high risk for violence</p> <p>Unit of allocation: school</p> <p>Unit of analysis: child</p> <p>Intention to treat: yes</p>
<b>Participants</b>	<p>Sample: 574</p> <p>Intervention n=293; Control n=281</p> <p>Age mean (SD) (years): Intervention: 11.2 (1.3); Control: 11.2 (1.3)</p> <p>Gender (Female): 48.6%</p> <p>Loss to follow-up: Intervention n=45; Control n=30</p>
<b>Intervention</b>	<p>Description of intervention: trained nutritionists gave monthly 1 hour sessions in the classrooms, which included games, theater sketches, movies and puppet shows, writing and drawing contests, to encourage changes in eating habits and food consumption</p> <p>Description of control: one-hour session of orientation on general health and advice on healthy eating, at the end of the study</p> <p>Duration of intervention: 9 months</p> <p>Length of follow-up: immediate post</p>
<b>Study/Location</b>	Daniels 2012 [22] Australia
<b>Objective</b>	To evaluate a universal obesity prevention intervention for infants

<b>Methods</b>	<p>Design: RCT</p> <p>Selection: recruitment 4 hospitals in Adelaide and 3 in Brisbane; consecutive sample of first-time mothers (<math>\geq 18</math> years old) who delivered a healthy term infant approached while still in hospital by hospital employed midwives, study-employed staff, or doctoral students; mothers who gave consent re-contacted when infant was 4-6 months</p> <p>Inclusion criteria: no documented history of domestic violence or intravenous drug use; no self-reported eating or psychiatric disorder; written and spoken English; ability to attend sessions; no serious infant health problems; score on the Kessler Psychological Distress Scale (K10) below 30 (not high maternal psychological distress).</p> <p>Unit of allocation: child</p> <p>Unit of analysis: child</p> <p>Intention to treat: yes</p>
<b>Participants</b>	<p>Sample: 698</p> <p>Intervention n=352; Control n=346</p> <p>Age mean (SD) (months): Intervention: 4.3 (1.0); Control: 4.3 (1.0)</p> <p>Gender (Female): Intervention 51%; Control 50%</p> <p>Loss to follow-up: Intervention n=92; Control n=65</p>
<b>Intervention</b>	<p>Description of intervention: comprehensive skills-based program that used a cognitive behavioural approach and focused on feeding and parenting practices; 2 modules of 6 fortnightly group sessions (10–15 mothers per group), each 1 to 1.5 hours; Module 1 delivered by 9 dietitians and 10 psychologists who worked in pairs</p> <p>Description of control: self-directed access to usual community child health services, which were similar in both states and largely targeted at high-risk families</p> <p>Duration of intervention: 3 months</p> <p>Length of follow-up: 15 months</p>
<b>Study/Location</b>	DeBar 2011 [23] United States; Companion paper: The HEALTHY Study Group [24]
<b>Objective</b>	To examine whether student's "public commitment" - voluntary participation as a peer communicator or in student-generated media opportunities - in a school-based intervention to prevent diabetes and reduce obesity predicted improved study outcomes
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: schools where at least 50% of children ineligible for federally subsidized, free, or reduced-priced meals and/or at least 50% of students' ethnicity was Black or Hispanic. Students enrolled in 6th grade in Fall 2006 who had no conditions that would preclude active participation in physical education classes</p> <p>Unit of allocation: school</p> <p>Unit of analysis: child</p> <p>Intention to treat: no</p>
<b>Participants</b>	Sample: 3,131



	<p>Intervention n=835; Control n=2,296</p> <p>Age mean (SD) (years): Intervention: 11.3 (0.5); Control: 11.3 (0.5)</p> <p>Gender (Female): Intervention 58.6%; Control 69.6%</p> <p>Race/Ethnicity: Intervention 51% Hispanic; Control 53.5% Hispanic</p> <p>Loss to follow-up: 0</p>
<b>Intervention</b>	<p>Description of intervention: HEALTHY intervention, delivered over five semesters (Spring 2007, Fall 2007, Spring 2008, Fall 2008, Spring 2009) comprised four components: nutrition, physical education, behaviour and communications</p> <p>Description of control: no intervention</p> <p>Duration of intervention: 3 years</p> <p>Length of follow-up: immediate post</p>
<b>Study/Location</b>	de Heer 2011 [25] United States
<b>Objective</b>	To evaluate the effectiveness and spillover of an after-school health education and physical activity program among Hispanic elementary school children
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: students recruited in third, fourth, and fifth grades by making announcements and passing out consent forms during PE classes</p> <p>Exclusion criteria: children were excluded if they were not in the target grades and/or if they had a condition that would endanger their own or others' safety</p> <p>Unit of allocation: individual</p> <p>Unit of analysis: individual</p> <p>Intention to treat: no</p>
<b>Participants</b>	<p>Sample: 646</p> <p>Intervention n=292; Control n=354</p> <p>Age mean (SD) (years): Intervention: 9.24 (0.87); Control: 9.10 (1.08)</p> <p>Gender (Female): 47.0%</p> <p>Loss to follow-up: Intervention n=50; Control n=28</p>
<b>Intervention</b>	<p>Description of intervention: after-school program ran twice weekly for 12 weeks for a total of 24 sessions at each school; each session took place in the schoolyard or in the multipurpose room and comprised a 20 to 30 minute health education component followed by 45 to 60 minutes of physical activity.</p> <p>Description of control: no treatment</p> <p>Duration of intervention: 3 months</p> <p>Length of follow-up: immediate post</p>
<b>Study/Location</b>	De Coen 2012 [26] Belgium
<b>Objective</b>	To evaluate the effects of a school-based, 2-year, multi-component intervention on BMI,

	eating and physical activity behaviour
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: six communities selected from research regions in Flanders based on five socio-economic characteristics: (i) number of births in underprivileged families; (ii) proportion of pupils in primary school with a school delay; (iii) rate of unemployment; (iv) number of persons on welfare support; and (v) number of underprivileged foreigners; recruitment in schools; all pre-primary and primary schools invited</p> <p>Unit of allocation: community</p> <p>Unit of analysis: child</p> <p>Intention to treat: no</p>
<b>Participants</b>	<p>Sample: 3,242</p> <p>Intervention n=2,034; Control n=1,208</p> <p>Age mean (SD) (years): Intervention: 4.86 (1.25); Control: 5.04 (1.29)</p> <p>Gender (Female): 50%</p> <p>Loss to follow-up: Intervention n=1,364; Control n=766</p>
<b>Intervention</b>	<p>Description of intervention: based on the 'Nutrition and Physical Activity Health Targets': (i) increasing daily consumption of water and decreasing soft drinks consumption; (ii) increasing daily milk consumption; (iii) increasing daily consumption of vegetables and fruit; (iv) decreasing daily consumption of sweets and savoury snacks; and (v) increasing daily PA and decreasing screen-time behaviour</p> <p>Description of control: no intervention</p> <p>Duration of intervention: 24 months</p> <p>Length of follow-up: immediate post</p>
<b>Study/Location</b>	de Ruyter 2012 [27] Netherlands; Companion paper: de Ruyter [28]
<b>Objective</b>	To examine the effect on weight gain of masked replacement of sugar-sweetened beverages with non-caloric, artificially sweetened beverages
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: recruited children at eight urban elementary schools near Amsterdam</p> <p>Inclusion criteria: children who commonly drank sugar-sweetened beverages</p> <p>Exclusion criteria: children with various medical conditions</p> <p>Unit of allocation: child</p> <p>Unit of analysis: child</p> <p>Intention to treat: yes</p>
<b>Participants</b>	<p>Sample: 641</p> <p>Intervention n=319; Control n=322</p> <p>Age mean (SD) (years): Intervention: 8.2 (1.8); Control: 8.2 (1.8)</p> <p>Gender (Female): Intervention 46%; Control 47%</p>

	Loss to follow-up: Intervention n=94; Control n=50
<b>Intervention</b>	Description of intervention: children received a box at school each week containing 8 cans, 1 for each day plus 1 spare in case a can was misplaced; teachers checked to see whether the children consumed their beverage during the morning break in class and reminded them to take cans home for the weekend and any holidays. The sugar-free beverages contained 24 mg of sucralose and 12 mg of acesulfame potassium per can. Description of control: control beverage contained 26 g of sucrose Duration of intervention: 18 months Length of follow-up: immediate post
<b>Study/Location</b>	Donnelly 2009 [29] United States; Companion paper: Gibson [30]
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	Dzewaltowski 2010 [31] United States
<b>Objective</b>	To evaluate the prevention of childhood obesity through building the capacity of after-school staff to increase physical activity and fruit and vegetable opportunities
<b>Methods</b>	Design: RCT Selection: all schools participating in an after-school program alliance of the Lawrence Public School District Exclusion criteria: if after-school programs were not on the elementary school grounds Unit of allocation: school Unit of analysis: individual Intention to treat: no
<b>Participants</b>	Sample: 273 Intervention n=148; Control n=125 Age mean (SD) (years): Intervention: 9.34 (0.65); Control: 9.19 (0.66) Gender (Female): Intervention 53.0%; Control 46.0% SES (% eligible for free/reduced lunch): Intervention 44%; Control 58% Loss to follow-up: Intervention n=14; Control n=13
<b>Intervention</b>	Description of intervention: the HOP'N intervention model included three levels: a community/government/human service agency (County Cooperative Extension), after-school staff training, and after-school program quality elements. Description of control: standard after-school program Duration of intervention: 24 months Length of follow-up: immediate post
<b>Study/Location</b>	Ebbeling 2006 [32] United States
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details

<b>Study/Location</b>	El Ansari 2010 [33] Egypt
<b>Objective</b>	To assess the association between a PA intervention and three anthropometric parameters (weight, body mass index, body fat) and four physiological parameters (cholesterol level, systolic blood pressure, diastolic blood pressure, heart rate) among adolescents
<b>Methods</b>	Design: RCT Selection: schools with sport facilities and sport equipment Unit of allocation: individual Unit of analysis: individual Intention to treat: no
<b>Participants</b>	Sample: 160 Intervention n=80; Control n=80 Age mean (SD) (years): Intervention: 15.7 (1.8); Control: 15.4 (1.6) Gender [Female n (%): Intervention n=45 (56%); Control n=45 (56%) Loss to follow-up: Intervention n=0; Control n=0
<b>Intervention</b>	Description of intervention: three, 60-minute PA sessions each week for three months Description of control: no intervention Duration of intervention: 3 months Length of follow-up: immediate post
<b>Study/Location</b>	Escribano 2012 [34] Germany/Spain; Companion paper: Koletzko [35]
<b>Objective</b>	To assess if the increases in weight gain velocity and BMI induced by protein intake early in life are related to an increase in fat or fat-free mass
<b>Methods</b>	Design: RCT Selection: 80 infants from the EU Childhood Obesity Programme sample; 37 from Germany and 43 from Spain; selected by recruitment order from 522 eligible subjects when they were 6 months old Unit of allocation: school Unit of analysis: individual Intention to treat: N/A
<b>Participants</b>	Sample: 66 Intervention 1 n=17; Intervention 2 n=24; Control n=25 Age: NR Gender [Female n (%): Intervention 1 n=8 (47%); Intervention 2 n=14 (58%); Control n=10 (40%) Loss to follow-up: NR
<b>Intervention</b>	Description of intervention: 41 infants randomized at birth to higher or lower protein content formula (HP=17 and LP=24); 25 breastfed infants also included; anthropometric measures assessed at baseline, 6, 12 and 24 months, and fat-free mass

	(FFM) and fat mass (FM) were assessed by isotope dilution at 6 months. Duration of intervention: 6 months Length of follow-up: 12, 24 months
--	--

<b>Study/Location</b>	Fitzgibbon 2005 [36] United States; Companion paper: Fitzgibbon [37]
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	Fitzgibbon 2006 [37] United States
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	Fitzgibbon 2011 [38] United States
<b>Objective</b>	To assess the feasibility and effectiveness of a teacher-delivered weight control intervention for black preschool children
<b>Methods</b>	Design: RCT Selection: no details regarding school recruitment Inclusion criteria: intervention took place during regular class time so all children in participating classrooms received intervention and were eligible to participate Unit of allocation: school Unit of analysis: individual Intention to treat: no
<b>Participants</b>	Sample: 589 Intervention n=309; Control n=280 Age mean (years): Overall: 4.3 Gender (Female): Intervention 52%; Control 55% Race/Ethnicity: 94% Black, 3% Latino Loss to follow-up: overall n=29
<b>Intervention</b>	Description of intervention: 14 week curriculum based intervention, 2 teacher delivered sessions per week each week on a specific theme/objective (one 20-minute lesson and one 20-minute physical activity component); parent involvement: weekly newsletter with parallel content, homework assignment with \$ reward for completion Description of control: general health intervention Duration of intervention: 14 weeks Length of follow-up: immediate post
<b>Study/Location</b>	Fitzgibbon 2013 [39] United States
<b>Objective</b>	To test the feasibility of Family-Based Hip-Hop to Health, a school-based obesity prevention intervention for 3-5-year-old Latino children and their parents, and estimate

	its effectiveness in producing smaller average changes in BMI
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: principals and preschool teachers from four Chicago Public Schools agreed to allow children to participate. Two half-day classrooms from each school participated</p> <p>Unit of allocation: ECE program</p> <p>Unit of analysis: child</p> <p>Intention to treat: no</p>
<b>Participants</b>	<p>Sample: 147</p> <p>Intervention n=73; Control n=74</p> <p>Age mean (SD) (months): Overall: 54.2 (5.0)</p> <p>Gender (Female): 50%</p> <p>Race/Ethnicity: 94% Hispanic</p> <p>Loss to follow-up: Intervention n=12; Control n=7</p>
<b>Intervention</b>	<p>Description of intervention: child component included a 14-week (three times weekly) intervention led by a bilingual/bicultural educator; each session included 20 min of nutrition instruction (included activities led by puppets) and 20 min of aerobic activity; parent component included classes and newsletters adapted for a lower-income, Hispanic population; parents encouraged to attend six weekly 90-min classes that included 60 min of interactive instruction on healthful eating and family exercise plus 30 min of moderate physical activity (e.g., salsa aerobics, walking group)</p> <p>Description of control: control schools received a once weekly intervention for 14 weeks (20 min each week) that taught general health concepts such as dental health, seat belt safety, and calling 911; parents received parallel weekly newsletters</p> <p>Duration of intervention: 14 weeks</p> <p>Length of follow-up: immediate post; 12 months</p>
<b>Study/Location</b>	Foster 2008 [40] United States
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	French 2011 [41] United States; Companion papers: Foster [42], The HEALTHY study group [24]
<b>Objective</b>	To evaluate an intervention to prevent weight gain among households
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: households recruited over 8 months; recruitment sources: community libraries, worksites, schools, daycare centers, health clinics, religious institutions, park and recreation centers, grocery stores, and food co-ops</p> <p>Exclusion criteria: living too far from the university, household TV viewing hours below enrollment criteria, household configuration not meeting enrollment criteria</p> <p>Unit of allocation: household</p>

	Unit of analysis: household/individual Intention to treat: no
<b>Participants</b>	Sample: 90 households Intervention n=45 households; Control n=45 households Age: aged <5 years n=23, aged 5-11 years n=84, aged 12-17 years n=75 Gender [Female n (%)]: only reported for adults as main respondents 93% Loss to follow-up: overall 4 households
<b>Intervention</b>	Description of intervention: 6 monthly face-to-face group sessions, monthly newsletters, and 12 home-based activities Description of control: no treatment Duration of intervention: 1 year Length of follow-up: immediate post
<b>Study/Location</b>	Fung 2012 [43] Canada
<b>Objective</b>	To examine the effectiveness of a Comprehensive School Health program by evaluating temporal changes in diets, activity levels and body weight
<b>Methods</b>	Design: RCT Selection: 10 schools selected from five jurisdictions in Alberta, all of which agreed to support healthy eating and active living initiatives among students Exclusion criteria: schools outside selected jurisdictions Unit of allocation: school Unit of analysis: child Intention to treat: N/A
<b>Participants</b>	Sample: 3,714 Intervention n=293; Control n=3,421 Age: NR (grade 5 students) Gender [Female n (%)]: Intervention n=149 (50.7%); Control n=1,762 (51.5%) Loss to follow-up: <10% dropout rate by the 2010 survey
<b>Intervention</b>	Description of intervention: “to make the healthy choice the easy choice” School Health Facilitators implemented healthy eating and active living strategies; engaged all stakeholders, including parents, staff and community; School Health Facilitators developed cross curriculum links and taught across curriculum; facilitated professional development days for teachers and staff, organized parent information nights, nutrition programs such as cooking clubs, after school physical activity programs, weekend events and celebrations, and circulated newsletters Description of control: no intervention Duration of intervention: 3 years Length of follow-up: -1 year

<b>Study/Location</b>	Gentile 2009 [44] United States
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details

<b>Study/Location</b>	Greening 2011 [45] United States
<b>Objective</b>	To evaluate a healthy lifestyle school-based obesity intervention in a rural southern community
<b>Methods</b>	Design: RCT Exclusion criteria: disabilities that precluded comprehending the questionnaires or performing the fitness tests Unit of allocation: school Unit of analysis: school Intention to treat: no
<b>Participants</b>	Sample: 450 Intervention n=204; Control n=246 Age mean (SD; range) (years): Overall: 8.34 (1.30; 6 to 10) Gender (Female): overall 48.0% Loss to follow-up: overall 11.0%
<b>Intervention</b>	Description of intervention: a 45 minute nutritional information session presented once during school year by a nutritionist; 45 minute physical education classes twice a week; healthy information incorporated into weekly class lectures; deep frying equipment replaced with baking ovens Description of control: standard health curriculum Duration of intervention: 8 months Length of follow-up: immediate post

<b>Study/Location</b>	Haerens 2006 [46] Belgium; Companion paper: Haerens [47]
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details

<b>Study/Location</b>	Hakanen 2010 [48] Finland
<b>Objective</b>	To evaluate the impact of individualized dietary and lifestyle counselling, primarily aimed to decrease serum low-density lipoprotein cholesterol, on the clustering of overweight-related cardiometabolic risk factors in children
<b>Methods</b>	Design: RCT Selection: recruited by nurses at well baby visit Exclusion criteria: children with chronic disease (e.g. chromosomal diseases, diabetes, familial hypercholesterolaemia)



	Unit of allocation: child Unit of analysis: child Intention to treat: no
<b>Participants</b>	Sample: 1,062 Intervention n=540; Control n=522 Age mean (months): Intervention: 7; Control: 7 Gender: NR Loss to follow-up: Intervention n=291; Control n=246
<b>Intervention</b>	Description of intervention: individualized dietary and lifestyle counselling at 1 to 3 month intervals until child was 2 years old and twice a year thereafter; all children continued regular visits at the wellbaby clinics and school health care for vaccinations, growth and development follow-up and basic health education Description of control: contacted by the counselling team twice a year until age 7 years and once a year after that; received similar basic health education as routinely given at Finnish wellbaby clinics and school health care Duration of intervention: 2 years Length of follow-up: every two years for 8 years
<b>Study/Location</b>	Harvey-Berino 2003 [49] United States
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	HEALTHY Study Group 2010 [42] United States
<b>Objective</b>	To examine the effects of a multicomponent, school-based program addressing risk factors for diabetes among children whose race or ethnic group and SES placed them at high risk for obesity and type 2 diabetes
<b>Methods</b>	Design: RCT Selection/Exclusion criteria: $\geq 50\%$ of children in school eligible for federally subsidized, free or reduced-price meals or $\geq 50\%$ students black or Hispanic; Black and Hispanic children of lower SES oversampled given that these children are at a high risk for obesity and type 2 diabetes; Students in 6th grade in fall 2006 eligible if no diabetes or conditions that would preclude regular participation in physical education Unit of allocation: school Unit of analysis: individual Intention to treat: no
<b>Participants</b>	Sample: 6,358 Intervention n=3,189; Control n=3,169 Age mean (SD) (years): Intervention: 11.3 (0.5); Control: 11.3 (0.6) Gender (Female): Intervention 52.6%; Control 52.9%

	Loss to follow-up: overall 27.6%
<b>Intervention</b>	Description of intervention: four integrated components: nutrition, physical activity, behavioural knowledge and skills, and communications and social marketing. Description of control: assessment only Duration of intervention: 3 years Length of follow-up: immediate post
<b>Study/Location</b>	Hoffman 2011 [50] United States
<b>Objective</b>	To examine the effects of a multi-component, theory-based, 2.5-year intervention on children's fruit and vegetable consumption, preferences, knowledge and BMI
<b>Methods</b>	Design: RCT Selection: multiple approaches to recruit (e.g., teacher meetings, principal support, classroom presentations); four urban public schools from the same school district Inclusion criteria: signed parental consent required for inclusion Unit of allocation: school Unit of analysis: individual Intention to treat: yes
<b>Participants</b>	Sample: 297 Intervention n=149; Control n=148 Age mean (years): Overall: 6.2 Gender (Female): Overall 49.0% Loss to follow-up: Intervention n=46, Control n=43
<b>Intervention</b>	Description of intervention: school wide (daily loud speaker announcements), classroom (instructional DVD), lunchroom (daily stickers contingent on a bite of fruit or vegetable), and family (take-home activity books) components to promote F&V consumption with emphasis on F&V in school lunch; role models delivering consistent information across multiple settings. Description of control: no intervention Duration of intervention: 2.5 years Length of follow-up: immediate post
<b>Study/Location</b>	Howe 2011 [51] United States
<b>Objective</b>	To evaluate the efficacy of a 10-month PA intervention on: (a) the prevention of excessive age-related increases in body fatness and (b) cardiovascular fitness
<b>Methods</b>	Design: RCT Selection: Black boys (8-12 years) recruited from five elementary schools using fliers Inclusion criteria: all 3 <sup>rd</sup> through 5 <sup>th</sup> grade Black boys eligible if: (a) weight <300 lbs (equipment limitation), (b) not taking medications known to affect metabolism, body composition, or fat distribution, and (c) no known CV, metabolic, or respiratory disease

	<p>or physical impairment that would limit participation in regular PA</p> <p>Unit of allocation: individual</p> <p>Unit of analysis: individual</p> <p>Intention to treat: no</p>
<b>Participants</b>	<p>Sample: 106</p> <p>Intervention n=62; Control n=44</p> <p>Age range (years): 8 to 12</p> <p>Gender: 100% boys</p> <p>Race/Ethnicity: African-American</p> <p>Loss to follow-up: NR</p>
<b>Intervention</b>	<p>Description of intervention: participants stayed at school at end of day (177+/- 8.6 days) to receive a 2-hour intervention; conducted by trained personnel with exercise-related education plus 1-2 trained classroom teachers; 30 minutes of homework time during which the boys provided with a healthy snack followed by 80 minutes of PA</p> <p>Description of control: no intervention, instructed not to change after-school routine</p> <p>Duration of intervention: 10 months</p> <p>Length of follow-up: immediate post</p>
<b>Study/Location</b>	James 2007 [52] England; Companion paper: James [53]
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	Jansen 2011 [54] Netherlands
<b>Objective</b>	To evaluate the effect of a school-based intervention program to reduce overweight and improve fitness in primary school children
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: primary schools in inner-city areas of Rotterdam; 27 schools applied</p> <p>Exclusion criteria: NR</p> <p>Unit of allocation: school</p> <p>Unit of analysis: individual</p> <p>Intention to treat: no</p>
<b>Participants</b>	<p>Sample: 1,386</p> <p>Intervention n=657; Control n=729</p> <p>Age mean (SD) (years): Intervention Grades 3-5: 7.7 (1.0), Grades 6-8: 10.8 (1.0); Control: Grades 3-5: 7.8 (1.0), Grades 6-8: 10.8 (1.0)</p> <p>Gender (Female): Grades 3-5 Intervention 50.5%; Control 51.0%; Grades 6-8 Intervention 52.8%; Control 49.0%</p> <p>Loss to follow-up: Intervention n=91; Control n=115</p>

<b>Intervention</b>	<p>Description of intervention: Lekker Fit! promoting healthy eating and active living; targeted individual behaviours, school policies and curriculum; 3 PA sessions/week by PA teacher for grades 3-8 (6-12 years), 3 classroom lessons on healthy nutrition, active living and healthy lifestyle choices adapted for each grade, administration of the Eurofit test, with measurements of height, weight and 9 fitness tests</p> <p>Description of control: usual curriculum</p> <p>Duration of intervention: 10 months</p> <p>Length of follow-up: immediate post</p>
<b>Study/Location</b>	Katz 2011 [55] United States
<b>Objective</b>	To evaluate the effects of a nutrition education program designed to teach elementary students and their parents to distinguish between more healthful and less healthful choices in diverse food categories
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: During the 2007-2008 school year, participants second to fourth grade students recruited from 5 elementary schools in Independence, Missouri</p> <p>Exclusion criteria: Students excluded from data collection and program evaluation if parental consent not received or if the student was unwilling or unable to comply</p> <p>Unit of allocation: school</p> <p>Unit of analysis: school</p> <p>Intention to treat: yes</p>
<b>Participants</b>	<p>Sample: 1,180</p> <p>Intervention n=628; Control n=552</p> <p>Age range (years): 7 to 9</p> <p>Gender (Female): Total: 51.1%; Intervention: 50.3%; Control: 52.2%</p> <p>Loss to follow-up: NR</p>
<b>Intervention</b>	<p>Description of intervention: The Nutrition Detectives program consists of 5 lessons (power point presentation plus hands on activity) presented by physical education instructors in four 20-minute sessions; a booster training session offered later in year</p> <p>Description of control: NR</p> <p>Duration of intervention: 1 school year</p> <p>Length of follow-up: 1 school year</p>
<b>Study/Location</b>	Klesges 2010 [56] United States
<b>Objective</b>	To determine the efficacy of a 2-year obesity prevention intervention in African-American girls
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: recruitment in 5 waves primarily through television and radio ads and flyers and community presentations; ads described GEMS as a study of healthy growth</p>

	<p>Inclusion criteria: identified as African-American or Black by parent/caregiver; aged 8-10 years; BMI <math>\geq</math>25th age-sex specific percentile, or at least one parent with BMI <math>\geq</math>25; Girls were excluded if they had BMI &gt;35 or conditions that would affect growth or limit participation in the study.</p> <p>Unit of allocation: individual</p> <p>Unit of analysis: individual</p> <p>Intention to treat: yes</p>
<b>Participants</b>	<p>Sample: 303</p> <p>Intervention n=153; Control n=150</p> <p>Age mean (SD) (years): Intervention: 9.3 (0.9); Control: 9.3 (0.9)</p> <p>Gender: 100% female</p> <p>Race/Ethnicity: African-American</p> <p>Loss to follow-up: 20%</p>
<b>Intervention</b>	<p>Description of intervention: girls and caregivers participated in the obesity prevention intervention through a combination of separate and joint sessions.</p> <p>Description of control: intervention on improving self-esteem and social efficacy</p> <p>Duration of intervention: 2 years</p> <p>Length of follow-up: immediate post</p>
<b>Study/Location</b>	Kriemler 2010 [57] Switzerland; Companion paper: Zahner [58]
<b>Objective</b>	To assess the effectiveness of a school based physical activity program during one school year on physical and psychological health in young schoolchildren
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: two provinces in Switzerland. Recruitment of participating schools based on willingness to be randomized either to an intervention group or a control group.</p> <p>Unit of allocation: school</p> <p>Unit of analysis: individual</p> <p>Intention to treat: yes</p>
<b>Participants</b>	<p>Sample: 502</p> <p>Intervention (grades 1 and 5 combined) n=297; Control (grades 1 and 5 combined) n=205</p> <p>Age mean (SD) (years): Intervention (grade 1): 6.9 (0.3); Intervention (grade 5): 11 (0.5); Control (grade 1): 6.9 (0.3); Control (grade 5): 11.3 (0.6)</p> <p>Gender [Female n (%): Intervention 1 n=64 (49%); Intervention 2 n=91 (55%); Control 1 n=50 (55%); Control 2 n=52 (46%)</p> <p>Loss to follow-up: NR</p>
<b>Intervention</b>	Description of intervention: children in both groups had three 45 minute PA lessons each week; intervention group had two more lessons on remaining school days; all intervention classes received same curriculum; 3-5 short activity breaks (2-5 minutes each) during academic lessons for motor skill tasks such as jumping or balancing on

	<p>one leg, power games or coordinative tasks; children received daily PA homework of about 10 minutes including aerobic, strength, or motor skill tasks such as brushing their teeth while standing on one leg, hopping up and down the stairs, rope jumping.</p> <p>Description of control: three physical education lessons each week</p> <p>Duration of intervention: 9 months</p> <p>Length of follow-up: immediate post</p>
--	---

<b>Study</b>	Lazaar 2001 [59] France
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	Li 2010 [60] China
<b>Objective</b>	To determine whether a large-scale physical activity intervention could affect body composition in primary school students in Beijing, China
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: two school districts randomly selected from eight in urban Beijing</p> <p>Inclusion criteria: NR</p> <p>Unit of allocation: school</p> <p>Unit of analysis: individual</p> <p>Intention to treat: no</p>
<b>Participants</b>	<p>Sample: 4,700</p> <p>Intervention n=2,329; Control n=2,371</p> <p>Age mean (SD) (years): Overall: 9.3 (0.7)</p> <p>Gender [Female n (%): 2,242 (47.7%)</p> <p>Loss to follow-up: Intervention n=301; Control n=279</p>
<b>Intervention</b>	<p>Description of intervention: 20 min of daily exercise in the classroom</p> <p>Description of control: no intervention in control schools</p> <p>Duration of intervention: 1 year</p> <p>Length of follow-up: immediate post</p>
<b>Study/Location</b>	Llargues 2012 [61] Spain; Companion paper: Llargues [62]
<b>Objective</b>	To assess whether the benefits seen in nutrition, physical activity and body mass index were maintained at 2 years of completion of the educational intervention
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: all children born in 2000 who attended any school in Granollers</p> <p>Exclusion criteria: school children requiring a special diet for a metabolic or digestive disorders, physical activity incapacity, no family acceptance of attendance to school</p>

	Unit of allocation: school Unit of analysis: individual Intention to treat: no
<b>Participants</b>	Sample: 464 Intervention n=233; Control n=231 Age mean (years): Intervention: 6.03; Control: 6.03 Gender [Female n (%): Intervention n=216 (48.2%); Control n=178 (47.5%) Loss to follow-up: Intervention n=9; Control n=23
<b>Intervention</b>	Description of intervention: promotion of healthy dietary habits and PA using IVAC (Intervention using research, Vision, Action and Change) educational pedagogy for two consecutive school years. The IVAC method is used in health strategies because the perceptions and knowledge elaborated by schoolchildren are directed towards change, so that they make their own decisions based on their concepts of health, determination of priorities, and change. Teachers act as moderators in conversations between schoolchildren and help them develop skills to be able to change these conditions. At study start, a group of educators specializing in community projects trained teachers in the intervention group in the above methodology Description of control: no intervention Duration of intervention: 2 years Length of follow-up: 2 years
<b>Study/Location</b>	Lloyd 2012 [63] United Kingdom
<b>Objective</b>	To assess the behavioural and weight status outcomes in English children in a feasibility study of a novel primary school-based obesity prevention program
<b>Methods</b>	Design: RCT Selection: schools recruited via the local network of primary school head teachers Inclusion criteria: all State schools in Exeter were eligible if they had at least one single age year 5 class (9-10-year-olds) (i.e., not mixed classes, 8-10- or 9-11- year-olds) Unit of allocation: school Unit of analysis: child Intention to treat: yes
<b>Participants</b>	Sample: 202 Intervention n=80; Control n=122 Age mean (SD) (years): Overall: 9.69 (0.3) Gender (Female): 50% Loss to follow-up: Intervention n=7; Control n=8
<b>Intervention</b>	Description of intervention: HeLP is a multicomponent four-phase program delivered over three school terms; program based on the Information, Motivation and Behavioural Skills Model, which proposes adequate information, motivation and

	<p>behavioural skills are essential to behaviour change; three key behaviours are emphasised: decrease in the consumption of sweetened fizzy drinks, increase in the proportion of healthy snacks to unhealthy snacks consumed and reduction in television viewing and other screen-based activities</p> <p>Description of control: no intervention</p> <p>Duration of intervention: 10 months</p> <p>Length of follow-up: 8 months, 14 months</p>
--	--

<b>Study/Location</b>	Lubans 2011 [64] Australia; Companion papers: Lubans [65], Morgan [66]
<b>Objective</b>	To evaluate the efficacy and feasibility of the Physical Activity Leaders program, an obesity prevention program for low-active adolescent boys from disadvantaged schools
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: 6 low SES co-educational secondary schools from the Hunter Region, NSW Australia were invited to participate and 4 consented. Schools were identified using the NSW DET Priority Schools Program (PSP) classification (identifies disadvantaged schools from communities with the highest concentrations of low SES families); physical education teachers were involved in identifying and recruiting low-active boys</p> <p>Inclusion criteria: adolescent boys in grade 9 attending one of the four study schools; students considered by the teachers to be disengaged in PE and/or not currently participating in organized team or individual sports</p> <p>Unit of allocation: school</p> <p>Unit of analysis: individual</p> <p>Intention to treat: NA</p>
<b>Participants</b>	<p>Sample: 100</p> <p>Intervention n=50; Control n=50</p> <p>Age mean (SD) (years): Intervention: 14.4 (0.7); Control: 14.2 (0.4)</p> <p>Gender: 100% boys</p> <p>SES: all schools had to be identified as disadvantaged schools (by PSP classification)</p> <p>Loss to follow-up: no loss</p>
<b>Intervention</b>	<p>Description of intervention: a multi-component school-based intervention including school sport sessions, interactive seminars, lunch-time activities, physical activity and nutrition handbooks, leadership sessions and pedometers for self-monitoring</p> <p>Description of control: program delivered at the wait-list control group schools at the completion of the study</p> <p>Duration of intervention: 6 months</p> <p>Length of follow-up: immediate post</p>
<b>Study/Location</b>	Lubans 2012 [67] Australia



<b>Objective</b>	To evaluate the impact of a multicomponent school-based obesity prevention program, Nutrition and Enjoyable Activity for Teen Girls
<b>Methods</b>	Design: RCT Selection: state-funded secondary schools located in New South Wales, Australia, in areas with lower SES; 18 schools invited to participate, 12 were recruited; eligible participants were adolescent girls in grade 8 (second year of secondary school) Unit of allocation: school Unit of analysis: individual Intention to treat: yes
<b>Participants</b>	Sample: 357 Intervention n=178; Control n=179 Age mean (SD) (years): Intervention: 13.15 (0.44); Control: 13.20 (0.45) Gender (Female): 100% Loss to follow-up: Intervention n=37; Control n=26
<b>Intervention</b>	Description of intervention: enhanced school sport sessions, interactive seminars, nutrition workshops, lunch-time PA sessions, handbooks and pedometers for self-monitoring, parent newsletters, text messaging for social support; school champions (i.e., teachers responsible for program delivery) attended 1-day training workshop which focused on promoting PA, reducing sedentary behaviours, and encouraging low-cost healthy eating; delivered during 4 school terms; enhanced sport sessions (60-80 minutes) delivered by teachers involved a range of activities organized into 4-week units; three practical nutrition workshops delivered by dietitians to provide students with the confidence to select, prepare, and consume healthy low-cost foods; parents sent 4 newsletters; girls sent weekly text messages during second and third terms and biweekly during fourth term (e.g., "Sitting down for long periods of time is bad for you, but what makes it worse is that people often eat junk while sitting down in front of the TV. Try to avoid eating dinner while watching TV"). Description of control: no intervention Duration of intervention: 12 months Length of follow-up: immediate post
<b>Study/Location</b>	Madsen 2013 [68] United States
<b>Objective</b>	To evaluate the impact of a community-based after-school soccer and youth development program, America SCORES, on students' physical activity, weight status, and fitness
<b>Methods</b>	Design: RCT Selection: the study was presented at a regularly scheduled principals' meeting Inclusion criteria: all 4th and 5th grade students enrolled in the after-school program at participating schools Unit of allocation: school

	Unit of analysis: child Intention to treat: yes
<b>Participants</b>	Sample: 156 Intervention n=82; Control n=74 Age mean (SD) (years): Overall: 9.8 (8.6) Gender (Female): 40% Race/Ethnicity: 12% African American; 32% Asian and 42% Latino SES (Mother's education): 56% had high school or less Loss to follow-up: Intervention n=3; Control n=3
<b>Intervention</b>	Description of intervention: soccer and writing coaches trained with a standard curriculum to lead the SCORES program in the after-school setting; students spent 2 to 3 days per week in soccer drills or games for up to 2 hours each day; the 2 non-soccer days dedicated to creative writing and performance in the 12-week fall session and to community service projects in the 12-week spring session Description of control: NR Duration of intervention: 8-10 months (1 school year) Length of follow-up: immediate post
<b>Study/Location</b>	Magnusson 2012 [69] Iceland
<b>Objective</b>	To assess the effects of a 2-year intervention program among elementary participants on body composition and cardiorespiratory fitness
<b>Methods</b>	Design: RCT Selection: three pairs of schools in city of Reykjavik were selected and matched on size; all children attending second grade were invited to participate Unit of allocation: school Unit of analysis: individual Intention to treat: no
<b>Participants</b>	Sample: 321 Intervention n=128; Control n=138 Age mean (SD) (years): Intervention: 7.3 (0.3); Control: 7.4 (0.3) Gender [Female n (%): Intervention n=65 (51%); Control n=83 (60%) Loss to follow-up: Intervention n=23; Control n=32
<b>Intervention</b>	Description of intervention: focused on increasing PA during school hours and promoting healthy dietary habits; teacher-led daily implementation of various intervention tactics, more frequent outdoor teaching, organized fieldtrips, promotion of active commute to and from school, one extra PA lesson per week (three 40-min sessions per week instead of two compulsory 40-min sessions at the control schools) and more dietary intervention aimed to have positive impact on dietary knowledge, awareness, preferences/taste, self-efficacy and parental influence; nutrition education

	<p>material was implemented during the latter intervention year; main focus of the dietary intervention was on fruit and vegetable intake</p> <p>Description of control: no intervention</p> <p>Duration of intervention: 2 years</p> <p>Length of follow-up: immediate post</p>
<b>Study</b>	Marcus 2009 [70] Sweden
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	Martínez-Vizcaino 2008 [71] Spain
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	Mihas 2010 [72] Greece
<b>Objective</b>	To assess short-term and long-term effects of a school-based health and nutrition education intervention on diet, nutrition intake and BMI
<b>Methods</b>	<p>Design: RCT</p> <p>Exclusion criteria: participants with an organic cause for high or low weight, who had received any medication that might interfere with growth or weight control, or who were on specific diets</p> <p>Unit of allocation: individual</p> <p>Unit of analysis: individual</p> <p>Intention to treat: no</p>
<b>Participants</b>	<p>Sample: 213</p> <p>Intervention n=108; Control n=105</p> <p>Age mean (SD) (years): Intervention: 13.1 (0.8); Control: 13.3 (0.9)</p> <p>Gender [Female n (%): Intervention n=50 (51.0%); Control n=43 (50.5%)</p> <p>Loss to follow-up: Intervention n=10; Control n=12</p>
<b>Intervention</b>	<p>Description of intervention: multi-component workbooks covering mainly dietary issues, but also dental health and consumption attitudes; health and nutrition components conducted by home economics teacher supervised by a health visitor or family doctor and incorporated 12 hours of classroom material during 12 weeks; modules designed to develop behavioural capability, expectations and self-efficacy for healthful eating and healthy foods selection; learning activities designed to influence expectancies that value achieving these behaviours; parental involvement included two meetings where they were given a file containing their child's screening results</p> <p>Description of control: no intervention</p> <p>Duration of intervention: 12 weeks</p> <p>Length of follow-up: immediate post</p>

<b>Study/Location</b>	Morgan 2011 [73] Australia
<b>Objective</b>	To evaluate the feasibility and efficacy of the Healthy Dads, Healthy Kids program, which was designed to help overweight fathers lose weight and be a role model of positive health behaviours for their children
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: Overweight or obese men with a child between 5 and 12 years of age were recruited through media releases, school newsletters and paid advertisements in local newspapers in; men were screened for eligibility through telephone interviews.</p> <p>Exclusion criteria: history of major medical problems (e.g., heart disease) in last 5 years, diabetes, orthopedic or joint problems that would be a barrier to PA, recent weight loss <math>\geq 4.5</math> kg, medication use that might affect body weight; a child with extreme obesity</p> <p>Unit of allocation: individual</p> <p>Unit of analysis: individual</p> <p>Intention to treat: yes</p>
<b>Participants</b>	<p>Sample: 53 dads, 71 children</p> <p>Intervention n=27; Control n=26</p> <p>Age mean (SD) (years): Intervention: 8.4 (2.1); Control: 7.9 (1.9)</p> <p>Gender [Female n (%): Intervention 48.7%; Control: 43.7%</p> <p>Loss to follow-up at 3 months: Intervention n=6; Control n=3</p> <p>Loss to follow-up at 6 months: Intervention n=7, Control n=2</p>
<b>Intervention</b>	<p>Description of intervention: fathers attended 8 face-to-face group sessions (75 min each); 5 sessions for fathers only, delivered by male researcher; 3 sessions practical and involved both fathers and children, delivered by two male researchers, both with expertise in physical education; total contact time was 600 minutes; PA sessions for fathers emphasized modeling, reinforcing and providing opportunities and removing barriers to PA; father/child PA sessions were i) fundamental movement skills ii) rough and tumble play iii) health related fitness and iv) fun and active games; dietician developed nutrition components modeled on a previous successful intervention; healthy eating focused on parental influence on children's dietary intake, incorporating Satter's 'trust' paradigm, which suggests parents should supply healthy foods and a supportive eating environment and children can decide when and how much to eat</p> <p>Description of control: waitlist</p> <p>Duration of intervention: 3 months</p> <p>Length of follow-up: 3 and 6 months</p>
<b>Study/Location</b>	Mo-suwan 1998 [74] Thailand
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	Muckelbauer 2012 [75] Germany; Companion paper: Muckelbauer [76]

<b>Objective</b>	To test whether a simple overweight prevention program promoting water consumption in elementary schools is equally effective in children with an immigration background and in those without
<b>Methods</b>	Design: RCT Selection: schools eligible if located in deprived areas, as defined by: unemployment rate $\geq 15\%$ and proportion of social welfare recipients $\geq 5\%$ , and proportion of non-German residents $\geq 5\%$ as indicated by local public authorities Unit of allocation: school Unit of analysis: individual Intention to treat: no
<b>Participants</b>	Sample: 3,190 Intervention n=1,641; Control n=1,309 Age mean (SD) (years): Intervention: 8.26 (0.73); Control: 8.34 (0.76) Gender [Female n (%): Intervention n=817 (49.8%); Control n= 651 (49.7%) Loss to follow-up: Intervention n=65; Control n=60
<b>Intervention</b>	Description of intervention: in each school, 1 water fountain, or 2 for schools with >150 participants, was installed; each child received a plastic water bottle (500 mL), and teachers encouraged to organize filling of bottles each morning; four 45-minute classroom lessons dealing with the body's water needs and the water circuit in nature; teachers received curriculum and materials to implement the lessons; 3 months into the study, teachers introduced a motivation unit (booster sessions) that used a goal-setting strategy to reach a sustained increase in water consumption by giving quantitative targets and feedback; 5 months after baseline, each participant received a new water bottle with an improved handling design Description of control: no intervention Duration of intervention: 10 months (1 school year) Length of follow-up: immediate post
<b>Study/Location</b>	Nemet 2011a [77] Israel
<b>Objective</b>	To prospectively examine the effects of a randomized school-based intervention on nutrition and physical activity knowledge and preferences, anthropometric measures, and fitness in low SES kindergarten children
<b>Methods</b>	Design: RCT Selection: 30 kindergartens from low SES communities Unit of allocation: classes Unit of analysis: individual Intention to treat: no
<b>Participants</b>	Sample: 795 Intervention n=417; Control n=378

	Age mean (SD) (years): Intervention: 5.20 (0.02); Control 5.24 (0.03) Gender (Female): Intervention 46%; Control 44% Loss to follow-up: NR
<b>Intervention</b>	Description of intervention: designed to improve nutritional knowledge, based on the nutritional program "It Fits Me" ("Tafur Alay") of the Israeli Ministry of Education; teaching topics included food groups, vitamins, healthy food choices, food preparation and cooking methods, and information on fast-food versus home cooking; topics taught through short lectures/talks, games and story reading; children participated in 45 minutes (three 15-minute sessions) per day of exercise training (6 days a week) Description of control: NR Duration of intervention: 1 school year Length of follow-up: 1 school year
<b>Study/Location</b>	Nemet 2011b [78] Israel
<b>Objective</b>	To examine the prevalence of obesity and to prospectively study the effects of a health promotion, school-based intervention on nutrition and physical activity knowledge and preferences, anthropometric measures, and fitness in Arab-Israeli kindergarten children
<b>Methods</b>	Design: RCT Selection: kindergarten classes randomly assigned by computerized program to participate in intervention or to serve as controls (6 control, 5 intervention) Exclusion criteria: students not coming from low SES communities Unit of allocation: school Unit of analysis: child Intention to treat: no
<b>Participants</b>	Sample: 342 Intervention n=154; Control n=188 Age mean (SD) (years): Intervention: 5.36 (0.03); Control: 5.40 (0.02) Gender (Female): 45% Race/Ethnicity: predominantly Arab-Israeli SES: schools were selected from low SES Loss to follow-up: Intervention n=20; Control n=25
<b>Intervention</b>	Description of intervention: preschool teachers attended an all-day training session that covered nutrition and physical activity; 2 additional days held to collect feedback on the program and introduce new materials; parents and children were invited to 2 Health Festival days that focused on the major themes of the program (introduction of healthy nutrition, prevention of childhood obesity and beneficial effects of exercise in children) Description of control: no intervention Duration of intervention: 1 school year Length of follow-up: immediate post

<b>Study/Location</b>	Neumark-Sztainer 2003 [79] United States
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	Neumark-Sztainer 2010 [80] United States
<b>Objective</b>	To evaluate a school-based program aimed at preventing weight-related problems in adolescent girls
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: girls in intervention and control schools invited to register for an all-girls physical education class as an alternative to the regular coeducational class; in participating schools, students were required to take one or two physical education classes to graduate; participation in the study class counted toward that requirement; recruitment materials designed to appeal to inactive girls interested in healthy weight management; class description included in the school catalogue; posters and flyers about the program were displayed at schools</p> <p>Exclusion criteria: high physical activity levels (<math>\geq 1</math> hour/day) and eating disorder behaviours (vomiting or laxative use weekly or more)</p> <p>Unit of allocation: school</p> <p>Unit of analysis: individual</p> <p>Intention to treat: no</p>
<b>Participants</b>	<p>Sample: 356</p> <p>Intervention n=182; Control n=174</p> <p>Age mean (SD) (years): Intervention: 15.7 (1.13); Control: 15.8 (1.22)</p> <p>Gender (Female): 100%</p> <p>Loss to follow-up: Intervention n=5; Control n=15</p>
<b>Intervention</b>	<p>Description of intervention: New Moves included (1) 16 week physical education class (Be Fit 4 days/week) which also incorporated nutrition (Be Fueled) and social support/self-empowerment (Be Fab) sessions 1 day/week; (2) individual counseling sessions using motivation interviewing techniques; (3) lunch get-togethers (lunch bunches) 1/week during maintenance period; (4) minimal parent outreach activities</p> <p>Description of control: all girls physical education class</p> <p>Duration of intervention: 9 months</p> <p>Length of follow-up: immediate post</p>
<b>Study/Location</b>	Ostbye 2012 [81] United States; Companion paper: Ostbye [82]
<b>Objective</b>	To evaluate the effects of Kids and Adults Now - Defeat Obesity! on enhancing healthy lifestyle behaviours in mother-preschooler (2-5 years old) dyads
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: mothers primarily identified from state birth certificates and screened for</p>

	<p>eligibility at 2-6 months postpartum</p> <p>Inclusion criteria: eligible mothers had a preschooler aged 2-5 years, self-reported pre-pregnancy (and measured postpartum) BMI <math>\geq 25</math>, no medical conditions preventing daily physical activity, English literacy, regular telephone access, <math>\geq 18</math> years of age</p> <p>Unit of allocation: dyads</p> <p>Unit of analysis: child</p> <p>Intention to treat: no</p>
<b>Participants</b>	<p>Sample: 400 mother-child dyads</p> <p>Intervention n=200; Control n=200</p> <p>Age mean (SD) (years): 3.06 (1.0)</p> <p>Gender (Female): Intervention 43.5%; Control 45%</p> <p>Loss to follow-up: Intervention n=50; Control n=49</p>
<b>Intervention</b>	<p>Description of intervention: 8 monthly mailed interactive kits, followed each month by a 20-30 minute telephone coaching session using motivational interviewing techniques; kits included child activities and incentives reinforcing the month's topic (e.g. a rewards chart, yoga mat, pedometer, portion plate)</p> <p>Description of control: monthly newsletters emphasizing pre-reading skills; retention encouraged by monetary incentives (up to \$100 for completing all assessments)</p> <p>Duration of intervention: 12 months</p> <p>Length of follow-up: immediate post</p>
<b>Study/Location</b>	Paineau 2008 [83] France
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	Papadaki 2010 [84] Netherlands, Denmark, United Kingdom, Greece, Germany, Spain, Bulgaria and Czech Republic; Companion paper: Larsen [85]
<b>Objective</b>	To investigate the effect of protein and glycemic index on body composition among European children in the DiOGenes (diet, obesity, and genes) family-based study
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: volunteer families from 8 countries (Netherlands, Denmark, United Kingdom, Greece, Germany, Spain, Bulgaria, and Czech Republic; families attended a screening examination to determine eligibility [eligible families were generally healthy, with at least 1 parent overweight (BMI<math>&lt;27</math>) and younger than 65 years, and at least 1 child between the age of 5 and 18 years]</p> <p>Exclusion criteria (for children): special diets, food intolerances, systemic infections or chronic diseases, use of medications that might influence study outcomes, drug or alcohol abuse</p> <p>Unit of allocation: family</p> <p>Unit of analysis: children</p> <p>Intention to treat: no</p>



<b>Participants</b>	<p>Sample: 465</p> <p>Intervention 1 (LP/LGI) n=102; Intervention 2 (LP/HGI) n=87; Intervention 3 (HP/LGI) n=92; Intervention 4 (HP/HGI) n=96; Control n=88</p> <p>Age mean (SD) (years): Overall males 11.9 (3.4); Overall females 12.4 (3.5)</p> <p>Gender (Female): 76%</p> <p>Loss to follow-up: 48%</p>
<b>Intervention</b>	<p>Description of intervention: trained dietician gave instructions on ad libitum diets; all diets were low in fat (25-30% of energy); target was for protein content to comprise 10-15% of energy intake in the low protein (LP) and 23-28% in the high protein (HP) groups, complying with the acceptable range (10-30%) for children aged 4 to 18 years; children in the low glycemic index (LGI) groups were advised to consume the LGI foods, and those in the high glycemic index (HGI) groups to consume the HGI foods</p> <p>Description of control: diet followed national dietary guidelines, with medium protein content and no specific instructions on glycemic index</p> <p>Duration of intervention: 6 months</p> <p>Length of follow-up: immediate post</p>
<b>Study/Location</b>	Peralta 2009 [86] Australia
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	Reed 2008 [87] Canada; Companion papers: Naylor [88,89]
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	Reilly 2006 [90] Scotland
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	Robinson 2003 [91] United States; Companion paper: Rochon [92]
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	Robinson 2010 [93] United States
<b>Objective</b>	To test a 2-year community- and family-based obesity prevention program for low-income African American girls: Stanford GEMS
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: recruited from schools, community centers, churches, and community events in low-income, predominantly African American neighbourhoods; identified as African American or black by parent/guardian; aged 8 to 10 years; to select a community-based group at higher risk, girls required to have BMI <math>\geq</math>25th percentile for age and/or at least 1 overweight parent/guardian (BMI <math>\geq</math>25)</p>

	<p>Exclusion criteria: girls with BMI &gt;35; diagnosed with medical condition or taking medications affecting growth; condition limiting participation in the interventions or assessments; unable to understand or complete the informed consent document; planned to move from the area; homeless; had no television</p> <p>Unit of allocation: families/households</p> <p>Unit of analysis: individual</p> <p>Intention to treat: yes</p>
<b>Participants</b>	<p>Sample: 284</p> <p>Intervention n=134; Control n=127</p> <p>Age mean (years): Intervention: 9.5; Control: 9.4</p> <p>Gender (Female): 100%</p> <p>Loss to follow-up: Intervention n=32; Control n=27</p>
<b>Intervention</b>	<p>Description of intervention: GEMS Jewels after-school dance intervention offered 5 days per week, 12 months per year (excluding school holidays), at community centers in selected neighborhoods; daily sessions lasted up to 2.5 hours and started with a 1-hour homework period and small snack followed by 45 to 60 minutes of learning and practicing dance routines; dance classes led by female African American college students and/or recent graduates from the local community</p> <p>Description of control: active-placebo health education comparison intervention consisting of culturally tailored, information-based health education on nutrition, physical activity, and reducing cardiovascular and cancer risk; 24 monthly newsletters for the girls and their parents/guardians and quarterly community center health lectures</p> <p>Duration of intervention: 2 years</p> <p>Length of follow-up: 6 months</p>
<b>Study/Location</b>	Rosario 2013 [94] Portugal; Companion paper: Rosario [95]
<b>Objective</b>	To examine the effects of a program run by teachers trained in nutrition, on consumption of low nutrient, energy-dense foods, by children attending elementary schools
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: 7 out of 80 public elementary schools from a city from the north of Portugal randomly selected and invited to participate</p> <p>Unit of allocation: school</p> <p>Unit of analysis: child</p> <p>Intention to treat: no</p>
<b>Participants</b>	<p>Sample: 464</p> <p>Intervention n=233; Control n=231</p> <p>Age mean (SD) (years): Intervention: 8.3 (1.2); Control: 8.2 (1.2)</p> <p>Gender (Female): Intervention 50.2%; Control 52.8%</p> <p>SES (mother's education up to 9 years): Intervention n=116 (58.6%); Control n=128</p>

	(69.9%); SES (father's education up to 9 years): Intervention n=122 (62.9%); Control n=132 (75.9%) Loss to follow-up: Intervention n=82; Control n=88
<b>Intervention</b>	Description of intervention: teachers attended 12 sessions on: health promotion and overweight/obesity prevention; food and nutrition and dietary guidelines (Portuguese Food Wheel); hydration and the importance of water; appropriate physical activity levels and healthy eating practices; teaching and learning strategies on healthy eating in the classroom; strategies to reduce screen time; healthy cooking and strategies to get children and families involved in healthy cooking; teachers delivered content to students and developed creative and engaging classroom activities about the topics Description of control: NR Duration of intervention: 24 months Length of follow-up: immediate post
<b>Study/Location</b>	Rosenkranz 2010 [96] United States
<b>Objective</b>	To evaluate the effectiveness of an intervention delivered through Girl Scout Junior troops designed to foster healthful troop meeting environments and increase obesity prevention behaviours at home
<b>Methods</b>	Design: RCT Selection: registered Girl Scout Junior troops, with girls in 4th and 5th grades; troops meet at least twice/month in facilities allowing physical activity and food preparation Exclusion criteria: troops not primarily composed of Girl Scout Juniors, not regularly meeting during the study period, or not having leader and parental consent for troop participation; individual girls were excluded if they could not speak or read English Unit of allocation: troops Unit of analysis: individual Intention to treat: no
<b>Participants</b>	Sample: 76 Intervention n=34; Control n=42 Age mean (SD) years: Intervention: 10.6 (1.1); Control: 10.5 (1.3) Gender (Female): 100% Race/Ethnicity: Intervention: Caucasian: 79.4%, Racial minority: 20.6%; Control: Caucasian: 75%, Racial minority: 25% Loss to follow-up: Intervention n=1; Control n=3
<b>Intervention</b>	Description of intervention: three main components: 1) an interactive educational curriculum delivered by troop leaders (8 modules, 60 to 90 minutes each, delivered over 4 months); 2) troop meeting policies implemented by troop leaders; and 3) badge assignments completed at home by girls with parental assistance Description of control: standard care

	Duration of intervention: 7 months Length of follow-up: immediate post
<b>Study/Location</b>	Rush 2012 [97] New Zealand; Companion papers: Graham [98], Cole [99]
<b>Objective</b>	To compare changes in blood pressure and body composition in children who attended Energize schools with children in control schools
<b>Methods</b>	Design: RCT Selection: NR Inclusion criteria: NR Unit of allocation: school Unit of analysis: children Intention to treat: N/A
<b>Participants</b>	Sample: 1,352 Intervention 1 n=492; Intervention 2 n=200; Control 1 n=434; Control 2 n=226 Age range (years): Intervention 1: 5-7; Intervention 2: 10-12; Control 1: 5-7; Control 2: 10-12 Gender (Female): Intervention 1 n=51%; Intervention 2 n=51%; Control 1 n=51%; Control 2 n=50% Race/Ethnicity: Intervention 1: European: 67%, Maori: 23%, Other: 9%; Intervention 2: European: 60%, Maori: 33%, Other: 7%; Control 1: European: 67%, Maori: 26%, Other: 7%; Control 2: European: 68%, Maori: 25%, Other: 7% Loss to follow-up: NR
<b>Intervention</b>	Description of intervention: program staff received training as a group in order to share experience, resources and skills; classes included fundamental movement skill training, ideas for 'huff and puff' fitness activities, modified games, and ball activities and sport-related games; teachers provided with ideas for managing children during physical activity sessions; program staff promoted active transport, lunchtime games, bike days and leadership training for students to be leaders of physical activities before and after school; program staff available to assist schools with healthy-eating initiatives Description of control: no intervention Duration of intervention: 2 years Length of follow-up: immediate post
<b>Study/Location</b>	Salcedo 2010 [100] Spain; Companion paper: Martínez-Vizcaíno [71]
<b>Objective</b>	To assess the impact of a 2-year recreational physical activity program in 1,044 fourth- and fifth-grade primary schoolchildren
<b>Methods</b>	Design: RCT Selection: 20 public schools in 20 towns in Cuenca Province Exclusion criteria: schools outside of Cuenca province

	<p>Unit of allocation: school</p> <p>Unit of analysis: individual</p> <p>Intention to treat: no</p>
<b>Participants</b>	<p>Sample: 1,119</p> <p>Intervention n=513; Control n=606</p> <p>Age mean (SD) (years): Intervention: 10.6 (1.1); Control: 10.5 (1.3)</p> <p>Gender [Female n (%): Intervention n=231 (45%); Control n=289 (48%)</p> <p>Loss to follow-up: Intervention n=138; Control n=60</p>
<b>Intervention</b>	<p>Description of intervention: MOVI was a non-competitive and recreational physical activity program consisting of three 90-minute sessions per week, during approximately 28 weeks every year; physical activity sessions were planned by 2 qualified physical education teachers and were supervised by sports instructors; standard physical education curriculum (3 hours per week of physical activity at low to moderate intensity) was also provided in intervention schools</p> <p>Description of control: standard physical education curriculum</p> <p>Duration of intervention: 7 months</p> <p>Length of follow-up: immediate post</p>
<b>Study/Location</b>	Shamah 2012 [101] Mexico
<b>Objective</b>	To assess the effectiveness of a nutrition and physical activity strategy, called “Nutrition on the Go” in maintaining the BMI values of school children in Mexico
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: 60 schools selected at random</p> <p>Exclusion criteria: schools outside of the State of Mexico</p> <p>Unit of allocation: class</p> <p>Unit of analysis: children</p> <p>Intention to treat: no</p>
<b>Participants</b>	<p>Sample: 1,019</p> <p>Intervention n=509; Control n=510</p> <p>Age mean (years): Intervention: 10; Control: 10</p> <p>Gender [Female n (%): Intervention n=263 (51.6 %); Control n=253 (49.7%)</p> <p>Loss to follow-up: Intervention n=13; Control n=12</p>
<b>Intervention</b>	<p>Description of intervention: nutrition and physical activity workshops; sale of fruit and vegetables and water in the school store; organized physical activity twice a week; banners; recipe calendar</p> <p>Description of control: no intervention</p> <p>Duration of intervention: 6 months</p> <p>Length of follow-up: immediate post</p>

<b>Study/Location</b>	Sichieri 2009 [102] Brazil
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	Siegrist 2013 [103] Germany
<b>Objective</b>	To investigate the effects of a school-based prevention program on physical activity, fitness, and obesity
<b>Methods</b>	Design: RCT Selection: 60 primary schools in Bavaria, Germany were invited by mail or telephone Inclusion criteria: attendance in 2 <sup>nd</sup> or 3 <sup>rd</sup> grade and written consent from parents Unit of allocation: school Unit of analysis: child Intention to treat: no
<b>Participants</b>	Sample: 902 Intervention n=486; Control n=340 Age mean (SD) (years): Overall 8.4 (0.7) Gender [Female n (%): n=350 (48.3%) Loss to follow-up: Intervention n=59; Control n=43
<b>Intervention</b>	Description of intervention: educating and encouraging children, teachers and parents to live active and healthy lifestyles; monthly lessons lasting 45 minutes with three parts: 10 minute warm-up of high intensity running games, 30 min of exercises to improve body awareness and self-esteem with conversation about health-related topics, and 5 min relaxation exercises; worksheets and homework assignments plus monthly newsletters to stimulate parent-child interaction and support physical activity at home and in sports clubs; school environment altered to promote more physical activity; 2 parent training sessions about health issues; teacher trainings to increase students' physical activity during lessons and breaks Description of control: usual physical education curriculum Duration of intervention: 12 months Length of follow-up: immediate post
<b>Study/Location</b>	Simon 2008 [104] France; Companion papers: Simon [105,106]
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	Singh 2009 [107] Netherlands
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	Singhal 2010 [108] India

<b>Objective</b>	To study the effectiveness of a multi-component intervention for nutrition and lifestyle education on behaviour, anthropometry and metabolic risk profile in urban adolescents
<b>Methods</b>	Design: RCT Selection: NR Inclusion criteria: NR Unit of allocation: school Unit of analysis: individual Intention to treat: no
<b>Participants</b>	Sample: 209 Intervention n=101; Control n=108 Age mean (SD) (years): Intervention: 16.04 (0.41); Control: 16.0 (0.5) Gender (Female %): Intervention 38.6%; Control 41.7% Loss to follow-up: Intervention n=2; Control n=6
<b>Intervention</b>	Description of intervention: multi-component model including seven components of nutrition and lifestyle education aimed at changing knowledge, behaviour and risk profile of urban Asian Indian adolescents Description of control: no intervention Duration of intervention: 6 months Length of follow-up: immediate post
<b>Study/Location</b>	Story 2003 [109] United States; Companion papers: Rochon [92], Story [6]
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	Story 2012 [110] United States
<b>Objective</b>	To develop and test the effectiveness of a school environment intervention, supplemented with family involvement, to reduce excessive weight gain by increasing physical activity and healthy eating practices among kindergarten and first-grade American Indian children
<b>Methods</b>	Design: RCT Selection: NR Exclusion criteria: NR Unit of allocation: school Unit of analysis: child Intention to treat: yes
<b>Participants</b>	Sample: 454 Intervention n=267; Control n=187 Age mean (SD) (years): Intervention: 5.87 (0.54); Control: 5.80 (0.51)

	<p>Gender (Female): 49%</p> <p>Race/Ethnicity: Native American</p> <p>Loss to follow-up: NR</p>
<b>Intervention</b>	<p>Description of intervention: at least 60 min of physical activity at school each day using school PE, class walks outdoors, in-class action breaks, and active recess; healthy eating promoted through offering 1% white milk instead of 2%, whole, chocolate or other flavoured milks, serving recommended portions, purchasing and using low-calorie/fat foods, offering low-fat portion-controlled salad dressing, providing more fruits and vegetables, offering second helpings only on fruits and vegetables, teachers trained to limit daily snacks; modify home environment to reduce excessive caloric intake, reduce television watching, and increase physical activity; 4 family events related to nutrition and physical activity held at the schools</p> <p>Description of control: no intervention</p> <p>Duration of intervention: 14 weeks or 31 weeks</p> <p>Length of follow-up: immediate post</p>
<b>Study/Location</b>	Telford 2012 [111] Australia; Companion paper: Telford [112]
<b>Objective</b>	To determine whether physical education taught by specialists contributed to academic development and prevention of obesity in elementary school children
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: government-funded schools in outer-city suburbs of similar average family income from an Australian education jurisdiction through invitations to principals; of 30 invited, 29 schools accepted; 13 schools (32 classes) randomly assigned to specialist-taught PE group and 16 schools (36 classes) to common-practice PE group</p> <p>Inclusion criteria: NR</p> <p>Unit of allocation: school</p> <p>Unit of analysis: child</p> <p>Intention to treat: no</p>
<b>Participants</b>	<p>Sample: 620</p> <p>Intervention n=312; Control n=308</p> <p>Age: NR</p> <p>Gender [Female n (%): Intervention n=154 (49%); Control n=149 (48%)</p> <p>Race/Ethnicity: White: 86%, Asian: 8%, Australian Aboriginal or Torres Strait Islander: 3%, Polynesian: 1%, Data missing: 2%</p> <p>Loss to follow-up: NR</p>
<b>Intervention</b>	<p>Description of intervention: students received 150 minutes per week of PE; specialist-taught PE included 90 minutes per week of PE from visiting specialists</p> <p>Description of control: common practice (PE from generalist classroom teachers)</p> <p>Duration of intervention: 2 years</p>



	Length of follow-up: immediate post
<b>Study/Location</b>	Thivel 2011 [113] France
<b>Objective</b>	To assess the effect of a 6-month physical activity program on body composition and physical fitness among primary school children
<b>Methods</b>	Design: RCT Selection: primary school children recruited from local public schools that agreed to participate in the study Inclusion criteria: attendance in 1 <sup>st</sup> or 2 <sup>nd</sup> grade, taking part in standard physical education classes, participating in no more than 3 hours of extracurricular sports activity per week, free of any known disease, not involved in any other study Unit of allocation: school Unit of analysis: individual Intention to treat: N/A
<b>Participants</b>	Sample: 457 Intervention n=229; Control n=228 Age: NR (1 <sup>st</sup> and 2 <sup>nd</sup> grade) Gender [Female n (%): Intervention n=117 (51%); Control n=112 (49%) Loss to follow-up: NR
<b>Intervention</b>	Description of intervention: 120 min (two times for 60 min) of supervised physical activity; 2 additional hours of physical education classes per week managed and taught by sports science students; sessions consisted of a 10-min warm-up followed by psychometric activities and exercises to improve coordination, flexibility, strength, speed, and endurance Description of control: regular 2 hours of physical education per week Duration of intervention: 6 months Length of follow-up: immediate post
<b>Study/Location</b>	Velez 2010 [114] United States
<b>Objective</b>	To examine the effects of a structured resistance training program on strength, body composition, and self-concept in normal and overweight Hispanic adolescents
<b>Methods</b>	Design: RCT Selection: recruited from a predominantly Hispanic high school in central New Jersey; Hispanic youth selected because of this population's greater propensity for obesity and their underrepresentation in resistance training research Exclusion criteria: known health (i.e., bone, joint, musculoskeletal, or cardiovascular) problems that would severely limit involvement in the resistance training sessions; already participating in structured resistance or aerobic training programs Unit of allocation: individual

	Unit of analysis: individual Intention to treat: no
<b>Participants</b>	Sample: 28 Intervention n=13; Control n=15 Age mean (SD) years: Overall: 16.14 (0.19) Gender [Female n (%): Intervention n=5 (38%); Control n=7 (47%) Loss to follow-up: 3
<b>Intervention</b>	Description of intervention: resistance training consisting of 35-40-minute sessions, 3 non-consecutive days/week, in lieu of PE class; workouts divided into upper body and lower body days; trainers met 3-4 students at a time at the school weight room and led them through planned workouts; instructed to maintain usual outside activities and diets Description of control: typical daily physical education/health class; total activity time per day was similar to intervention participants Duration of intervention: 12 weeks Length of follow-up: immediate post
<b>Study/Location</b>	Webber 2008 [115] United States
<b>Comments</b>	See Cochrane Review by Waters et al.[2] for details
<b>Study/Location</b>	Weeks 2012 [116] Australia
<b>Objective</b>	To determine the effect of a twice-weekly, school-based 10-minute jumping regime on muscle and fat tissue in healthy adolescent boys and girls
<b>Methods</b>	Design: RCT Selection: adolescents in the 9 <sup>th</sup> grade of a local high school were recruited Inclusion criteria: sound general health, fully ambulatory and had written consent of a parent or guardian Exclusion criteria: endocrine disorder, metabolic disease or chronic renal pathology, taking medication known to affect the musculoskeletal system, recovering from lower limb injury or affected by any condition not compatible with intense physical activity Unit of allocation: individual Unit of analysis: individual Intention to treat: yes
<b>Participants</b>	Sample: 99 Intervention n=52; Control n=47 Age mean (SD) years: Overall boys 13.8 (0.4), Overall girls 13.7 (0.4) Gender (Female):54% Loss to follow-up: Intervention: 9, Control: 9

<b>Intervention</b>	Description of intervention: 10 minutes of supervised jumping activity at the start of each physical education class, 2 times per week for 8 months Description of control: regular PE warm-ups and stretching Duration of intervention: 1 school year Length of follow-up: 1 school year
<b>Study/Location</b>	Wen 2012 [117] Australia
<b>Objective</b>	To assess the effectiveness of a home based early intervention on BMI at age 2
<b>Methods</b>	Design: RCT Selection: research assistants gave pregnant women attending antenatal clinics a letter of invitation and information about the study Inclusion criteria: women were eligible if aged $\geq 16$ , expecting first child, between weeks 24-34 of pregnancy, able to communicate in English, lived in the local area Unit of allocation: mother Unit of analysis: child Intention to treat: yes
<b>Participants</b>	Sample: 667 Intervention n=337; Control n=330 Age range (years): Overall 0 to 2 Gender: NR Loss to follow-up: Intervention n=88; Control n=96
<b>Intervention</b>	Description of intervention: 4 community nurses recruited and trained to make 8 home visits, once at 30-36 weeks' gestation and 7 times after the birth (at 1, 3, 5, 9, 12, 18 and 24 months); at each visit, the nurse spent about one to two hours with the mother and infant and teaching specific skills and knowledge in relation to healthy infant feeding practices and active play and discussing family physical activity, nutrition, and social support as well as any issues and concerns raised by the mother Description of control: usual childhood nursing service from community health service nurses (at least one nurse visit for general support at home; some vulnerable families are offered multiple home visits) Duration of intervention: 24 months Length of follow-up: immediate post
<b>Study/Location</b>	Williamson 2012 [118] United States; Companion paper: Williamson [119]
<b>Objective</b>	To test the efficacy of two-school based programs for prevention of body weight/fat gain in all participants and in overweight children
<b>Methods</b>	Design: RCT Selection: students recruited through presentations, fliers and word of mouth

	<p>Inclusion criteria: for schools: one of the 28 schools or elementary feeder schools in the LA GEAR UP program, located in a rural section of Louisiana, minimum of 100 students available for study; for students: in grades 4 to 6</p> <p>Unit of allocation: school</p> <p>Unit of analysis: child</p> <p>Intention to treat: no</p>
<b>Participants</b>	<p>Sample: 2,097</p> <p>Intervention 1 n=713; Intervention 2 n=760; Control n=587</p> <p>Age mean (SD) years: Overall 10.5 (1.2)</p> <p>Gender (Female): 45%</p> <p>Loss to follow-up: Intervention 1 n=224; Intervention 2 n=207, Control n=196</p>
<b>Intervention</b>	<p>Description of intervention: Intervention 1: emphasis on modification of environmental cues, enhancement of social support, and promotion of self-efficacy for health behaviour change; goals compatible with conventional nutrition recommendations; promotion of 60 minutes of moderate to vigorous activity per day; meeting USDA guidelines for the National School Lunch Program and legislated requirements related to advertising fast foods and contents of vending machines and concessions in schools</p> <p>Intervention 2: emphasis on behaviour modification approaches designed to change personal factors (i.e., increased healthy eating habits, increased physical activity, and decreased sedentary behaviour); used internet-based HIPTeens program as a part of regular classroom instruction, combined with synchronous (on-line) internet counseling and asynchronous (email) communications for children and their parents; frequent prompts to promote sustained website usage</p> <p>Description of control: none of the prevention components hypothesized to yield weight gain prevention; a nonspecific control condition</p> <p>Duration of intervention: 28 months</p> <p>Length of follow-up: immediate post</p>
<b>Study/Location</b>	Yin 2012 [120] United States
<b>Objective</b>	To determine the effects of a 3-year after-school physical activity program, without restriction of dietary energy intake, on cardiometabolic outcomes
<b>Methods</b>	<p>Design: RCT</p> <p>Selection: recruitment of children in 2<sup>nd</sup> and 3<sup>rd</sup> grades</p> <p>Exclusion criteria: NR</p> <p>Unit of allocation: school</p> <p>Unit of analysis: child</p> <p>Intention to treat: yes</p>
<b>Participants</b>	<p>Sample: 617</p> <p>Intervention n=324; Control n=293</p>

	<p>Age mean (SD) years: Overall 8.7 (0.5)</p> <p>Gender (Female): 53%</p> <p>Loss to follow-up: Intervention n=129; Control n=88</p>
<b>Intervention</b>	<p>Description of intervention: 120 min structured after-school program consisting of 40 min snack and teacher-assisted homework; 20 min skill based PA; 40 min vigorous PA; 20 min stretching/cool down; weekly health-related lesson</p> <p>Description of control: NR</p> <p>Duration of intervention: 33 months</p> <p>Length of follow-up: immediate post</p>

## References

1. Amaro S, Viggiano A, Di Costanzo A, Madeo I, Viggiano A, Baccari ME, et al. Kalédo, a new educational board-game, gives nutritional rudiments and encourages healthy eating in children: a pilot cluster randomized trial. *Eur J Pediatr* 2006 Sep;165(9):630-5.
2. Waters E, de Silva-Sanigorski A, Hall BJ, Brown T, Campbell KJ, Gao Y, et al. Interventions for preventing obesity in children. *Cochrane Database Syst Rev* 2011;(12):CD001871.
3. Baranowski T, Baranowski JC, Cullen KW, Thompson DI, Nicklas T, Zakeri IE, et al. The Fun, Food, and Fitness Project (FFFP): the Baylor GEMS pilot study. *Ethn Dis* 2003;13(Suppl 1):S30-9.
4. Barkin S, Gesell S, Po'e E, Escarfuller J, Tempesti T. Culturally tailored, family-centered, behavioral obesity intervention for Latino-American preschool-aged children. *Pediatrics* 2012 Sep;130(3):445-56.
5. Beech BM, Klesges RC, Kumanyika SK, Murray DM, Klesges L, McClanahan B, et al. Child- and parent-targeted interventions: the Memphis GEMS pilot study. *Ethn Dis* 2003;13(Suppl 1):S40-53.
6. Story M, Sherwood NE, Obarzanek E, Beech BM, Baranowski JC, Thompson NS, et al. Recruitment of African-American pre-adolescent girls into an obesity prevention trial: the GEMS pilot studies - Project Account [Child of RefID 75039]. *Ethn Dis* 2003;13(1 Suppl 1):S78-S87.
7. Bellows LL, Davies PL, Anderson J, Kennedy C. Effectiveness of a physical activity intervention for Head Start preschoolers: a randomized intervention study. *Am J Occup Ther* 2013;67(1):28-36.
8. Black MM, Hager ER, Le K, Anliker J, Arteaga SS, Diclemente C, et al. Challenge! Health promotion/obesity prevention mentorship model among urban, black adolescents. *Pediatrics* 2010 Aug;126(2):280-8.
9. Bonsergent E, Agrinier N, Thilly N, Tessier S, Legrand K, Lecomte E, et al. Overweight and obesity prevention for adolescents: a cluster randomized controlled trial in a school setting. *Am J Prev Med* 2013;44(1):30-9.
10. Briancon S, Bonsergent E, Agrinier N, Tessier S, Legrand K, Lecomte E, et al. PRALIMAP: study protocol for a high school-based, factorial cluster randomised interventional trial of three overweight and obesity prevention strategies. *Trials* 2010;11:119.
11. Brandstetter S, Klenk J, Berg S, Galm C, Fritz M, Peter R, et al. Overweight prevention implemented by primary school teachers: a randomised controlled trial. *Obes Facts* 2012;5(1):1-11.

12. Brown B, Noonan C, Harris KJ, Parker M, Gaskill S, Ricci C, et al. Developing and piloting the Journey to Native Youth Health program in Northern Plains Indian communities. *Diabetes Educ* 2013;39(1):109-18.
13. Bürgi F, Niederer I, Schindler C, Bodenmann P, Marques-Vidal P, Kriemler S, et al. Effect of a lifestyle intervention on adiposity and fitness in socially disadvantaged subgroups of preschoolers: a cluster-randomized trial (Ballabeina). *Prev Med* 2012;54(5):335-40.
14. Puder JJ, Marques-Vidal P, Schindler C, Zahner L, Niederer I, Bürgi F, et al. Effect of multidimensional lifestyle intervention on fitness and adiposity in predominantly migrant preschool children (Ballabeina): Cluster randomised controlled trial. *BMJ (Online)* 2011;343:d6195.
15. Niederer I, Bürgi F, Ebenegger V, Marques-Vidal P, Schindler C, Nydegger A, et al. Effects of a lifestyle intervention on adiposity and fitness in overweight or low fit preschoolers (Ballabeina). *Obesity (Silver Spring)* 2013;21(3):E287-93.
16. Caballero B, Clay T, Davis SM, Ethelbah B, Rock BH, Lohman T, et al. Pathways: a school-based, randomized controlled trial for the prevention of obesity in American Indian schoolchildren. *Am J Clin Nutr* 2003 Nov;78(5):1030-8.
17. Campbell KJ, Lioret S, McNaughton SA, Crawford DA, Salmon J, Ball K, et al. A parent-focused intervention to reduce infant obesity risk behaviors: a randomized trial. *Pediatrics* 2013;131(4):652-60.
18. Campbell K, Hesketh K, Crawford D, Salmon J, Ball K, McCallum Z. The Infant Feeding Activity and Nutrition Trial (INFANT) an early intervention to prevent childhood obesity: cluster-randomised controlled trial. *BMC Public Health* 2008;8:103.
19. Crespo NC, Elder JP, Ayala GX, Slymen DJ, Campbell NR, Sallis JF, et al. Results of a multi-level intervention to prevent and control childhood obesity among Latino children: the Aventuras Para Niños Study. *Ann Behav Med* 2012;43(1):84-100.
20. Elder JP, Arredondo EM, Campbell N, Baquero B, Duerksen S, Ayala G, et al. Individual, family, and community environmental correlates of obesity in Latino elementary school children [Erratum in *J SCH HEALTH* 2010 Mar;80(3):159]. *J Sch Health* 2010;80(1):20-30.
21. Cunha DB, de Souza B, Pereira RA, Sichieri R. Effectiveness of a randomized school-based intervention involving families and teachers to prevent excessive weight gain among adolescents in Brazil. *PLoS ONE* 2013;8(2):e57498.
22. Daniels LA, Mallan KM, Battistutta D, Nicholson JM, Perry R, Magarey A. Evaluation of an intervention to promote protective infant feeding practices to prevent childhood obesity: outcomes of the NOURISH RCT at 14 months of age and 6 months post the first of two intervention modules. *Int J Obes (Lond)* 2012;36(10):1292-8.

23. DeBar LL, Schneider M, Drews KL, Ford EG, Stadler DD, Moe EL, et al. Student public commitment in a school-based diabetes prevention project: impact on physical health and health behavior. *BMC Public Health* 2011;11:711.
24. The HEALTHY Study Group. HEALTHY study rationale, design and methods: moderating risk of type 2 diabetes in multi-ethnic middle school students. *Int J Obes (Lond)* 2009;33(Suppl 4):S4-20.
25. de Heer HD, Koehly L, Pederson R, Morera O. Effectiveness and spillover of an after-school health promotion program for Hispanic elementary school children. *Am J Public Health* 2011 Oct;101(10):1907-13.
26. De Coen V, De Bourdeaudhuij I, Vereecken C, Verbestel V, Haerens L, Huybrechts I, et al. Effects of a 2-year healthy eating and physical activity intervention for 3-6-year-olds in communities of high and low socio-economic status: the POP (Prevention of Overweight among Pre-school and school children) project. *Public Health Nutr* 2012;15(9):1737-45.
27. de Ruyter JC, Olthof MR, Seidell JC, Katan MB. A trial of sugar-free or sugar-sweetened beverages and body weight in children. *N Engl J Med* 2012 Oct 11;367(15):1397-406.
28. de Ruyter JC, Olthof MR, Kuijper LDJ, Katan MB. Effect of sugar-sweetened beverages on body weight in children: design and baseline characteristics of the Double-blind, Randomized INtervention study in Kids. *Contemp Clin Trials* 2012;33(1):247-57.
29. Donnelly JE, Greene JL, Gibson CA, Smith BK, Washburn RA, Sullivan DK, et al. Physical Activity Across the Curriculum (PAAC): a randomized controlled trial to promote physical activity and diminish overweight and obesity in elementary school children. *Prev Med* 2009 Oct;49(4):336-41.
30. Gibson CA, Smith BK, DuBose KD, Greene JL, Bailey BW, Williams SL, et al. Physical activity across the curriculum: year one process evaluation results. *Int J Behav Nutr Phys Act* 2008;5:36.
31. Dziewaltowski DA, Rosenkranz RR, Geller KS, Coleman KJ, Welk GJ, Hastmann TJ, et al. HOP'N after-school project: an obesity prevention randomized controlled trial. *Int J Behav Nutr Phys Act* 2010;7:90.
32. Ebbeling CB, Feldman HA, Osganian SK, Chomitz VR, Ellenbogen SJ, Ludwig DS. Effects of decreasing sugar-sweetened beverage consumption on body weight in adolescents: a randomized, controlled pilot study. *Pediatrics* 2006 Mar;117(3):673-80.
33. El Ansari W, El Ashker S, Moseley L. Associations between physical activity and health parameters in adolescent pupils in Egypt. *Int J Environ Res Public Health* 2010 Apr;7(4):1649-69.



34. Escribano J, Luque V, Ferre N, Mendez-Riera G, Koletzko B, Grote V, et al. Effect of protein intake and weight gain velocity on body fat mass at 6 months of age: the EU Childhood Obesity Programme. *Int J Obes (Lond)* 2012;36(4):548-53.
35. Koletzko B, von Kries R, Closa R, Escribano J, Scaglioni S, Giovannini M, et al. Lower protein in infant formula is associated with lower weight up to age 2 y: a randomized clinical trial. *Am J Clin Nutr* 2009 Jun;89(6):1836-45.
36. Fitzgibbon ML, Stolley MR, Schiffer L, Van Horn L, KauferChristoffel K, Dyer A. Two-year follow-up results for Hip-Hop to Health Jr.: a randomized controlled trial for overweight prevention in preschool minority children. *J Pediatr* 2005 May;146(5):618-25.
37. Fitzgibbon ML, Stolley MR, Schiffer L, Van Horn L, KauferChristoffel K, Dyer A. Hip-Hop to Health Jr. for Latino preschool children. *Obesity (Silver Spring)* 2006 Sep;14(9):1616-25.
38. Fitzgibbon ML, Stolley MR, Schiffer LA, Braunschweig CL, Gomez SL, Van Horn L, et al. Hip-Hop to Health Jr. Obesity Prevention Effectiveness Trial: postintervention results. *Obesity (Silver Spring)* 2011 May;19(5):994-1003.
39. Fitzgibbon ML, Stolley MR, Schiffer L, Kong A, Braunschweig CL, Gomez-Perez SL, et al. Family-based hip-hop to health: outcome results. *Obesity (Silver Spring)* 2013;21(2):274-83.
40. Foster GD, Sherman S, Borradaile KE, Grundy KM, Vander Veur SS, Nachmani J, et al. A policy-based school intervention to prevent overweight and obesity. *Pediatrics* 2008 Apr;121(4):e794-802.
41. French SA, Gerlach AF, Mitchell NR, Hannan PJ, Welsh EM. Household obesity prevention: Take action--a group-randomized trial. *Obesity (Silver Spring)* 2011 Oct;19(10):2082-8.
42. HEALTHY study group, Foster GD, Linder B, Baranowski T, Cooper DM, Goldberg L, et al. A school-based intervention for diabetes risk reduction. *N Engl J Med* 2010 Jul 29;363(5):443-53.
43. Fung C, Kuhle S, Lu C, Purcell M, Schwartz M, Storey K, et al. From "best practice" to "next practice": the effectiveness of school-based health promotion in improving healthy eating and physical activity and preventing childhood obesity. *Int J Behav Nutr Phys Act* 2012;9:27.
44. Gentile DA, Welk G, Eisenmann JC, Reimer RA, Walsh DA, Russell DW, et al. Evaluation of a multiple ecological level child obesity prevention program: Switch what you Do, View, and Chew. *BMC Med* 2009;7:49.

45. Greening L, Harrell KT, Low AK, Fielder CE. Efficacy of a school-based childhood obesity intervention program in a rural southern community: TEAM Mississippi Project. *Obesity* (Silver Spring) 2011 Jun;19(6):1213-9.
46. Haerens L, Deforche B, Maes L, Cardon G, Stevens V, De B, I. Evaluation of a 2-year physical activity and healthy eating intervention in middle school children. *Health Educ Res* 2006 Dec;21(6):911-21.
47. Haerens L, Deforche B, Maes L, Stevens V, Cardon G, De B, I. Body mass effects of a physical activity and healthy food intervention in middle schools - Project Account [child of 75061]. *Obesity* (Silver Spring) 2006 May;14(5):847-54.
48. Hakanen M, Lagström H, Pahkala K, Sillanmaki L, Saarinen M, Niinikoski H, et al. Dietary and lifestyle counselling reduces the clustering of overweight-related cardiometabolic risk factors in adolescents. *Acta Paediatr* 2010 Jun;99(6):888-95.
49. Harvey-Berino J, Rourke J. Obesity prevention in preschool native-american children: a pilot study using home visiting. *Obes Res* 2003 May;11(5):606-11.
50. Hoffman JA, Thompson DR, Franko DL, Power TJ, Leff SS, Stallings VA. Decaying behavioral effects in a randomized, multi-year fruit and vegetable intake intervention. *Prev Med* 2011 May;52(5):370-5.
51. Howe CA, Harris RA, Gutin B. A 10-month physical activity intervention improves body composition in young black boys. *J Obes* 2011;2011:358581.
52. James J, Thomas P, Cavan D, Kerr D. Preventing childhood obesity by reducing consumption of carbonated drinks: cluster randomised controlled trial.[Erratum in *BMJ*. 2004 May 22;328(7450):1236.]. *BMJ* 2004 May 22;328(7450):1237.
53. James J, Thomas P, Kerr D. Preventing childhood obesity: two year follow-up results from the Christchurch obesity prevention programme in schools (CHOPPS). *BMJ* 2007;335:762.
54. Jansen W, Borsboom G, Meima A, Zwanenburg EJ, Mackenbach JP, Raat H, et al. Effectiveness of a primary school-based intervention to reduce overweight. *Int J Pediatr Obes* 2011 Jun;6(2-2):e70-7.
55. Katz DL, Katz CS, Treu JA, Reynolds J, Njike V, Walker J, et al. Teaching healthful food choices to elementary school students and their parents: the Nutrition Detectives™ program. *J Sch Health* 2011;81(1):21-8.
56. Klesges RC, Obarzanek E, Kumanyika S, Murray DM, Klesges LM, Relyea GE, et al. The Memphis Girls' health Enrichment Multi-site Studies (GEMS): an evaluation of the efficacy of a 2-year obesity prevention program in African American girls. *Arch Pediatr Adolesc Med* 2010 Nov;164(11):1007-14.

57. Kriemler S, Zahner L, Schindler C, Meyer U, Hartmann T, Hebestreit H, et al. Effect of school based physical activity programme (KISS) on fitness and adiposity in primary schoolchildren: cluster randomised controlled trial. *BMJ* 2010;340:c785.
58. Zahner L, Puder JJ, Roth R, Schmid M, Guldimann R, Puhse U, et al. A school-based physical activity program to improve health and fitness in children aged 6-13 years ("Kinder-Sportstudie KISS"): study design of a randomized controlled trial [ISRCTN15360785]. *BMC Public Health* 2006;6:147.
59. Lazaar N, Aucouturier J, Ratel S, Rance M, Meyer M, Duché P. Effect of physical activity intervention on body composition in young children: influence of body mass index status and gender. *Acta Paediatr* 2007 Sep;96(9):1315-20.
60. Li YP, Hu XQ, Schouten EG, Liu AL, Du SM, Li LZ, et al. Report on childhood obesity in China (8): effects and sustainability of physical activity intervention on body composition of Chinese youth. *Biomed Environ Sci* 2010 Jun;23(3):180-7.
61. Llargués E, Recasens A, Franco R, Nadal A, Vila M, Pérez MJ, et al. Medium-term evaluation of an educational intervention on dietary and physical exercise habits in schoolchildren: the Avall 2 study. *Endocrinol Nutr* 2012;59(5):288-95.
62. Llargues E, Franco R, Recasens A, Nadal A, Vila M, Perez MJ, et al. Assessment of a school-based intervention in eating habits and physical activity in school children: the AVall study. *J EPIDEMIOLOGICAL COMMUNITY HEALTH* 2011 Oct;65(10):896-901.
63. Lloyd JJ, Wyatt KM, Creanor S. Behavioural and weight status outcomes from an exploratory trial of the Healthy Lifestyles Programme (HeLP): a novel school-based obesity prevention programme. *BMJ Open* 2012;2(3):pii-e000390.
64. Lubans DR, Morgan PJ, Aguiar EJ, Callister R. Randomized controlled trial of the Physical Activity Leaders (PALs) program for adolescent boys from disadvantaged secondary schools. *Prev Med* 2011 Mar;52(3-4):239-46.
65. Lubans DR, Morgan PJ, Callister R. Potential moderators and mediators of intervention effects in an obesity prevention program for adolescent boys from disadvantaged schools. *J Sci Med Sport* 2012;15(6):519-25.
66. Morgan PJ, Saunders KL, Lubans DR. Improving physical self-perception in adolescent boys from disadvantaged schools: psychological outcomes from the Physical Activity Leaders randomized controlled trial. *Pediatr Obes* 2012;7(3):e27-32.
67. Lubans DR, Morgan PJ, Okely AD, Dewar D, Collins CE, Batterham M, et al. Preventing obesity among adolescent girls: one-year outcomes of the Nutrition and Enjoyable Activity for Teen Girls (NEAT Girls) Cluster Randomized Controlled Trial. *Arch Pediatr Adolesc Med* 2012;166(9):821-7.

68. Madsen K, Thompson H, Adkins A, Crawford Y. School-community partnerships: a cluster-randomized trial of an after-school soccer program. *JAMA Pediatr* 2013;167(4):321-6.
69. Magnusson KT, Hrafnkelsson H, Sigurgeirsson I, Johannsson E, Sveinsson T. Limited effects of a 2-year school-based physical activity intervention on body composition and cardiorespiratory fitness in 7-year-old children. *Health Educ Res* 2012;27(3):484-94.
70. Marcus C, Nyberg G, Nordenfelt A, Karpmyr M, Kowalski J, Ekelund U. A 4-year, cluster-randomized, controlled childhood obesity prevention study: STOPP. *Int J Obes (Lond)* 2009 Apr;33(4):408-17.
71. Martínez-Vizcaino V, Salcedo AF, Franquelo Gutiérrez R, Solera Martínez M, Sánchez López M, Serrano Martínez M, et al. Assessment of an after-school physical activity program to prevent obesity among 9- to 10-year-old children: a cluster randomized trial. *Int J Obes (Lond)* 2008 Jan;32(1):12-22.
72. Mihas C, Mariolis A, Manios Y, Naska A, Arapaki A, Mariolis-Sapsakos T, et al. Evaluation of a nutrition intervention in adolescents of an urban area in Greece: short- and long-term effects of the VYRONAS study. *Public Health Nutr* 2010 May;13(5):712-9.
73. Morgan PJ, Lubans DR, Callister R, Okely AD, Burrows TL, Fletcher R, et al. The 'Healthy Dads, Healthy Kids' randomized controlled trial: efficacy of a healthy lifestyle program for overweight fathers and their children. *Int J Obes (Lond)* 2011 Mar;35(3):436-47.
74. Mo-suwan L, Pongprapai S, Junjana C, Puetpaiboon A. Effects of a controlled trial of a school-based exercise program on the obesity indexes of preschool children. *Am J Clin Nutr* 1998 Nov;68(5):1006-11.
75. Muckelbauer R, Libuda L, Clausen K, Toschke A, Reinehr T, Kersting M. Promotion and provision of drinking water in schools for overweight prevention: randomized, controlled cluster trial. *Nutr Today* 2012 Jul 2;47(45):S27-34.
76. Muckelbauer R, Libuda L, Clausen K, Toschke AM, Reinehr T, Kersting M. Immigrational background affects the effectiveness of a school-based overweight prevention program promoting water consumption. *Obesity (Silver Spring)* 2010 Mar;18(3):528-34.
77. Nemet D, Geva D, Eliakim A. Health promotion intervention in low socioeconomic kindergarten children. *J Pediatr* 2011;158(5):796-801.
78. Nemet D, Geva D, Pantanowitz M, Igarria N, Meckel Y, Eliakim A. Health promotion intervention in Arab-Israeli kindergarten children. *J Pediatr Endocrinol Metab* 2011;24(11-12):1001-7.

79. Neumark-Sztainer D, Story M, Hannan PJ, Rex J. New Moves: a school-based obesity prevention program for adolescent girls. *Prev Med* 2003 Jul;37(1):41-51.
80. Neumark-Sztainer DR, Friend SE, Flattum CF, Hannan PJ, Story MT, Bauer KW, et al. New moves-preventing weight-related problems in adolescent girls a group-randomized study. *Am J Prev Med* 2010 Nov;39(5):421-32.
81. Østbye T, Krause KM, Stroom M, Lovelady CA, Evenson KR, Peterson BL, et al. Parent-focused change to prevent obesity in preschoolers: Results from the KAN-DO study. *Prev Med* 2012;55(3):188-95.
82. Ostbye T, Zucker NL, Krause KM, Lovelady CA, Evenson KR, Peterson BL, et al. Kids and adults now! Defeat Obesity (KAN-DO): rationale, design and baseline characteristics. *Contemp Clin Trials* 2011;32(3):461-9.
83. Paineau DL, Beaufils F, Boulier A, Cassuto DA, Chwalow J, Combris P, et al. Family dietary coaching to improve nutritional intakes and body weight control: a randomized controlled trial. *Arch Pediatr Adolesc Med* 2008 Jan;162(1):34-43.
84. Papadaki A, Linardakis M, Larsen TM, van Baak MA, Lindroos AK, Pfeiffer AF, et al. The effect of protein and glycemic index on children's body composition: the DiOGenes randomized study. *Pediatrics* 2010 Nov;126(5):e1143-52.
85. Larsen TM, Dalskov S, van BM, Jebb S, Kafatos A, Pfeiffer A, et al. The Diet, Obesity and Genes (Diogenes) Dietary Study in eight European countries - a comprehensive design for long-term intervention. *Obes Rev* 2010 Jan;11(1):76-91.
86. Peralta LR, Jones RA, Okely AD. Promoting healthy lifestyles among adolescent boys: the Fitness Improvement and Lifestyle Awareness Program RCT. *Prev Med* 2009 Jun;48(6):537-42.
87. Reed KE, Warburton DE, Macdonald HM, Naylor PJ, McKay HA. Action Schools! BC: a school-based physical activity intervention designed to decrease cardiovascular disease risk factors in children. *Prev Med* 2008 Jun;46(6):525-31.
88. Naylor PJ, Macdonald HM, Zebedee JA, Reed KE, McKay HA. Lessons learned from Action Schools! BC--an 'active school' model to promote physical activity in elementary schools. *J Sci Med Sport* 2006 Oct;9(5):413-23.
89. Naylor PJ, Macdonald HM, Reed KE, McKay HA. Action Schools! BC: a socioecological approach to modifying chronic disease risk factors in elementary school children. *Prev Chronic Dis* 2006 Apr;3(2):A60.
90. Reilly JJ, Kelly L, Montgomery C, Williamson A, Fisher A, McColl JH, et al. Physical activity to prevent obesity in young children: cluster randomised controlled trial. *BMJ* 2006 Nov 18;333(7577):1041.

91. Robinson TN, Killen JD, Kraemer HC, Wilson DM, Matheson DM, Haskell WL, et al. Dance and reducing television viewing to prevent weight gain in African-American girls: the Stanford GEMS pilot study. *Ethn Dis* 2003;13(Suppl 1):S65-77.
92. Rochon J, Klesges RC, Story M, Robinson TN, Baranowski T, Obarzanek E, et al. Common design elements of the Girls health Enrichment Multi-site Studies (GEMS) - Project Account [Child of RefID 75039]. *Ethn Dis* 2003;13(1 Suppl 1):S6-14.
93. Robinson TN, Matheson DM, Kraemer HC, Wilson DM, Obarzanek E, Thompson NS, et al. A randomized controlled trial of culturally tailored dance and reducing screen time to prevent weight gain in low-income African American girls: Stanford GEMS. *Arch Pediatr Adolesc Med* 2010 Nov;164(11):995-1004.
94. Rosário R, Araújo A, Oliveira B, Padrão P, Lopes O, Teixeira V, et al. Impact of an intervention through teachers to prevent consumption of low nutrition, energy-dense foods and beverages: a randomized trial. *Prev Med* 2013;57(1):20-5.
95. Rosário R, Oliveira B, Araújo A, Lopes O, Padrão P, Moreira A, et al. The impact of an intervention taught by trained teachers on childhood overweight. *Int J Environ Res Public Health* 2012;9(4):1355-67.
96. Rosenkranz RR, Behrens TK, Dzewaltowski DA. A group-randomized controlled trial for health promotion in Girl Scouts: healthier troops in a SNAP (Scouting Nutrition & Activity Program). *BMC Public Health* 2010;10:81.
97. Rush E, Reed P, McLennan S, Coppinger T, Simmons D, Graham D. A school-based obesity control programme: Project Energize. Two-year outcomes. *Br J Nutr* 2012;107(4):581-7.
98. Graham D, Appleton S, Rush E, McLennan S, Reed P, Simmons D. Increasing activity and improving nutrition through a schools-based programme: Project Energize. 1. Design, programme, randomisation and evaluation methodology. *Public Health Nutr* 2008 Oct;11(10):1076-84.
99. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* 2000 May 6;320(7244):1240-3.
100. Salcedo AF, Martínez-Vizcaíno V, Sánchez López M, Solera Martínez M, Franquelo Gutiérrez R, Serrano Martínez M, et al. Impact of an after-school physical activity program on obesity in children. *J Pediatr* 2010 Jul;157(1):36-42.
101. Shamah Levy T, Morales Ruán C, Amaya Castellanos C, Salazar Coronel A, Jiménez Aguilar A, Méndez Gómez Humarán I. Effectiveness of a diet and physical activity promotion strategy on the prevention of obesity in Mexican school children. *BMC Public Health* 2012;12:152.

102. Sichieri R, Paula TA, de Souza RA, Veiga GV. School randomised trial on prevention of excessive weight gain by discouraging students from drinking sodas. *Public Health Nutr* 2009 Feb;12(2):197-202.
103. Siegrist M, Lammel C, Haller B, Christle J, Halle M. Effects of a physical education program on physical activity, fitness, and health in children: the Juven TUM project. *Scan J Med Sci Sports* 2013 Jun;23(3):323-30.
104. Simon C, Schweitzer B, Oujaa M, Wagner A, Arveiler D, Triby E, et al. Successful overweight prevention in adolescents by increasing physical activity: a 4-year randomized controlled intervention.[Erratum in *Int J Obes (Lond)*. 2008 Oct;32(10):1606.]. *Int J Obes (Lond)* 2008 Oct;32(10):1489-98.
105. Simon C, Wagner A, DiVita C, Rauscher E, Klein-Platat C, Arveiler D, et al. Intervention centred on adolescents' physical activity and sedentary behaviour (ICAPS): concept and 6-month results - Project Account [Child of 75047]. *Int J Obes Relat Metab Disord* 2004 Nov;28(Suppl 3):S96-103.
106. Simon C, Wagner A, Platat C, Arveiler D, Schweitzer B, Schlienger JL, et al. ICAPS: a multilevel program to improve physical activity in adolescents. *Diabetes Metab* 2006 Feb;32(1):41-9.
107. Singh AS, Chin APM, Brug J, van Mechelen W. Dutch obesity intervention in teenagers: effectiveness of a school-based program on body composition and behavior. *Arch Pediatr Adolesc Med* 2009 Apr;163(4):309-17.
108. Singhal N, Misra A, Shah P, Gulati S. Effects of controlled school-based multi-component model of nutrition and lifestyle interventions on behavior modification, anthropometry and metabolic risk profile of urban Asian Indian adolescents in North India. *Eur J Clin Nutr* 2010 Apr;64(4):364-73.
109. Story M, Sherwood NE, Himes JH, Davis M, Jacobs DR, Jr., Cartwright Y, et al. An after-school obesity prevention program for African-American girls: the Minnesota GEMS pilot study. *Ethn Dis* 2003;13(1 Suppl 1):S54-64.
110. Story M, Hannan PJ, Fulkerson JA, Rock BH, Smyth M, Arcan C, et al. Bright start: description and main outcomes from a group-randomized obesity prevention trial in american Indian children. *Obesity (Silver Spring)* 2012;20(11):2241-9.
111. Telford RD, Cunningham RB, Fitzgerald R, Olive LS, Prosser L, Jiang X, et al. Physical education, obesity, and academic achievement: a 2-year longitudinal investigation of Australian elementary school children. *Am J Public Health* 2012;102(2):368-74.
112. Telford RD, Bass SL, Budge MM, Byrne DG, Carlson JS, Coles D, et al. The lifestyle of our kids (LOOK) project: outline of methods. *J Sci Med Sport* 2009 Jan;12(1):156-63.

113. Thivel D, Isacco L, Lazaar N, Aucouturier J, Ratel S, Dóre E, et al. Effect of a 6-month school-based physical activity program on body composition and physical fitness in lean and obese schoolchildren. *Eur J Pediatr* 2011;170(11):1435-43.
114. Velez A, Golem DL, Arent SM. The impact of a 12-week resistance training program on strength, body composition, and self-concept of Hispanic adolescents. *J Strength Cond Res* 2010 Apr;24(4):1065-73.
115. Webber LS, Catellier DJ, Lytle LA, Murray DM, Pratt CA, Young DR, et al. Promoting physical activity in middle school girls: Trial of Activity for Adolescent Girls. *Am J Prev Med* 2008 Mar;34(3):173-84.
116. Weeks BK, Beck BR. Twice-weekly, in-school jumping improves lean mass, particularly in adolescent boys. *Pediatr Obes* 2012;7(3):196-204.
117. Wen LM, Baur LA, Simpson JM, Rissel C, Wardle K, Flood VM. Effectiveness of home based early intervention on children's BMI at age 2: randomised controlled trial. *BMJ* 2012;344:e3732.
118. Williamson DA, Champagne CM, Harsha DW, Han H, Martin CK, Newton Jr, et al. Effect of an environmental school-based obesity prevention program on changes in body fat and body weight: a randomized trial. *Obesity (Silver Spring)* 2012;20(8):1653-61.
119. Williamson DA, Champagne CM, Harsha D, Han H, Martin CK, Newton R, Jr., et al. Louisiana (LA) Health: design and methods for a childhood obesity prevention program in rural schools. *Contemp Clin Trials* 2008 Sep;29(5):783-95.
120. Yin Z, Moore JB, Johnson MH, Vernon MM, Gutin B. The impact of a 3-year after-school obesity prevention program in elementary school children. *Child Obes* 2012;8(1):60-70.