

Table 1. Expression of R-cadherin in clinical breast cancer specimens

Case	Intraductal			Invasive			Cyto ⁴
	Cells (%) ¹	Intensity ²	Score ³	Cells (%)	Intensity	Score	
1	100	3	1.00	90	2	0.60	
2	95	3	0.95	90	2	0.60	
3				90	2	0.60	+
4				70	2	0.47	
5				50	2	0.33	
6	30	2	0.20	30	2	0.20	
7				60	2	0.40	
8				80	2	0.53	+
9				10	1	0.03	
10	100	3	1.00	50	2	0.33	+
11				70	2	0.47	
12	80	3	0.80	20	2	0.13	
13	95	3	0.95	60	3	0.60	
14				15	2	0.10	
15				80	2	0.53	+
16	70	2	0.47	30	2	0.20	
17				70	2	0.47	
18	80	2	0.53	60	2	0.40	
Avg	81	2.6	0.74	57	2.0	0.39	
SD	23.4	0.5	0.30	26.3	0.3	0.19	
p < ⁵				0.0172		0.0007	

¹ Per centage of positive tumor cells

² Average staining intensity - 0, 1, 2, or 3

³ Product of positive tumor cells X intensity, normalized to the maximal value of 300.

⁴ Cytoplasmic staining in a significant fraction of tumor cells. Case 15 had only cytoplasmic staining and three others had membrane and cytoplasmic staining.

⁵ Comparison Intraductal and Invasive Carcinoma; Student's T-test, one-tailed, equal variance