

SUPPLEMENTARY TABLE S2. HAPLOTYPE DISTRIBUTION BETWEEN PERIODONTITIS PATIENTS AND CONTROLS

<i>Haplotype</i>	<i>Control</i>	<i>Case</i>	<i>OR (95% CI)</i>	<i>p-Value</i>
TGCC	125 (31.3)	123 (30.8)	Reference	
TGTC	74 (18.4)	74 (18.4)	1.02 (0.68–1.52)	0.938
CTCC	55 (13.8)	28 (7.1)	0.52 (0.31–0.87)	0.012
TGCT	47 (11.9)	32 (8.1)	0.69 (0.41–1.16)	0.158
CTTC	21 (5.4)	25 (6.3)	1.21 (0.64–2.27)	0.554
TTCC	20 (5.1)	15 (3.8)	0.76 (0.37–1.56)	0.455
CGCC	11 (2.7)	8 (2.1)	0.74 (0.29–1.90)	0.529
TTCT	10 (2.6)	14 (3.5)	1.42 (0.61–3.32)	0.414
TTTC	10 (2.6)	13 (3.4)	1.32 (0.56–3.13)	0.525
CGTC	9 (2.3)	11 (2.6)	1.24 (0.50–3.10)	0.642
CGCT	7 (1.7)	8 (1.9)	1.16 (0.41–3.30)	0.780
CTTT	4 (0.9)	8 (2.1)	2.03 (0.60–6.92)	0.248
TGTT	4 (0.9)	15 (3.7)	3.18 (1.23–11.8)	0.014
TTTT	2 (0.5)	0 (0.0)	—	—
CTCT	0 (0.0)	22 (5.6)	—	—
CGTT	0 (0.0)	2 (0.6)	—	—

Haplotype was constructed using SNPs of *IL1A* and *IL1B* gene polymorphisms in the following order rs1800587, rs17561, rs16944, and rs1143634.

p-Value < 0.05, statistically significant.