## Appendix 1: Web references

[Web reference 1, Appendix 1] Blythe MJ, Fortenberry JD and Orr DP. Douching behaviours reported by adolescent and young adult women at high risk for sexually transmitted infections. *Journal of Paediatric and Adolescent Gynaecology* 2003;**16**(2):95-100.

[Web reference 2, Appendix 1] Brotman RM, Bradford LL, Gajer P et al. Assocoiation between Trichomonas vaginalis and vaginal bacterial community composition among reporductive-age women. *Sex Transm Dis* 2012;**39**(10):807-12.

[Web reference 3, Appendix 1] Nathan B, Appiah J, Heron D et al. Evaluation of five different tests for Trichomonas vaginalis and cost effective planning for clinical implementation. Poster presented at the British Association for Sexual Health and HIV Spring meeting. May 2013. Abstract available at *Int J STD AIDS* May 2013 24: 1-60, doi:10.1177/0956462413484421

[Web reference 4, Appendix 1] Muir P. Economic Evaluation of the Aptima TV TMA. 2013. Available at http://public.ukcrn.org.uk/search/StudyDetail.aspx?StudyID=13827 (accessed on 20<sup>th</sup> June 2013).

[Web reference 5, Appendix 1] Chapin KC, Andrea SB, Florczyk et al. Comparison of Transcription Mediated Amplification (TMA) and Polymerase Chain Reaction (PCR) methods for detection of *Trichomonas vaginalis* from SUN study patients [abstract]. Natl. STD Prev. Conf. Centers for Disease Control and Prevention, Atlanta, GA 2010:P75.