

Supplemental Table 18. Studies reporting the incidence of congestive heart failure in multiple sclerosis

Study (Year)	Region/ Sub- Region	Source/ Population	Prevalence Day/Perio d	Diagnostic Criteria: MS/ Established By	Diagnostic Criteria: CHF/ Established By	Crude Overall Incidence (95%CI)	Age- Standardized Overall Incidence (95%CI)	Crude Incidence (95%CI)		Standardized Incidence (95%CI)		Quality Score
								Males	Females	Males	Females	
Christiansen ⁵ (2010)	Denmark	DNRP, incident (hospitalized) MS	1977-2006	ICD-10 G35.9, ICD-8 340 (prior to 1993)	ICD-8: 427.09, 427.10, 427.11, 427.19 ICD-10: I50, I11.0, I13.0, I13.2	1 year follow-up: 23/13,963 = 0.16% 2-30 years follow-up: 376/13,963 = 2.69%	NR	NR	NR	NR	NR	6/8
Jadidi ²⁸ (2013)	Sweden	M&H Co Database, incident MS	1987-2009	ICD-8: 340; ICD-9: 340; ICD-10: G35	ICD-8: 427.0, 427.1; ICD-9: 428; ICD-10: I50	68/7,664 = 0.89%	NR	26/5215 = 0.50%	49/2449 = 2.00%	NR	NR	5/8

NR: Not Reported, DNRP- Danish National Registry of Patients, M&H Co Database: Migration and Health Cohort Database (contains over 15 linkable Swedish national registers).