

S5 Table. Biodistribution of rFVIIIIFc as calculated %ID/organ as determined by QWBA or scintillation counting in FVIII/VWF DKO Mice

FVIII/VWF DKO mice	rFVIIIIFc by QWBA (%ID/organ)					rFVIIIIFc by Scintillation Counts (%ID/organ)	
	Organ	5 min	15 min	1 hr	2 hr	6 hr	5 min
Blood	28.42	19.33	11.45	8.17	3.24	41.89±1.18	33.15±1.12
Liver	29.23	30.45	13.55	8.35	2.92	25.49±1.55	25.22±5.28
Kidney	1.63	2.32	2.01	1.56	0.89	2.65±0.49	2.52±0.06
Lung	1.85	0.88	0.57	0.40	0.17	0.28±0.09	0.24±0.07
Muscle	3.59	2.57	1.73	2.57	1.57	3.91±0.77	3.4±0.21
Spleen	0.64	0.68	0.43	0.24	0.09	0.74±0.14	0.65±0.2
Heart	0.61	0.58	0.25	0.17	0.11	0.37±0.05	0.25±0.02
Total (%ID)	65.97	56.81	29.99	21.46	8.99	75.5±4.32	65.46±7.01
Urine						2.06±2.84	8.61±0.86