

S6 Table. Percentage of injected dose per gram of tissue (%ID/g) of rFVIII_hFc as determined by QWBA or scintillation counting in FVIII/VWF DKO Mice

FVIII/VWF DKO mice	rFVIII _h Fc by QWBA (%ID/g)					rFVIII _h Fc by Scintillation Counts (%ID/g)	
	5 min	15 min	1 hr	2 hr	16 hr	5 min	15 min
Blood	22.98	9.18	5.91	4.04	1.51	19.9±0.9	16.5±0.7
Liver	29.50	18.04	8.73	5.16	1.70	15.1±0.7	15.7±3.4
Kidney	7.02	5.87	5.55	4.11	2.22	6.7±1.4	6.7±0.1
Lung	21.35	5.94	4.18	2.83	1.13	1.9±0.6	1.7±0.5
Muscle	0.58	0.24	0.18	0.25	0.15	0.4±0.1	0.3±0
Spleen	9.93	6.14	4.20	2.24	0.81	6.7±1.4	6.2±1.9
Heart	7.30	6.95	5.87	0.63	0.56	3.3±0.5	2.3±0.2
Bile	5.48	3.56	18.06	18.99	14.37	ND	ND
Cecum	0.54	0.19	0.14	3.90	4.48	ND	ND
Large intestine	ND	0.52	0.30	1.43	8.77	0.4±0	0.4±0
Urine	55.24	50.20	3.55	99.22	4.84	43.3±53.8	168.4±123.9