

Online Table 5: Amblyopic-Eye Visual Acuity by Treatment Group

Amblyopic Eye	4-week Visit				10-week Visit				16-week Visit				18-week Primary Outcome Visit			
	Levodopa N=88		Placebo N=47		Levodopa N=88		Placebo N=48		Levodopa N=87		Placebo N=46		Levodopa N=86*		Placebo N=45	
	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%
Acuity Score																
20/25 (78-82 letters)	0	0	0	0	0	0	0	0	1	1%	0	0	0	0	0	0
20/32 (73-77 letters)	1	1%	1	2%	3	3%	0	0	2	2%	0	0	5	6%	1	2%
20/40 (68-72 letters)	8	9%	2	4%	12	14%	6	13%	13	15%	6	13%	7	8%	6	13%
20/50 (63-67 letters)	15	17%	10	21%	12	14%	11	23%	13	15%	10	22%	20	23%	5	11%
20/63 (58-62 letters)	15	17%	6	13%	14	16%	7	15%	17	20%	6	13%	18	21%	9	20%
20/80 (53-57 letters)	13	15%	11	23%	19	22%	5	10%	16	18%	7	15%	12	14%	5	11%
20/100 (48-52 letters)	19	22%	4	9%	16	18%	7	15%	16	18%	7	15%	17	20%	9	20%
20/125 (43-47 letters)	8	9%	3	6%	4	5%	5	10%	4	5%	3	7%	4	5%	3	7%
20/160 or worse (≤ 42)	9	10%	10	21%	8	9%	7	15%	5	6%	7	15%	3	3%	7	16%
Mean (SD) letter score	55.0 (10.9)		53.7 (12.0)		56.5 (10.5)		55.1 (12.0)		58.2 (9.6)		55.5 (11.8)		58.7 (8.9)		54.8 (12.3)	
	~20/80		~20/80 ⁻¹		~20/80 ⁺²		~20/80		~20/63 ⁻²		~20/80 ⁺¹		~20/63 ⁻¹		~20/80	

SD = standard deviation; CI = confidence interval

*Note: one participant completing the 18-week visit 288 days (41 weeks) post randomization is not included in the analysis as pre-specified in the analysis plan.