Online Table 7 – Fellow-Eye Visual Acuity by Treatment Group

| Fellow Eye | 18-week Primary Outcome | | | | 26-week Outcome* | | | | |
|-----------------------|----------------------------|-----|----------------------|-----|----------------------|-----|----------------------|-----|------------------|
| | | | | | | | | | Levodopa N=87 |
| | N | % | N | % | N | % | N | % | |
| | Acuity Score | | | | | | | | |
| | 20/12 (93-97 letters) | 12 | 14% | 4 | 9% | 15 | 17% | 7 | 16% |
| 20/16 (88-92 letters) | 44 | 51% | 22 | 49% | 34 | 40% | 22 | 49% | |
| 20/20 (83-87 letters) | 22 | 25% | 16 | 36% | 28 | 33% | 13 | 29% | |
| 20/25 (78-82 letters) | 8 | 9% | 3 | 7% | 9 | 10% | 2 | 4% | |
| 20/32 (73-77 letters) | 1 | 1% | 0 | 0 | 0 | 0 | 1 | 2% | |
| Mean (SD) | 88.1 (4.4) | | 88.0 (3.2) | | 88.0 (4.5) | | 88.2 (4.1) | | |
| | ~20/16 ⁻² | | ~20/16 ⁻² | | ~20/16 ⁻² | | ~20/16 ⁻² | | |
| Change from Baseline | | | | | | | | | |
| ≥15 letters worse | 0 | 0 | 0 | 0 | 1 | 1% | 0 | 0 | |
| 10-14 letters worse | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 5-9 letters worse | 4 | 5% | 3 | 7% | 2 | 2% | 4 | 9% | |
| within 4 letters (±4) | 68 | 78% | 39 | 87% | 74 | 86% | 35 | 78% | |
| 5-9 letters better | 15 | 17% | 3 | 7% | 7 | 8% | 6 | 13% | |
| 10-14 letters better | 0 | 0 | 0 | 0 | 2 | 2% | 0 | 0 | |
| Mean (SD) letters | 1.5 (3.0) | | 0.5 (3.1) | | 1.4 (3.6) | | 0.8 (3.3) | | |

SD = standard deviation

^{*} All participants continued in the study regardless of whether they continued with study medication after the 18-week visit; 49 (56%) participants in the levodopa group and 24 (53%) participants in the placebo group continued study medication at the 18-week visit.